

Review Article

GABAergic inhibitory neurons as therapeutic targets for cognitive impairment in schizophrenia

Meng-yi XU¹, Albert H C WONG^{2, *}

¹Departments of Pharmacology, University of Toronto; Centre for Addiction and Mental Health, Toronto, Ontario, Canada; ²Departments of Psychiatry, Pharmacology and the Institute of Medical Science, University of Toronto, and the Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, Ontario, Canada

Abstract

Schizophrenia is considered primarily as a cognitive disorder. However, functional outcomes in schizophrenia are limited by the lack of effective pharmacological and psychosocial interventions for cognitive impairment. GABA (gamma-aminobutyric acid) interneurons are the main inhibitory neurons in the central nervous system (CNS), and they play a critical role in a variety of pathophysiological processes including modulation of cortical and hippocampal neural circuitry and activity, cognitive function-related neural oscillations (eg, gamma oscillations) and information integration and processing. Dysfunctional GABA interneuron activity can disrupt the excitatory/inhibitory (E/I) balance in the cortex, which could represent a core pathophysiological mechanism underlying cognitive dysfunction in schizophrenia. Recent research suggests that selective modulation of the GABAergic system is a promising intervention for the treatment of schizophrenia-associated cognitive defects. In this review, we summarized evidence from postmortem and animal studies for abnormal GABAergic neurotransmission in schizophrenia, and how altered GABA interneurons could disrupt neuronal oscillations. Next, we systemically reviewed a variety of up-to-date subtype-selective agonists, antagonists, positive and negative allosteric modulators (including dual allosteric modulators) for $\alpha 5/\alpha 3/\alpha 2$ GABA_A and GABA_B receptors, and summarized their pro-cognitive effects in animal behavioral tests and clinical trials. Finally, we also discuss various representative histone deacetylases (HDAC) inhibitors that target GABA system through epigenetic modulations, GABA prodrug and presynaptic GABA transporter inhibitors. This review provides important information on current potential GABA-associated therapies and future insights for development of more effective treatments.

Key words: schizophrenia; cognitive impairment; GABA; GABA prodrug; HDAC inhibitor; positive allosteric modulator; negative allosteric modulator; GAT-1 inhibitor; GABA_B receptor antagonists

Acta Pharmacologica Sinica (2018) 39: 733–753; doi: 10.1038/aps.2017.172; published online 22 Mar 2018

Introduction

Schizophrenia is a devastating psychiatric disease that affects approximately 1% of the population worldwide. This disorder is characterized by a heterogenous combination of symptoms that can be divided into positive symptoms including delusions and hallucinations, negative symptoms of impaired motivation, social withdrawal and affective flattening, and cognitive deficits such as impairments in attention, reasoning, processing speed as well as verbal and working memory^[1, 2]. The onset of psychosis typically occurs in late adolescence or early adulthood, but there are rare cases in which symptoms emerge during childhood or old age. A decline in cognitive

ability or prodrome often precedes the onset of the first psychotic episode^[3]. Cognitive symptoms persist throughout the course of the illness, and are the most important factors in long-term functional outcomes in schizophrenia^[1, 2, 4, 5].

Although schizophrenia is typically classified as a psychotic disorder^[1, 5], some suggest that it should be “considered primarily and foremost as a cognitive disorder”^[1]. Current pharmaceutical treatments are partially effective in reducing positive symptoms, but not for negative and cognitive symptoms. Therefore, functional outcomes in schizophrenia are limited by the lack of effective pharmacological and psychosocial interventions for cognitive impairment^[1, 2, 6].

γ -Aminobutyric acid (GABA) interneurons are the main inhibitory neurons in the central nervous system (CNS), and they play a critical role in a variety of physiological processes including modulation of cortical and hippocampal neural circuitry and activity^[7, 8], cognitive function-related neural oscilla-

*To whom correspondence should be addressed.

E-mail albert.wong@utoronto.ca

Received 2017-10-10 Accepted 2017-12-25

tions (eg gamma oscillations)^[9] and information integration and processing^[10]. Multiple lines of evidence strongly support that the GABAergic system is a major convergence point for both genetic and environmental risk factors of schizophrenia^[11]. For instance, a recent large genome-wide association study (GWAS) involving 11 355 schizophrenia patients and 16 416 controls identified copy number variations (CNVs) enriched in genes related to GABA neurotransmission^[12]. Duplications were strongly enriched for components of GABA_A receptor complexes, where the most highly associated genes were $\alpha 5$, $\beta 3$, and δ receptor subunits. It is known that excessive $\alpha 5$ -GABA receptor signaling can contribute to cognitive impairment of schizophrenia (discussed below), thereby providing strong support for the view that tonic inhibition mediated by $\alpha 5$ and δ subunits-containing GABA_A receptors could be disrupted in schizophrenia^[12]. Mutations in schizophrenia-linked genes such as *DISC1*^[13], *NRG1* and *ERBB4*^[14] can lead to disrupted GABA interneuron development, perturbed GABA circuitry and impaired synchrony of neural oscillations. Environmental factors such as prenatal infection and hypoxia, stress, smoking and cannabis use, when interacting with risk genes, can also contribute to schizophrenia^[15, 16]. Dysfunctional GABA interneuron activity can disrupt the excitatory/inhibitory (E/I) balance in the cortex^[17], which could be a core pathophysiological mechanism underlying cognitive dysfunction in schizophrenia. In this review, we will discuss evidence from postmortem and animal studies for abnormal GABA neurotransmission in schizophrenia, and how altered GABA interneurons could disrupt neuronal oscillations underlying cognitive impairment in schizophrenia. We will also discuss potential therapeutic targets and pharmacological treatments for cognitive deficits, with a focus on selective GABA_A and GABA_B receptor modulators, epigenetic modulations, GABA prodrug and presynaptic GABA transporter inhibitors.

GABA system and cognitive dysfunction in schizophrenia

Reductions in parvalbumin and GAD67 expression are associated with alterations in GABA transmission in schizophrenia

Several markers of GABA neurotransmission are altered in cortical regions of patients with schizophrenia. The most conserved and consistent finding from multiple imaging, animal and postmortem studies is the reduction in mRNA and protein levels for the 67 kDa isoform of glutamic acid decarboxylase (GAD67). This enzyme is responsible for synthesizing the majority of cytosolic and vesicular GABA^[18] in the dorsal lateral prefrontal cortex (DLPFC), which is responsible for working memory and selective attention^[19–21]. Decreased GAD67 mRNA is selectively observed in a subpopulation of prefrontal cortex (PFC) GABA neurons that express the calcium-binding protein parvalbumin (PV)^[19, 22]. Around 45% of PV-mRNA positive neurons have undetectable levels of GAD67 mRNA and such alteration is due to reduction in PV mRNA in neurons rather than a decreased density of PV neurons. By contrast, approximately 10% of PV neurons do not express GAD67 mRNA in healthy controls^[22–24]. Furthermore, since the reduction in PV mRNA is also found in non-medicated

schizophrenia patients and PV mRNA is not altered in the PFC of primates treated long-term with antipsychotics^[22–24], alterations in PV mRNA are unlikely to be due to antipsychotic drugs.

Parvalbumin functions as a slow calcium buffer, and can influence GABA release. Genetic elimination of this protein leads to various alterations in GABA neurotransmission. Taken together, these findings support the hypothesis that the capacity of PV neurons to synthesize and release GABA is impaired in the cortex of patients with schizophrenia^[25]. The reduction in GAD67 mRNA, however, does not necessarily mean that the GABA concentration is decreased in schizophrenia, because the reduced GAD67 mRNA may also be associated with slower GABA metabolism^[26]. Interestingly, a recent study found that the cerebrospinal fluid concentration of GABA analyzed by high-performance liquid chromatography was significantly reduced in first-episode psychosis patients compared with healthy controls, and patients with low GABA concentrations tend to have poor attention. Therefore, this study provides clinical evidence for a potential role of GABA in an early-stage schizophrenia^[27].

GABA inhibitory circuit deficits lead to impaired neural oscillations in schizophrenia

Many important brain functions depend on the coordinated activity of large populations of neurons within one region or across brain regions. Neural oscillations in the gamma frequency range (30–80 Hz) in the PFC have been studied extensively because of their strong relationship with complex cognitive functions, and because disruption of gamma oscillations could be an important mechanism underlying cognitive deficits in schizophrenia^[28]. Functionally, gamma-band oscillations in the human PFC increase in proportion to working memory load^[29]. Compared to healthy controls, schizophrenia patients have a marked decrease in the amplitude and phase synchronization of gamma oscillations in the frontal cortex, and they tend to perform poorly on executive and working memory tasks^[28, 30]. These deficits in gamma oscillations are observed in schizophrenia patients, independently of antipsychotic medication treatment^[30, 31].

In the hippocampus, gamma-band oscillations originating in the CA3 and CA1 regulate network activity that promotes the encoding of spatial information and formation of episodic memories^[32]. Slow theta-frequency oscillations (4–8 Hz) are complementary to gamma oscillations, and are especially important for episodic memory formation. Both gamma and theta oscillations are observed independently in the cortex and hippocampus but they are also coupled to each other^[9]. Aberrant theta-gamma coupling can affect cognitive function in schizophrenia, such as visuospatial working memory^[33]. In addition, ketamine, a pharmacological agent often to create animal models of schizophrenia, has been shown to alter gamma-theta oscillatory coupling in the hippocampus^[34]. Taken together, these findings support the view that disrupted functional connectivity of cortical and hippocampal neural networks is a core mechanism underlying cognitive impair-

ment in schizophrenia^[31].

It has been suggested that GABAergic inhibitory circuits play a crucial role for the generation of gamma oscillations and synchrony^[28]. Despite the presence of many cortical interneuron subtypes, only some contribute to the generation of gamma oscillations. Several lines of evidence have linked disturbances in the subset of perisomatic-targeting, fast-spiking, PV-expressing (FS PV) neurons to impaired gamma oscillations and working memory deficits in schizophrenia. The synchronous activity of FS PV neurons can generate gamma oscillations in mice *in vivo*. Optogenetic stimulation^[35,36] and inhibition^[35] of FS PV neurons selectively enhances and suppresses gamma oscillations *in vivo*, respectively. A subset of the FS PV interneurons, namely the parvalbumin-containing basket neurons (PVBC) that synapse on the somas of target pyramidal neurons and mediate fast, strong and shunting inhibition, are primarily responsible for gamma oscillations^[37]. Networks of these basket neurons connected by gap junctions have been shown to produce large synchronous inhibitory postsynaptic potentials (IPSPs) to pyramidal neurons, thus exerting precise inhibitory control over the temporal coding of information in pyramidal neurons^[25,37]. PV interneurons are also important for intrinsic theta rhythm generation in the hippocampus^[38]. Detailed neuronal mechanisms by which PV neurons mediate neural oscillations have been reviewed elsewhere^[25,26,28,37,39-42].

The reduced expression of GAD67 and PV in schizophrenia is accompanied by disinhibition of cortical excitatory neurons and diminished neuronal oscillation and synchrony. For example, GAD67^[26] and PV^[43] protein levels are lower in PVBC boutons in DLPFC of human postmortem brain tissues^[43], and downregulation of these two proteins in PV neurons probably reduces gamma oscillatory activity^[44] in PFC. In addition, by conditionally knocking out one allele of the *Gad1* gene (the gene encoding for GAD67) in PV neurons in rats, Lazarus *et al* found that the decrease in GAD67 mRNA reduced PV-neuron synaptic output^[45]. This in turn, disinhibited local pyramidal neurons in the DLPFC^[45]. Other subtypes of interneurons such as parvalbumin-expressing chandelier cells (PVChCs) and cholecystokinin-expressing (CCK) interneurons also promote gamma synchronization *in vivo*^[46]. As a whole, these findings suggest that reduced PV neuron output could result from decreased GABA synthesis and calcium buffering. This could be the cause of impaired gamma oscillations and working memory deficits in schizophrenia. Restoring the activity of GAD67 and PV within PV neurons, in particular PVBC, could be a promising strategy for improving cognitive deficits in schizophrenia^[43].

Furthermore, interneurons that innervate the perisomatic regions of the targeted pyramidal neurons such as axo-somatic PV basket neurons, axo-axonic PV chandelier neurons and CCK-containing neurons, also appear to be essential for generating and maintaining the fast oscillations (*eg*, gamma oscillations) and slow theta-frequency oscillations in the hippocampus, which play a critical role in different aspects of episodic memory^[47].

Alterations in other GABA genes related to GABA neurotransmission

In addition to GAD67 and PV, the expression of other key proteins in GABAergic pathways such as GABA transporter type 1 (GAT-1), REELIN (which is encoded by the gene *RELN* in humans), NMDA receptor subunits (*eg*, NR2A, NR3A), nicotine acetylcholine receptor (nAChR) $\alpha 4$ and $\alpha 7$ subunits, and brain-derived neurotrophic factor (BDNF)^[48] are reduced in the brains of schizophrenia patients^[49,50].

Reelin is essential for regulating the growth, maturation, synaptic plasticity^[51] and positioning of interneurons in the developing and adult brain^[52], and hence plays a crucial role in the pathophysiology of schizophrenia^[51]. The decreased number of dendritic spines observed in postmortem brains of schizophrenia patient is likely due to a deficit in reelin^[51]. At the molecular level, reelin can enhance GABA inhibitory signals through inhibition of GAT-1 internalization and increase KCC2 (potassium chloride cotransporter 2) expression^[53]. Hypermethylation of the *RELN* promoter can result in silencing of reelin^[54], with reduced transcription^[53,54].

As molecular alterations of these proteins have been implicated in the pathophysiology of prefrontal cortex dysfunction, they are potential targets for novel pharmacological interventions.

The GABA_A receptor as a therapeutic target in schizophrenia

The GABA_A receptor is a heteropentameric ligand-gated chloride channel, widely distributed in the mammalian CNS, that mediates synaptic and extrasynaptic inhibition. These receptors are also the site of action of a number of clinically important drugs, including benzodiazepines (BZs), barbiturates, and anesthetics^[55]. GABA_ARs consist of five different subunits, composed of 19 known subtypes ($\alpha 1-6$, $\beta 1-3$, $\gamma 1-3$, δ , ϵ , π , θ and $\rho 1-3$), although three subunits ($\rho 1-3$) are also thought to form the GABA_C receptor^[56]. Figure 1(A) depicts the structure and GABA/benzodiazepine binding sites of GABA_A receptors, whereas (B) depicts phasic *vs* tonic inhibition of the postsynaptic membrane mediated by synaptic *vs* extrasynaptic GABA_A receptors, respectively.

Benzodiazepine-like ligands for the treatment of cognitive defects in schizophrenia and other neuropsychiatric disorders

In general, the sedative and addictive effects of classical non-selective GABA_A receptor benzodiazepines have limited their usefulness in treating psychosis and cognitive impairments in schizophrenia. Consequently, a GABA_A receptor subtype-selective compound could overcome these limitations of the classical BZDs without unwanted side effects. Compared to full agonists or antagonists, allosteric modulators like BZDs that possess selective affinity and/or efficacy for different GABA_A receptor subtypes have been widely used for various purposes in schizophrenia^[67]. Despite some positive reports, there is no consistent evidence of efficacy of BZDs as adjunctive treatment for positive symptoms in schizophrenia^[67]. Convincing evidence for long-term cognitive benefits is also lacking^[56].




Figure 1. (A) The GABA_A receptor generally contains two α , two β and one γ subunit that are arranged in a $\alpha\beta\alpha\beta\gamma$ fashion, of which the γ subunit may be replaced by either an δ , ϵ or θ subunit, and the β subunit may be replaced by $\alpha^{[57]}$. Benzodiazepines enhance the action of the neurotransmitter GABA at GABA_A receptors by interaction with their allosteric modulatory benzodiazepine binding site (BZ site) that is formed by one of the α subunits ($\alpha 1$ -3 and $\alpha 5$) and $\gamma 2$ subunit^[56]. The BZ site is distinct from the endogenous ligand GABA binding site that occurs at the interface of the α and β subunits^[56, 58]. The α subunit is of importance in determining the pharmacological properties of the benzodiazepine drugs. (B) GABA-induced chloride ion influx hyperpolarizes postsynaptic neurons, generating inhibitory postsynaptic potentials (IPSPs)^[56]. GABA_A receptors mediate two distinct effects on the postsynaptic membrane: synaptic GABA_A receptors containing the $\alpha 1$ -, $\alpha 2$ - and $\alpha 3$ - subunits and the ubiquitous $\gamma 2$ subunit^[59] have been shown to possess a low affinity for GABA and mediate fast but short-lasting phasic inhibition, whereas extrasynaptic GABA_A receptors with a high-affinity for GABA that contain the relatively rare subunits $\alpha 4$, $\alpha 5$, $\alpha 6$, and δ mediate slow but long-lasting tonic inhibition^[56, 59, 60, 61, 62]. The function of phasic inhibition is critically dependent on the synaptic location of the subunit and the IPSP timing. One crucial role of phasic inhibition is to provide timing-based signaling for setting the temporal window of neuron networks firing^[63], therefore it is important for the generation and regulation of gamma (γ) or theta (θ) oscillations and synchrony^[64]. In contrast, tonic inhibition is responsible for generating about 75% of the total inhibitory conductance received by hippocampal neurons^[65] and regulates neuronal excitability through its effects on the magnitude and duration of the postsynaptic excitatory postsynaptic potential (EPSP)^[66].

GABA_A receptors containing the $\alpha 1$ subunit are the most prevalent BZDs-sensitive GABA_A receptors in the brain^[68]. There is insufficient evidence that selective allosteric agonists at $\alpha 1$ subunit-containing GABA_A receptors could improve cognition in schizophrenia^[68]. Two such compounds, triazolam and zolpidem, impair cognition in adult rhesus monkeys in the object retrieval with detours (ORD) task of executive function^[69]. Furthermore, GABA_A receptors containing $\alpha 3$ subunits inhibit the dopamine system^[70], and reduction in $\alpha 3$ subunit expression could contribute to a hyperdopaminergic state in schizophrenia^[71]. Partial positive agonists at $\alpha 3$ subunits (eg, ELB139) could thus be potential antipsychotics^[70].

Reduction in $\alpha 5$ -GABA_A receptor signaling might indirectly contribute to increased dopamine signaling in schizophrenia^[72, 73]. Therefore, selective allosteric $\alpha 5$ subunit activators could also have potential as antipsychotic medications. Multiple lines of evidence implicate GABA_A receptors containing $\alpha 2$ and/or $\alpha 3$ and $\alpha 5$ subunits in cognition. The following sections will focus on these two GABA_A subtypes and how selective modulators at these subtypes can improve cognition in

neuropsychiatric disorders such as schizophrenia, Alzheimer's disease and Down syndrome.

Role of $\alpha 5$ GABA_A receptor in cognitive function

The $\alpha 5$ subtype constitutes 5%-10% of total brain GABA_A receptors, but 25% in the hippocampus^[55]. The $\alpha 5$ -GABA_A receptor is mainly localized in the dendritic regions of the CA1-CA3 hippocampal areas^[55, 74], where they can modulate excitatory glutamatergic input^[75]. $\alpha 5$ subunits are also expressed in olfactory bulb, amygdala, hypothalamus and neocortex (layer V and VI) but to a lesser extent^[55, 74].

Studies in rodents have confirmed the role of $\alpha 5$ -GABA_A receptors in cognition through a variety of genetic and pharmacological approaches^[55, 59, 76]. For example, mice deficient for $\alpha 5$ subunits ($\alpha 5^{-/-}$) had better spatial learning in the water maze^[55]. Moreover, $\alpha 5$ (H105R) mutant mice with reduced $\alpha 5$ GABA_A receptors on hippocampal pyramidal neurons had stronger trace fear conditioning, but not in hippocampal-independent delay or contextual fear conditioning^[59].

The $\alpha 5$ GABA_A receptor subunit mediates tonic inhibition

in hippocampal neurons^[77], and regulated gamma oscillations^[77, 78]. Genetic knock-down or pharmacological inhibition, negative modulation of $\alpha 5$ -GABA_A receptors promotes hippocampal gamma oscillations^[77], long-term potentiation (LTP), and learning^[79]. For example, genetic reduction of $\alpha 5$ and δ subunits blocks tonic inhibition in CA3 pyramidal neurons, and produces spontaneous gamma oscillation *in vitro*. Together, these results suggest that reducing $\alpha 5$ -GABA signaling could improve hippocampus-based cognitive functions. However, other studies involving genetic reduction of $\alpha 5$ subunit expression found behavioural abnormalities related to schizophrenia, arguing against the potential therapeutic value of this approach. For example, $\alpha 5$ (H105R) mutant mice also display attenuated prepulse inhibition (PPI), increased spontaneous locomotor activity^[72] and latent inhibition (LI) deficits.

$\alpha 5$ subtype selective ligands as nootropic drugs

$\alpha 5$ negative allosteric modulators

Inverse agonists at $\alpha 5$ -GABA_A receptors could have potential to enhance cognition. Several $\alpha 5$ GABA_A receptor-selective ligands have tested in animal models. Non-selective BZD inverse agonists may enhance cognitive function in rodents (eg, $\beta 3$ -CCM facilitated spatial learning)^[80] but can cause anxiety-like behavior (eg, $\beta 3$ -CCM; FG 7142)^[79, 80, 81] and seizures (eg, DMCM and FG 7142)^[58, 80, 82, 83] and cause increased vigilance. These side effects prevent these compounds from being clinically useful^[79]. Reduced expression of $\alpha 5$ GABA_A receptors can enhance cognition in some contexts, while $\alpha 1$ subunit inverse agonists can promote seizures. Hence GABA_A receptors containing the $\alpha 5$ subunit would appear to be more promising as a target for the development of cognition-enhancing compounds.

The imidazo-benzodiazepine L-655,708 (also known as FG-8094) has weak inverse agonist efficacy at all four GABA receptor subtypes, but a 50–100-fold higher selectivity for the $\alpha 5$ subunit. This drug enhances performance in normal rats in both the learning and recall phases of the water maze, at a dose previously shown to not be pro-convulsant. Unfortunately, the pharmacokinetic profile of this compound makes it unsuitable for further development as a drug. Other similar $\alpha 5$ selective compounds, including RY-023, RY-024 and RY-080, can promote seizures, which obviously precludes them from clinical use^[79].

To address the side effects associated with targeting GABA_A receptors containing the $\alpha 1$ subunit, some have investigated compounds with selectivity at the $\alpha 5$ receptor^[84]. One example of this is RO4938581, an imidazo-triazolo-benzodiazepine with both binding and functional selectivity at the $\alpha 5$ receptor. This compound reversed scopolamine-induced working memory impairment in the delayed match-to-position task (DMTP task) and diazepam-induced spatial learning impairment in the water maze. More importantly, RO4938581 did not produce anxiety or seizures at ~30% occupancy of hippocampal GABA_A $\alpha 5$ -receptors^[84, 85]. RO4938581 improved performance in a prefrontal cortex-mediated executive function task in monkeys^[84]. This compound also improved cognitive deficits

in rats induced by sub-chronic and neonatal administration of phencyclidine (PCP) (a NMDAR antagonist) in novel object recognition (NOR) and intradimensional/extradimensional attentional set-shifting (ID/ED) task, respectively^[86].

NOR and ID/ED tasks are two preclinical behavioral assays that are related to cognitive tests used to evaluate patients with schizophrenia, such as the MATRICS (measurement and treatment research to improve cognition in schizophrenia) and CNTRICS (cognitive neuroscience treatment research to improve cognition in schizophrenia) test packages^[61, 86–90].

RO4882224, a functional selective inverse agonist for the $\alpha 5$ -GABA_A receptor from Roche, is another imidazo-triazolo-benzodiazepine. RO4882224 enhanced hippocampal LTP and reversed scopolamine-induced working memory impairment in the DMTP task in rodents^[85]. One derivative of RO4938581, Basmisanil (RG1662 or RO5186582), failed at Phase II clinical trial for the treatment of cognitive impairments in Down syndrome^[91] and schizophrenia. There is an ongoing Phase II clinical trial with Basmisanil for treating cognitive deficits in schizophrenia (NCT02953639). Other compounds, such as the thiophene MRK-536, enhanced performance in a Morris water maze of spatial memory^[92] without pro-convulsant effects.

PWZ-029 is another compound with binding selectivity and moderate functional selectivity at $\alpha 5$ -containing GABA_A receptors. It enhanced encoding and consolidation of memory in normal rats tested with the passive avoidance task at doses that did not cause anxiety-like effects or seizure^[93], but had no effect in the active avoidance task. Moreover, PWZ-029 attenuated scopolamine-induced impairment of pavlovian fear-conditioned contextual memory in mice^[94], and reversed scopolamine-induced deficits in novel object recognition, but not the water maze^[95]. PWZ-029 also improved cognitive deficits induced by MK-801 in rodents tested on novel object recognition in the water maze. PWZ-029 did not improve deficits in social recognition memory^[96]. An important caveat with studies of GABA_A $\alpha 5$ drugs in animal models is the difficulty in translating the results to schizophrenia, since many features are difficult to accurately reproduce or measure in animals^[97].

Another compound, $\alpha 5$ IA, is a triazolophthalazine with an equivalent affinity for GABA_A receptors-containing either an $\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 5$ subunit but with greater inverse agonist efficacy than L-655,708 at the $\alpha 5$ subtype (40% for $\alpha 5$ IA vs 17% for L-655,708)^[79, 92, 98]. $\alpha 5$ IA enhances LTP, and improves encoding and recall in the DMTP task, without producing anxiety-like effects or seizure^[98–99]. In addition, $\alpha 5$ IA can reverse ethanol-induced memory impairment in healthy volunteers^[100]. However, $\alpha 5$ IA actually worsened some cognitive impairments in elderly people^[98]. This drug also has a nephrotoxic metabolite that prevents long-term clinical use in humans. The structurally similar compound $\alpha 5$ IA-II has improved oral bioavailability and efficacy selectivity but has pro-convulsant effects^[92]. Another compound MRK-016 (pyrazolotriazine) with high efficacy selectivity at the $\alpha 5$ subtype had promising cognitive effects in animal models but was poorly tolerated in humans^[92, 101].

In summary, inverse agonists at $\alpha 5$ GABA_A receptors have

cognition-enhancing effects in pre-clinical studies. However, many of these drugs have side effects such as anxiety or seizure, and there is the potential for them to amplify psychotic symptoms since they can impair PPI and latent inhibition in animals.

Dual allosteric modulators of $\alpha 5$ GABA_A receptors and $\alpha 7$ nAChRs

The $\alpha 7$ neuronal nicotinic-acetylcholine receptor ($\alpha 7$ nAChR) has been investigated as a valid therapeutic target for treating cognitive deficits in schizophrenia^[102]. Genome-wide association studies have linked deletion of a genetic locus containing the $\alpha 7$ nAChR to increased risk for schizophrenia^[103]. In addition, linkage studies have strongly associated variants in the $\alpha 7$ nAChR gene with deficient P50 auditory gating in schizophrenic patients^[104]. Emerging evidence suggests that P50 sensory gating deficits reflect various cognitive (eg, executive functioning) and perceptual dysfunctions in schizophrenia^[105].

Interestingly, individuals diagnosed with schizophrenia have among the highest rates of cigarette smoking^[106]. Reduction of $\alpha 7$ nAChR function or expression has been identified as a potential mechanism for elevated tobacco use in schizophrenia^[107]. Moreover, multiple studies have found that nicotine, the major psychoactive component of tobacco, can improve cognitive function (eg, attention and spatial working memory) in schizophrenic patients^[106,108] by activating $\alpha 7$ nAChRs. Enhancing the activity of $\alpha 7$ nAChRs is a potential strategy for ameliorating cognitive impairment in schizophrenia.

$\alpha 7$ nAChRs are densely expressed in the hippocampus, especially in interneurons where the density of $\alpha 7$ nAChRs is decreased in schizophrenia^[109]. Given the preferential expression of $\alpha 7$ nAChRs in interneurons, the effects of $\alpha 7$ nAChR on hippocampal neurotransmission is mainly mediated by activation of GABA interneurons^[110]. Pre-synaptic $\alpha 7$ receptors facilitate the release of glutamate and GABA^[102], whereas postsynaptic $\alpha 7$ receptors can regulate GABA_A receptor signaling^[109]. $\alpha 7$ nAChR selective positive allosteric modulators (PAMs) are able to preserve the temporal integrity of neurotransmission^[111] and can improve cognitive deficits in animal models^[102].

522-054, a novel "dual modulator" that acts both as an $\alpha 5$ GABA_A receptor negative allosteric modulator (NAM) and a selective $\alpha 7$ nAChR PAM, was able to restore scopolamine-disrupted deficits in the five-choice serial reaction time test (5-CSRTT) and the eight-arm radial arm maze (RAM). 5-CSRTT is a behavioral test that measures the visual attention and impulsivity in rodents. The test chamber has five holes on one wall, and a reward dispenser on the opposite wall. The task requires the animals to detect a briefly illuminated light presented in one of the five holes and identify the correct spatial location of the hole with nose pokes in order to obtain reward^[112]. The accuracy of visual stimulus discrimination reflects the attentional capacity of the animals^[112]. Intraperitoneal injection of 522-054 in the rats can significantly improve the baseline performance of scopolamine-treated (1.25 mg/kg, ip) animals at a dose of 0.003 mg/kg, suggesting beneficial effects on attention.

The eight-arm RAM task evaluates hippocampus-based spa-

tial learning and memory in rodents. Animals must find the food reward at the end of four randomly chosen maze arms based on spatial navigation cues. At a dose of 0.03 mg/kg, 522-054 had a trend towards reversing the acquired short-term and long-term memory impairments caused by scopolamine (1 mg/kg, ip) in rats.

Flumazenil (an $\alpha 1$ - and $\alpha 5$ -subunit-selective antagonist) and methyllycaconitine (a selective $\alpha 7$ nAChR antagonist) blocked the effect of 522-054 on scopolamine-induced attentional and cognitive deficits, suggesting that simultaneous allosteric modulation of different receptors mediating related functions can have synergistic effects on cognition^[111]. It is possible that compounds with relatively low specificity and moderate potency could be effective^[113].

$\alpha 5$ positive allosteric modulators

$\alpha 5$ -GABA_A receptors mediate the majority of tonic inhibition in hippocampal neurons, and some have suggested that reduced GABA inhibitory input could lead to hyperactivity in the ventral hippocampal dopamine system^[114]. Neural activity in the PFC can synchronize with the neural oscillations in ventral hippocampus, so it is possible that PAMs at $\alpha 5$ GABA_A receptors could restore dopamine tone by selectively reducing vHPC output and vHPC-PFC oscillatory activity, with positive effects on cognition^[115]. PAMs selective for GABA_A $\alpha 5$ receptors improve hippocampal-dependent memory in a rodent model of age-related memory impairment where CA3 neurons have excess firing rates^[116]. In contrast, other $\alpha 5$ -selective PAMs can worsen cognition.

For example, in methylazoxymethanol acetate (MAM)-treated rats, the $\alpha 5$ -selective partial agonist SH-053-2'F-R-CH3 impaired cognitive performance^[58] but normalized the aberrant increase in the number of spontaneously firing dopamine neurons in the VTA to levels comparable to saline-treated rats, and reduced locomotor response to amphetamine^[117]. SH-053-2'F-R-CH3 has no effect on visual recognition and spatial working memory in rhesus monkeys^[118]. Similarly, in an immune-neurodevelopmental model of schizophrenia, the S-isomer of SH-053-2'F-R-CH3 (SH-053-2'F-S-CH3) has detrimental effects on both cognitive function and social interaction^[119]. In contrast, SH-053-2'F-S-CH3 reduces amphetamine-induced hyperactivity^[119]. These findings suggest that $\alpha 5$ -selective PAMs are not suitable for treating cognitive deficits in schizophrenia. However, positive modulation at $\alpha 5$ GABA_A receptors could be a promising adjunctive treatment for targeting positive symptoms in schizophrenia.

Cognitive enhancement through $\alpha 2$ and/or $\alpha 3$ GABA_A receptor modulation

GABA_A- $\alpha 2$ receptor and schizophrenia

GABA_A receptors containing $\alpha 2$ subunits comprise 15%–20% of all GABA_A receptors^[120]. Within the cortex, $\alpha 2$ -containing GABA_A receptors are enriched on the axon-initial segments (AIS) and perisomatic region of pyramidal neurons that are opposed to PVChC and CCK-basket cell (CCK-BCs) terminals, respectively^[121]. PVChCs have arrays of boutons (cartridges)

immunoreactive for the GABA membrane transporter 1 (GAT-1) that innervate the AIS of postsynaptic target neurons^[122]. PVChCs are important for facilitating synchronization of large populations of pyramidal neurons and are thus critical for working memory^[122]. In addition, animal studies have supported an important role for $\alpha 2$ -containing GABA_A receptors in schizophrenia-related cognitive impairment. For example, mice with down-regulated $\alpha 2$ subunits in the frontal cortex have PPI deficits, reduced gamma oscillations, and impairments in working memory^[121].

Postmortem and animal studies have reported that the density of PVChC axon cartridges appears to be reduced in DLPFC layer 2-4 in schizophrenia^[26, 122-123]. Moreover, the immunoreactivity for the GABA_A receptor $\alpha 2$ subunit is markedly elevated in schizophrenia, while the density of $\alpha 2$ -labeled AIS is negatively correlated with the density of PVChCs cartridges^[124]. These findings may be interpreted as compensatory responses to the diminished presynaptic GABA input^[25, 120]. Based on this interpretation, one would predict that augmenting GABA neurotransmission from chandelier neurons through GABA_A receptors containing the $\alpha 2$ subunit could restore gamma-frequency synchronized neuronal activity required for working memory^[125-126]. Therefore, a positive allosteric modulation of $\alpha 2$ subunit-containing GABA_A receptors could be a promising therapeutic strategy in schizophrenia.

GABA_A $\alpha 2$ / $\alpha 3$ positive allosteric modulators

MK-0777 (also known as TPA-023 or L-830982) is a selective partial positive allosteric modulator at $\alpha 2$ and $\alpha 3$ subtypes. MK-0777 can improve performance on several cognitive tasks in patients with schizophrenia. Moreover, MK-0777 improved frontal gamma band power^[127]. However, a subsequent larger clinical trial ($n=60$) failed to replicate these promising findings^[128]. Since MK-0777 is a partial agonist of $\alpha 2$ subunits, a more selective agonist with a greater potency at the GABA_A receptor-containing $\alpha 2$ subunit might work better^[128].

There are questions about the role of PVChCs as neural substrates for the reduced frontal gamma oscillations in patients with schizophrenia^[26]. PVChCs may depolarize cortical pyramidal neurons rather than hyperpolarizing them, and so PVChCs may be a potent source of a slow depolarizing current that mimics the type of slow, NMDA-like depolarization of pyramidal cells^[26]. Moreover, the slow kinetics of $\alpha 2$ GABA_A receptors does not appear to meet the requirement for the strong and fast inhibition required for gamma-band neural oscillations^[26, 129, 130]. In contrast, other studies have indicated that $\alpha 2$ GABA_A receptors are strongly coupled to theta oscillations^[26, 131]. Consequently, the compensatory effects or lower presynaptic GAT-1 and higher postsynaptic GABA_A $\alpha 2$ receptors, in response to decreased GABA neurotransmission may in fact increase EPSCs at the AIS. This could be a way to increase excitation and restore the E/I imbalance in schizophrenia^[26].

GABA_B receptor ligands as cognitive enhancers

GABA_B receptors and schizophrenia

GABA_B receptors are G-protein-coupled receptors consist-

ing of GABBR1 and GABBR2. Unlike GABA_AR, GABA_BRs are found outside the synapse and have high affinity for GABA^[132]. They are widely distributed in the brain and regulate neuronal network activity^[133], neurodevelopment^[134] and synaptic plasticity^[135]. Given their prevalence and widespread distribution in the CNS, it is not surprising that dysfunctions of GABA_B receptors have been implicated in numerous CNS disorders such as major depression^[136], schizophrenia^[137], bipolar disorder^[138] and seizures.

Multiple lines of evidence implicate GABA_B receptors in the pathophysiology of schizophrenia^[139]. GABA_B receptors are markedly reduced in the cerebellum^[139], hippocampus^[140], entorhinal cortex, inferior temporal cortex^[141] of patients with schizophrenia compared with healthy controls. GABA_B receptors in the entorhinal cortex and hippocampus are important for memory^[142]. Agonist activation of GABA_B inhibits neuronal excitation and gamma oscillations^[143], while antagonists promote theta and gamma oscillations^[144]. Inhibition of postsynaptic GABA_B receptors can enhance LTP by lengthening NMDA receptor-mediated currents^[145]. Inhibition of presynaptic GABA_B receptor enhances GABA release, thereby decreasing calcium conductance and subsequent GABA release^[146].

GABA_B receptor antagonists

GABA_B receptor antagonists can improve cognition^[145]. Proposed mechanisms include facilitating synaptic plasticity and LTP^[147-148] and entraining neuronal oscillations^[144, 149]. For example, intrahippocampal infusion of the antagonist 2-OH saclofen can markedly reverse scopolamine-induced impairments in LTP and Y-maze performance^[148]. Similarly, the antagonists CGP 55845 and CGP 52432 enhance impaired LTP in the dentate gyrus of Ts65Dn mice, a genetic mouse model of Down Syndrome^[147]. Moreover, CGP 55845 significantly increased gamma and theta oscillations in rat brain slices^[149]. Another antagonist SCH 50911 was also found to increase gamma power^[150].

So far, a variety of GABA_B receptor antagonists have displayed cognition-enhancing effects in animal models of psychiatric disorders. Representative GABA_B receptor antagonists and their effects on cognition are summarized in Table 1. CGP 36742 (also known as SGS742) was the first^[151] and the only antagonist tested in clinical trials for mild cognitive impairment. SGS742 enhances cognition in animal models and in clinical phase II trials (Table 1) (For a comprehensive review, see^[145, 151]). However, no current data are available with regards to treating cognitive deficits in schizophrenia using SGS742.

Despite evidence for cognitive enhancing effects of GABA_B receptor antagonists, few studies have examined these compounds in animal models of schizophrenia. For example, in a recent apomorphine-susceptible (APO-SUS) young rat model displaying schizophrenia-relevant features^[146], the level of GAD67 and the density of GAD67-positive cells were reduced. However, basal synaptic input to pyramidal neurons was unaltered. In contrast, the paired-pulse ratio (PPR) at lon-

Table 1. Effects of representative GABA(B) receptor antagonists on cognition in preclinical/clinical trials.

Compound	Pharmacological properties	Species	Behavioural test/clinical trial	Effect on cognition	References
CGP 35348 (IC ₅₀ =34 μmol/L)	Showed affinity for the GABA(B) receptor only 200 and 500 times more potent than the GABA(B) receptor antagonists OH-saclofen and phaclofen, respectively [153] Displayed higher affinity for postsynaptic vs presynaptic GABA(B) receptors	Scopolamine-induced amnesia in CD1 mice	Passive avoidance test	Amnesic effect ↓	[152]
		Baclofen-treated male Sprague-Dawley rats Streptozotocin-induced dementia in rats Hypoxia ischemia insult induced-neonatal brain damage albino mice	T maze footshock avoidance test Novel object recognition Morris water maze Morris water maze	Retention memory↑ Recognition memory↑ Recognition memory↑ Spatial learning and memory M↑,F↔	[153] [154] [155] [156, 157]
CGP 36742 (SGS742) (IC ₅₀ =36 μmol/L)	Interacted very weakly with GABA(A) receptor (IC ₅₀ =500 μmol/L)	Streptozotocin-induced dementia in rats Insular injection of CGP 35348 in Sprague-Dawley rats	Morris water maze Intellicage	Retention memory↑ Position learning, punitive learning and punitive reversal learning ↓; position reversal learning↔	[155] [158]
		Mice Male Sprague-Dawley rats CD1 mice Rhesus monkeys Baclofen- and scopolamine-induced deficits in Wistar rats GAERS rats Long-Evans rats	Footshock passive avoidance test Social recognition test Eight arm radial maze Conditional space-color test Morris water maze	Retention memory F↑ (male mice were not tested) Retention memory↑ Spatial learning and memory↑ Cognitive function↑ Spatial learning and memory↑	[159, 160] [159, 161] [162] [159] [163]
CGP 55845 (IC ₅₀ =5 nmol/L)	Acted at pre- and post-synaptic GABA(B) receptors in the hippocampus [169]	1.10 patients aged 59–85 with mild cognitive impairment (MCI) were treated for 8 weeks at dose of 600 mg t.i.d. of SGS742 280 patients diagnosed with mild to moderate Alzheimer's disease	Two-way active avoidance test Eight- and twelve-arm radial mazes Double-blind, placebo-controlled Phase IIa clinical trial	Cognitive function↑ Spatial learning and memory↑ attention↑ (i.e. choice reaction time, visual information processing); working memory↑ (i.e. pattern recognition speed) Failed to meet the efficacy end points	[164] [165] [165] [166]
		Rats GHL-treated rats GHL-treated mice GHL-treated rats GHL-treated mice Cognitively-impaired aged Fischer 344 rats Ts65Dn mice Ts65Dn mice Ts65Dn mice Ts65Dn mice	Active avoidance test Active avoidance test Active avoidance test Passive avoidance test Passive avoidance test Olfactory discrimination test Novel place recognition Contextual fear conditioning task Novel object recognition T-maze Morris water maze	Learning and retention memory↑ Learning and retention memory↑ Learning and retention memory↑ Learning and retention memory↑ Learning and retention memory↔ Learning↑ Recognition memory↑ Learning and memory↑ Recognition memory↑ Memory and spatial learning↔ Spatial learning and memory↔ in both M&F (1 mg/mL solvent/kg body weight)	[167] [168] [168] [168] [168] [170] [147] [147] [147] [147] [171]
CGP 55845 (IC ₅₀ =5 nmol/L)	~1000 times more potent than CGP 35348	Hypoxia ischemia insult induced-neonatal brain damage albino mice	Morris water maze	Spatial learning and memory↔ in both M&F (1 mg/mL solvent/kg body weight)	[171]

(To be continued)

Compound	Pharmacological properties	Species	Behavioural test/clinical trial	Effect on cognition	References
CGP 62349 (IC ₅₀ =2 nmol/L)	The antagonists reversed the effects of an EC70 concentration of GABA (1 µmol/L) and forskolin (10 µmol/L) with the rank order of potency: CGP62349>CGP54626A>SCH 50911>phaclofen=saclofen [172]	Rats GHBL-treated rats GHBL-treated mice	Active avoidance test Active avoidance test Active avoidance test	Learning and retention memory ↑ Learning and retention memory ↔ Learning and retention memory ↑ (more potent than CGP 55845) Learning and retention memory ↔ Learning and retention memory ↔	[167] [168] [168] [168] [168]
CGP 71982 (IC ₅₀ =8 nmol/L)	NA	Rats GHBL-treated rats GHBL-treated mice GHBL-treated rats GHBL-treated mice	Active avoidance test Active avoidance test Active avoidance test Passive avoidance test Passive avoidance test	Learning and retention memory↑ Learning and retention memory ↔ Learning and retention memory ↔ Learning and retention memory↑ Learning and retention memory ↑	[167] [168] [168] [168] [168]

Male (M) Female (F) Increase↓ Decrease↑ No effect ↔
 Gamma-hydroxybutyrolactone (GHBL)
 Ts65Dn mice: a genetic mouse model of Down Syndrome
 Genetic absence epilepsy rats of Strasbourg (GAERS): a model of epilepsy
 NA: Data not available

ger inter-stimulus intervals was decreased in APO-SUS rats, indicative of a reduced GABA release. This reduction could be caused by enhanced GABA_B receptor signaling. Interestingly, the application of CGP 55845 can completely restore the level of PPR and cause a decrease in GABA_B signaling. The authors showed that CGP 55845 is likely to act at presynaptic GABA_B receptors rather than postsynaptic receptors. Therefore, these findings are in line with the hypothesis that inhibition of presynaptic GABA_B receptors could enhance GABA release. Decreased inhibitory drive from interneurons (eg, PV neurons) onto their postsynaptic targeted pyramidal neurons could be a core pathophysiological feature of schizophrenia^[173], and that reduced GABA neurotransmission is highly correlated to cognitive defects in schizophrenia^[146]. GABA_B receptor antagonists require further investigations, especially in animal models related to schizophrenia in order to determine their potential for treating cognitive impairments in schizophrenia.

GABA_B receptor agonists

The prototypic GABA_B receptor agonist Baclofen is currently the only marketed drug targeting GABA_B receptors. Numerous studies have reported that the Baclofen and other GABA_B receptor agonists impair cognition in animal models^[174]. However, these results are not consistent. Baclofen can ameliorate the recognition memory impairment induced by methamphetamine^[175] and the spatial working memory deficits induced by chronic cerebral hypoperfusion in rats^[176] (for a more comprehensive review, see^[174]). More studies need to be conducted to confirm the effects of Baclofen on cognition.

GABA_B receptor allosteric modulators

The data for the effects of PAM and NAM targeting GABA_B receptors on cognition is limited. In the mouse passive avoidance cognition paradigm, a selective GABA_B receptor PAM GS39783 showed no deleterious effects on cognition in contrast to Baclofen (1 mg/kg), which significantly impaired cognitive performance^[177]. In 2015, the novel compound **7** from Astellas Pharma, a sulfur-containing bicyclic compound functioning as a GABA_B receptor PAM, was used for the prevention or treatment of cognitive disorders, schizophrenia and pain (publication number for patent application: WO2015056771)^[178].

Epigenetic therapies for cognitive impairment

Alterations in epigenetic regulations in schizophrenia

Epigenetic mechanisms (eg, histone modification, chromatin remodeling, DNA methylation) can synergistically interact with genetics^[53] to mediate GABA system abnormalities in schizophrenia. Prominent genes include *GAD1*, *RELN*, *BDNF* and *GABAB3*^[53, 179]. Several excellent reviews discuss epigenetic mechanisms in the neurobiology of neuropsychiatric disorders including schizophrenia^[51, 53, 180-182]. These articles highlight several key findings about epigenetic alterations in schizophrenia.

Histone modifications in schizophrenia are shifted from open chromatin (H3K4-trimethylation, which positively regu-

lates gene expression)^[53] to repressive histone methylation (H3K27-methylation that negatively regulates gene expression) at GABAergic gene promoters (eg, *GAD1*) in PFC of some subjects with schizophrenia^[51]. In addition, the expression of histone deacetylases (HDAC) such as HDAC1 that facilitate downregulation of gene expression is increased in schizophrenic brains and correlated with reduced *GAD67* expression^[51, 183]. Interestingly, overexpression of neuronal HDAC1 in mouse mPFC (but not in dorsal or ventral hippocampus) resulted in schizophrenia-like phenotypes such as impaired short-term memory, PPI and synaptic plasticity^[184]. In fact, many CNS disorders with cognitive impairment also have reduced histone acetylation^[185].

DNA (cytosine-5)-methyltransferase (DNMT) (eg, DNMT1 and 3a) expression and DNA hypermethylation are abnormally increased in gene promoters in schizophrenia, which consequently results in MecP2-mediated gene silencing of GABAergic candidate genes such as *GAD1*, *RELN*, *SOX10*^[51] and *BDNF*^[51, 53]. Therefore, epigenetic modulators acting in GABAergic system could remediate the epigenetic alterations observed in schizophrenia.

HDAC and DNMT inhibitors for the treatment of cognitive defects neuropsychiatric disorders

In animal models, treatment with HDAC inhibitors such as trichostatin A (TSA), valproate (VPA) and MS-275 can increase the expression of reelin and *GAD67* by activating demethylation^[49, 186]. In addition, co-administration of VPA with antipsychotics (eg, clozapine^[187], olanzapine and quetiapine) can synergistically potentiate VPA-induced promoter demethylation and chromatin remodeling (see review by^[49]) and enhance antipsychotic effects^[188]. VPA can elevate GABA concentration via synthesis, reuptake, and metabolism. In spite of a few positive reports^[189, 190], VPA is likely to be detrimental to cognition as shown in animal studies^[191]. Class I HDACs play an important role in neuronal and brain development^[192], and are the best-studied HDACs with respect to cognition (for comprehensive review on this topic, see^[185]). HDAC inhibition/deletion facilitates upregulation of a key set of genes involved in cognitive functions and enhances synaptic plasticity and long-term memory^[193, 194].

Class I HDACs HDAC 2 and 3^[185, 188, 195-197] and Class II HDACs HDAC 5^[198] and 6^[188, 195, 197] could be potential targets for improving cognition in neuropsychiatric disorders. HDAC 1 appears to have inconsistent effects on cognition^[184, 188, 195, 198-200]. Representative HDAC inhibitors that have effects on cognition in preclinical and/or clinical studies for a variety of neuropsychiatric disorders are summarized in Table 2. However, these inhibitors are also reported to cause impairment in memory, learning and cognition in some other studies^[182] in a brain region- and HDAC isoform-specific manner^[188]. Development of selective inhibitors may reduce undesirable side effects, while still retaining pro-cognitive effects. (For comprehensive reviews on Epigenetics in CNS diseases/cognition, see^[49, 182, 185, 188, 201-202].)

Similar to HDAC inhibitors, DNMT (eg, DNMT1 and 3a)

inhibitors such as 5AZA and zebularine are also able to restore the expression of reelin and *GAD67*. But the majority of these compounds or drugs are clinically used for cancer treatment (eg, 5AZA, zebularine, RG108, decitabine). So far, there are few reported studies of DNMT inhibitors in animal models of schizophrenia^[182].

In summary, the field of epigenetic drugs for the treatment of cognitive impairment in neuropsychiatric diseases such as schizophrenia is at an early stage and development of these drugs for cognition remains a great challenge. Although HDAC and DNMT inhibitors could be of potential therapeutic value in ameliorating cognitive deficits among high-risk individuals with schizophrenia, obstacles such as the lack of subtype- or brain region-specificity^[188] or capacity to cross the blood-brain-barrier (BBB)^[182, 240], difficulty with the establishment of dose ranges or treatment duration in clinical trials^[188], and severe toxicity^[188] have hindered the translation to the clinic.

GABA prodrug with pro-cognitive and antipsychotic effects

BL-1020 (also known as CYP-1020) is being developed by Bio-LineRx. It is an ester that combines the dopamine antagonism of perphenazine (a typical antipsychotic D₂/5-HT₂ antagonist drug) with GABA agonist activity^[241]. This drug has promising pro-cognitive and antipsychotic effects in rodent models of schizophrenia^[242-243] and phase II clinical trials for schizophrenia (NCT00480571, NCT00567710, NCT00722176)^[241, 244].

Preclinical studies indicate that BL-1020 can cross the BBB^[244] and is less likely to cause neurological or metabolic side effects than current antipsychotics^[243, 244]. In phase II a/b clinical trials, patients with chronic schizophrenia or schizoaffective disorder treated with BL-1020 demonstrated significant improvements in cognition and psychotic symptoms. However, the most recent IIb-III clinical trial (NCT01363349) designed to compare the cognitive effects of treatment with CYP-1020 to risperidone was terminated in 2013 because CYP-1020 did not meet its standard efficacy end points. Therefore, future phase II/III clinical trials are required to determine the clinical efficacy of BL-1020 compared with the established antipsychotics such as risperidone or clozapine. Taken together, BL-1020 has shown promising signs as a novel antipsychotic and pro-cognitive compound with excellent therapeutic effects for psychosis and cognitive impairments, and produces fewer side effects that commonly occur with typical and atypical antipsychotic medications. (For review, see^[244])

GAT-1 inhibitors

GAT-1, the main plasma membrane GABA transporter in brain^[132], is localized almost exclusively to axon terminals^[245], which mediates the uptake of extracellular GABA^[132]. This activity is generally thought to terminate the synaptic effects of GABA^[132]. It has been shown that GAT-1-mediated GABA transport regulates GABA_B receptor electrophysiological activity through synaptic GABA^[132]. Blockade of GAT-1 can enhance postsynaptic GABA_BR-mediated IPSPs^[246] and pre-

Table 2. Summary of HDAC inhibitors and their effects on cognition in preclinical and clinical studies for neuropsychiatric disorders.

HDACi group	HDAC inhibitor	Target HDACs	Diseases targeted	Model organisms	Effects on cognition	References
Benzamide derivatives	MS-275 (Entinostat)	HDAC1, 2, 3, 9	Mood and anxiety disorders [203]	129S1/SvImJ (S1) mice; stressed Sprague Dawley male rats at the prepubertal age	Rescued extinction-consolidation deficits and temporal order recognition (TOR) memory deficits	[204,205]
		HDAC1,2,3,6,8 [206]	SZ	Mice suffering early life stress (i.e. maternal separation)	↑short-term memory, synaptic plasticity and PPI	[184]
		HDAC6/HDAC1	AD	Mouse cortical primary neurons & transgenic Egr1-EGFP mice	↑associative learning, BDNF and memory-related genes; enhance synaptogenesis	[206]
Hydroxamic acids	Tubastatin or Tubastatin A	HDAC6/HDAC1	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice; rTg4510 mice	↑working memory; ↑spatial and reversal learning and memory; ↓hyperphosphorylated tau and Aβ	[207]
		HDAC1, 2, 3	PTSD	Mice model of fear extinction	Attenuated remote fear memories; upregulated a key set of neuroplasticity-related genes	[208]
Short chain fatty acids	Suberoylanilide hydroxamic acid (SAHA, Vorinostat)	Classes I, II &IV	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice; aged mice	↑hippocampus-dependent associative learning	[189]
		Class I and II	RTS	Tg2576 mice	Failed to rescue memory deficits due to low brain availability; ↑ LTP and ↓ LTD <i>in vitro</i>	[209]
		Class I and II	?	CBP ^{+/−} Mice	↑hippocampus-dependent associative learning; enhance late phase of LTP	[210]
		Classes I and II	?	HDAC2 over-expressed mice	↑hippocampus-dependent associative learning; enhance dendritic spine density, synaptogenesis and LTP	[200]
Short chain fatty acids	Trichostatin A (TSA)	Classes I and II	RTS	CREB ^{ca} mutant mice	↑contextual fear memory	[211]
		Class I and II	Metabolic syndrome	High-fat diet fed mice	↑spatial learning and memory and BDNF; alleviate oxidative stress	[212]
		Class I and II	AD	hAPP 3xTg	↑spatial learning and memory; ↑tau phosphorylation and Aβ	[213]
		Classes I, IIa	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice; CKp-25 mice; aged mice	↑associative, spatial, contextual and object recognition memory; ↑dendritic spine density	[189, 214, 215, 216, 217]
		Classes I, IIa	PD	6-Hydroxydopamine (6-OHDA)-treated rats	↑executive function	[218]
Short chain fatty acids	Sodium butyrate (NaB)	Depression	Depression	Maternal deprivation and chronic mild stress treated rats	↑object recognition memory; ↑ BDNF	[219]
		ASD	ASD	Mice given prenatal exposure to VPA	↑object recognition memory; ↑ CA1 region dendritic spine density	[190]
		SZ	SZ	Quadruple blind randomized placebo controlled phase II trial (NCT03010865)	To evaluate a series of cognitive functions in patients with SZ	https://clinicaltrials.gov

(To be continued)

HDACi group	HDAC inhibitor	Target HDACs	Diseases targeted	Model organisms	Effects on cognition	References
	Phenylbutyrate/ Sodium Phenylbutyrate	-	SZ AD	Rats with neonatal ventral hippocampus lesions Tg2576 mice; AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice	↑associative learning ↑spatial reference memory deficits; ↔Aβ _{42/40} and senile plaques;	[220] [221,222]
	Valproate/ Valproic Acid (VPA)	Classes I & IIa	HD PD ASD AD	N171-82Q mice Aged Y39C mice Mice given prenatal exposure to VPA AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice	↓hyperphosphorylated tau; alleviate ER stress ↑survival and ↓ gross brain and neuronal atrophy ↑spatial learning and memory ↑object recognition memory, ↑ CA1 region dendritic spine density ↑contextual memory	[223] [224] [190] [189]
	Sodium Valproate	Class I [227]	Epilepsy Epilepsy	Patients with epilepsy Rats with convulsive status epilepticus (CSE)	↓ cognition ↑spatial cognitive dysfunction of P35 CSE rats, but ↓ cognition of normal P15 and P35 rats	[225, 226] [228]
Mercapto acetamide Carbamide based inhibitor	W2 BRD4884	Class II HDACs HDAC2	AD neuropsychiatric diseases (eg. SZ/PTSD)	hAPP 3xTg CKp-25 mice	↑spatial learning and memory; ↓tau phosphorylation and Aβ Rescued memory defects in contextual fear conditioning	[213] [197]
Urea based inhibitor	BRD6688	HDAC2	neuropsychiatric diseases (eg SZ/PTSD)	CKp-25 mice	Rescued memory defects in contextual fear conditioning	[197]
Carboxamide	RGFP963	HDAC1, 2 and 3	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice	↑contextual fear memory deficits by inducing synaptogenesis and increasing hippocampal spine density	[196]
	RGFP968	HDAC1, 2 and 3	PTSD	Male C57BL/6J mice Primary cortical neuron	Enhanced consolidation of cued fear extinction Induced synaptogenesis and increased spine density in hippocampus	[229] [196]
	RGFP966	HDAC3	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice	Failed to rescue contextual fear memory deficits or to stimulate synaptogenesis	[196]
			PTSD	Male C57BL/6J mice	Failed to enhance consolidation of cued fear extinction	[229]
			-	Auditory memory rat model	Enhanced memory of a acoustic details associated with sound-to-reward learning; affects auditory cortical plasticity	[194]
			HD	HdhQ111 knock-in mice	↑spatial and recognition long-term memory and motor learning deficits; normalizes hippocampal expression of memory-dependent genes; ↓ mutant Huntingtin oligomers	[230]

(To be continued)

HDACi group	HDAC inhibitor	Target HDACs	Diseases targeted	Model organisms	Effects on cognition	References
	RGFP136	HDAC3	-	Homozygous deletions of <i>Hdac3</i> in CA1 of the dorsal hippocampus	↑ long-term memory for object recognition and location	[231]
	Rocilinostat (ACY-1215)	HDAC6	AD	AβPP ^{swE} /PS1 ^{ΔE9} (PAP) double-transgenic mice	↑ non-spatial working memory; ↑ hippocampus-dependent spatial learning and memory	[207]
Quinazolin-4-one derivatives	HDACi-4b EXX001688; EVX	HDAC1,3,6 Class I and IIb	HD AD	N171-82Q transgenic mice Aged rats	↑ cognitive ability in T-maze task Had no effect on age-related spatial learning and memory	[232] [233]
Amide of nicotinic acid	Nicotinamide (nicotinamide)	SIRT1	AD	3xTg mice	↑ spatial and contextual learning; has no effect on object recognition memory	[234]
Sulfonic acid	Suramin	SIRT1	ASD/SZ?	Maternal immune activation (MIA) mice; Fragile X (<i>Fmr1</i> knockout) mice	Reversed social behavior, novelty preference and metabolism deficits	[235, 236]
-	CM-414 (compound 7)	-	ASD AD	Double-blind randomized placebo-controlled phase I/II trial (NCT02508259) Tg2576 mice	↑ in language, social interaction; ↓ restricted or repetitive behaviors Rescued associative fear/spatial learning and memory impairment; ↑ synaptic plasticity, dendritic spine density and memory-related genes; ↓ Aβ and tau	[237] [238]

Table adapted from [239]

Schizophrenia (SZ), Alzheimer's disease (AD), Post-traumatic stress disorder (PTSD), Rubinstein-Taybi syndrome (RTS), Parkinson's disease (PD)

Autism spectrum disorder (ASD), Huntington's disease (HD)

Increase ↑ Decrease ↓ No effect ↔

amyloid-β (Aβ)

synaptic effects (For a detailed review, see^[132]). Therefore, GAT-1 inhibitors may enhance cognition.

In a lipopolysaccharide (LPS)-treated rat model which mimics the prenatal inflammation thought to contribute to schizophrenia^[247], the GAT-1 inhibitor Tiagabine (TGB) (also known as Gabitril) prevents impairments in LTP and LTD (long-term depression) in male offspring but had no effect on LTD in control rats^[248]. So far, several studies, including double-blind, placebo-controlled trials, have reported that TGB monotherapy or adjunctive therapy for CNS disease (eg, epilepsy) has no negative effects on cognitive function^[249, 250]. In contrast, data for cognitive enhancing effects of this drug is lacking. Currently, a phase III clinical trial evaluating the effects of TGB on brain deficits including neurocognitive functions and clinical symptoms during early-stage schizophrenia is in progress (NCT00179465).

Conclusion

In conclusion, the literature reviewed above suggests that restoring pyramidal neuronal inhibition could normalize aberrant cortical and hippocampal neuronal oscillations in schizophrenia. This could ameliorate cognitive impairments such as episodic memory, working memory and executive function in schizophrenia and other neuropsychological disorders. Pharmacological modulation of synaptic or extrasynaptic GABAergic signaling mediated by GABA_A and GABA_B receptors could restore disrupted neuronal network synchronization within or between brain regions-associated with learning and memory, which can in turn restore E/I imbalance and cognitive deficits in patients with schizophrenia. Electrophysiological measurements such as electroencephalography (EEG) signals, can provide an index of functional connectivity in the brain^[251], that could serve as endophenotypes for screening candidate cognitive enhancing drugs for schizophrenia.

Overall, most of the potential cognitive-enhancing pharmacological treatments targeting the GABA neurotransmitter system have shown promise in pre-clinical studies with animals. This is also true for $\alpha 5$ subunit-selective negative allosteric modulators. The paucity of data demonstrating therapeutic effects of these drugs in clinical studies, however, has raised questions about how valid these pre-clinical studies are for predicting clinical therapeutic effects in patients. This problem is generic to many aspects of animal models for psychiatric disease and treatment^[97], and emphasizes the need for continuing development of more powerful translational animal behavioural assays with better predictive validity. One promising approach is the use of touchscreen-based cognitive tests that can deliver cognitive tests that are nearly identical for both humans and rodents^[252].

Moreover, dual allosteric modulators acting at two different receptors mediating similar functions could produce synergic effects on cognition. This synergistic strategy could reduce the dosage required for achieving optimal therapeutic efficacy, reduce side effects caused by individual drugs, and potentiate the pro-cognitive effects. Therefore, deliberately targeting multiple receptors could be a promising strategy for improv-

ing the pharmacological treatment of cognitive impairment in schizophrenia and other neuropsychiatric disorders.

Finally, epigenetic therapies, in particular selective class I HDAC inhibitors, require further modifications in order to increase brain regional selectivity, capacity to cross the blood-brain-barrier, and to reduce systemic toxicity. The epigenetic machinery is difficult to manipulate with specificity, and this is especially problematic for pharmacological manipulation of higher mental functions such as cognition. However, there are examples that show promise, such as inhibiting histone methyltransferases to treat anxiety and depression^[253]. Given the complexity of human cognition, and the heterogeneity of patients with schizophrenia, it is likely that targeting multiple systems and individualizing pharmacological treatment to each patient, is the way forward.

References

- 1 Kahn RS, Keefe RS. Schizophrenia is a cognitive illness: time for a change in focus. *JAMA Psychiatry* 2013; 70: 1107–12.
- 2 Kahn RS, Sommer IE, Murray RM, Meyer-Lindenberg A, Weinberger DR, Cannon TD, et al. Schizophrenia. *Nat Rev Dis Primers* 2015; 1: 15067.
- 3 Larson MK, Walker EF, Compton MT. Early signs, diagnosis and therapeutics of the prodromal phase of schizophrenia and related psychotic disorders. *Expert Rev Neurother* 2010; 10: 1347–59.
- 4 Lewis DA, Levitt P. Schizophrenia as a disorder of neurodevelopment. *Annu Rev Neurosci* 2002; 25: 409–32.
- 5 Bora E. Neurodevelopmental origin of cognitive impairment in schizophrenia. *Psychol Med* 2015; 45: 1–9.
- 6 Volk DW, Lewis DA. Early developmental disturbances of cortical inhibitory neurons: contribution to cognitive deficits in schizophrenia. *Schizophr Bull* 2014; 40: 952–7.
- 7 Benes FM, Berretta S. GABAergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology* 2001; 25: 1–27.
- 8 Kelsom C, Lu W. Development and specification of GABAergic cortical interneurons. *Cell Biosci* 2013; 3: 19.
- 9 Moran LV, Hong LE. High vs low frequency neural oscillations in schizophrenia. *Schizophr Bull* 2011; 37: 659–63.
- 10 Kann O, Papageorgiou IE, Draguhn A. Highly energized inhibitory interneurons are a central element for information processing in cortical networks. *J Cereb Blood Flow Metab* 2014; 34: 1270–82.
- 11 Schmidt MJ, Mirnic K. Neurodevelopment, GABA system dysfunction, and schizophrenia. *Neuropsychopharmacology* 2015; 40: 190–206.
- 12 Pocklington AJ, Rees E, Walters JT, Han J, Kavanagh DH, Chambert KD, et al. Novel findings from CNVs implicate inhibitory and excitatory signaling complexes in schizophrenia. *Neuron* 2015; 86: 1203–14.
- 13 Lee FH, Zai CC, Cordes SP, Roder JC, Wong AH. Abnormal interneuron development in disrupted-in-schizophrenia-1 L100P mutant mice. *Mol Brain* 2013; 6: 20.
- 14 Fazzari P, Paternain AV, Valiente M, Pla R, Lujan R, Lloyd K, et al. Control of cortical GABA circuitry development by Nrg1 and ErbB4 signalling. *Nature* 2010; 464: 1376–80.
- 15 Dean K, Murray RM. Environmental risk factors for psychosis. *Dialogues Clin Neurosci* 2005; 7: 69–80.
- 16 Tsuang MT, Stone WS, Faraone SV. Genes, environment and schizophrenia. *Br J Psychiatry Suppl* 2001; 40: s18–24.
- 17 Uhlhaas PJ. High-frequency oscillations in schizophrenia. *Clin EEG*

- Neurosci 2011; 42: 77–82.
- 18 Patel AB, de Graaf RA, Martin DL, Battaglioli G, Behar KL. Evidence that GAD65 mediates increased GABA synthesis during intense neuronal activity *in vivo*. *J Neurochem* 2006; 97: 385–96.
- 19 Blum BP, Mann JJ. The GABAergic system in schizophrenia. *Int J Neuropsychopharmacol* 2002; 5: 159–79.
- 20 Akbarian S, Huang HS. Molecular and cellular mechanisms of altered GAD1/GAD67 expression in schizophrenia and related disorders. *Brain Res Rev* 2006; 52: 293–304.
- 21 Lett TA, Voineskos AN, Kennedy JL, Levine B, Daskalakis ZJ. Treating working memory deficits in schizophrenia: a review of the neurobiology. *Biol Psychiatry* 2014; 75: 361–70.
- 22 Hashimoto T, Volk D, Eggan S, Pierri J, Sun Z, Sampson A, et al. Altered gene expression in parvalbumin-containing GABA neurons in the prefrontal cortex of subjects with schizophrenia. *Schizophr Res* 2003; 60: 71.
- 23 Volk DW, Austin MC, Pierri JN, Sampson AR, Lewis DA. Decreased glutamic acid decarboxylase67 messenger RNA expression in a subset of prefrontal cortical gamma-aminobutyric acid neurons in subjects with schizophrenia. *Arch Gen Psychiatry* 2000; 57: 237–45.
- 24 Hashimoto T, Volk DW, Eggan SM, Mirnics K, Pierri JN, Sun Z, et al. Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J Neurosci* 2003; 23: 6315–26.
- 25 Lewis DA. Inhibitory neurons in human cortical circuits: substrate for cognitive dysfunction in schizophrenia. *Curr Opin Neurobiol* 2014; 26: 22–6.
- 26 Lewis DA, Curley AA, Glausier JR, Volk DW. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci* 2012; 35: 57–67.
- 27 Orhan F, Fatouros-Bergman H, Goiny M, Malmqvist A, Piehl F, Karolinska Schizophrenia Project C, et al. CSF GABA is reduced in first-episode psychosis and associates to symptom severity. *Mol Psychiatry* 2017. doi: 10.1038/mp.2017.25.
- 28 Uhlhaas PJ, Singer W. Abnormal neural oscillations and synchrony in schizophrenia. *Nat Rev Neurosci* 2010; 11: 100–13.
- 29 Howard MW, Rizzuto DS, Caplan JB, Madsen JR, Lisman J, Aschenbrenner-Scheibe R, et al. Gamma oscillations correlate with working memory load in humans. *Cereb Cortex* 2003; 13: 1369–74.
- 30 Chen CM, Stanford AD, Mao X, Abi-Dargham A, Shungu DC, Lisanby SH, et al. GABA level, gamma oscillation, and working memory performance in schizophrenia. *Neuroimage Clin* 2014; 4: 531–9.
- 31 Gallinat J, Winterer G, Herrmann CS, Senkowski D. Reduced oscillatory gamma-band responses in unmedicated schizophrenic patients indicate impaired frontal network processing. *Clin Neurophysiol* 2004; 115: 1863–74.
- 32 Montgomery SM, Buzsaki G. Gamma oscillations dynamically couple hippocampal CA3 and CA1 regions during memory task performance. *Proc Natl Acad Sci U S A* 2007; 104: 14495–500.
- 33 Lynn PA, Sponheim SR. Disturbed theta and gamma coupling as a potential mechanism for visuospatial working memory dysfunction in people with schizophrenia. *Neuropsychiatry Electrophysiol* 2016; 2: 7.
- 34 Caixeta FV, Cornelio AM, Scheffer-Teixeira R, Ribeiro S, Tort AB. Ketamine alters oscillatory coupling in the hippocampus. *Sci Rep* 2013; 3: 2348.
- 35 Sohal VS, Zhang F, Yizhar O, Deisseroth K. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* 2009; 459: 698–702.
- 36 Cardin JA, Carlen M, Meletis K, Knoblich U, Zhang F, Deisseroth K, et al. Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature* 2009; 459: 663–7.
- 37 Bartos M, Vida I, Jonas P. Synaptic mechanisms of synchronized gamma oscillations in inhibitory interneuron networks. *Nat Rev Neurosci* 2007; 8: 45–56.
- 38 Amilhon B, Huh CY, Manseau F, Ducharme G, Nichol H, Adamantidis A, et al. Parvalbumin interneurons of hippocampus tune population activity at theta frequency. *Neuron* 2015; 86: 1277–89.
- 39 Gonzalez-Burgos G, Lewis DA. GABA neurons and the mechanisms of network oscillations: implications for understanding cortical dysfunction in schizophrenia. *Schizophr Bull* 2008; 34: 944–61.
- 40 Gonzalez-Burgos G, Hashimoto T, Lewis DA. Alterations of cortical GABA neurons and network oscillations in schizophrenia. *Curr Psychiatry Rep* 2010; 12: 335–44.
- 41 Shin YW, O'Donnell BF, Youn S, Kwon JS. Gamma oscillation in schizophrenia. *Psychiatry Investig* 2011; 8: 288–96.
- 42 Curley AA, Lewis DA. Cortical basket cell dysfunction in schizophrenia. *J Physiol* 2012; 590: 715–24.
- 43 Glausier JR, Fish KN, Lewis DA. Altered parvalbumin basket cell inputs in the dorsolateral prefrontal cortex of schizophrenia subjects. *Mol Psychiatry* 2014; 19: 30–6.
- 44 Volman V, Behrens MM, Sejnowski TJ. Downregulation of parvalbumin at cortical GABA synapses reduces network gamma oscillatory activity. *J Neurosci* 2011; 31: 18137–48.
- 45 Lazarus MS, Krishnan K, Huang ZJ. GAD67 deficiency in parvalbumin interneurons produces deficits in inhibitory transmission and network disinhibition in mouse prefrontal cortex. *Cereb Cortex* 2015; 25: 1290–6.
- 46 Tukker JJ, Fuentealba P, Hartwich K, Somogyi P, Klausberger T. Cell type-specific tuning of hippocampal interneuron firing during gamma oscillations *in vivo*. *J Neurosci* 2007; 27: 8184–9.
- 47 Nyhus E, Curran T. Functional role of gamma and theta oscillations in episodic memory. *Neurosci Biobehav Rev* 2010; 34: 1023–35.
- 48 Weickert CS, Hyde TM, Lipska BK, Herman MM, Weinberger DR, Kleinman JE. Reduced brain-derived neurotrophic factor in prefrontal cortex of patients with schizophrenia. *Mol Psychiatry* 2003; 8: 592–610.
- 49 Guidotti A, Auta J, Chen Y, Davis JM, Dong E, Gavin DP, et al. Epigenetic GABAergic targets in schizophrenia and bipolar disorder. *Neuropharmacology* 2011; 60: 1007–16.
- 50 Nakazawa K, Zsiros V, Jiang Z, Nakao K, Kolata S, Zhang S, et al. GABAergic interneuron origin of schizophrenia pathophysiology. *Neuropharmacology* 2012; 62: 1574–83.
- 51 Gavin DP, Sharma RP. Histone modifications, DNA methylation, and schizophrenia. *Neurosci Biobehav Rev* 2010; 34: 882–8.
- 52 D'Arcangelo G. Reelin in the Years: controlling neuronal migration and maturation in the mammalian brain. *Adv Neurosci* 2014; 2014: 1–19.
- 53 Shrestha S, Offer SM. Epigenetic regulations of GABAergic neurotransmission: relevance for neurological disorders and epigenetic therapy. *Med Epigenet* 2016; 4: 1–19.
- 54 Costa E, Chen Y, Davis J, Dong E, Noh JS, Tremolizzo L, et al. REELIN and schizophrenia: a disease at the interface of the genome and the epigenome. *Mol Interv* 2002; 2: 47–57.
- 55 Collinson N, Kuenzi FM, Jarolimek W, Maubach KA, Cothliff R, Sur C, et al. Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the alpha 5 subunit of the GABA_A receptor. *J Neurosci* 2002; 22: 5572–80.
- 56 Rudolph U, Knoflach F. Beyond classical benzodiazepines: novel therapeutic potential of GABA_A receptor subtypes. *Nat Rev Drug Discov* 2011; 10: 685–97.
- 57 Clayton T, Poe MM, Rallapalli S, Biawat P, Savic MM, Rowlett JK, et al. A review of the updated pharmacophore for the Alpha 5 GABA(A)

- benzodiazepine receptor model. *Int J Med Chem* 2015; 2015: 430248.
- 58 Savic MM, Huang S, Furtmuller R, Clayton T, Huck S, Obradovic DI, *et al*. Are GABA_A receptors containing alpha5 subunits contributing to the sedative properties of benzodiazepine site agonists? *Neuropsychopharmacology* 2008; 33: 332–9.
- 59 Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, *et al*. Trace fear conditioning involves hippocampal alpha5 GABA(A) receptors. *Proc Natl Acad Sci U S A* 2002; 99: 8980–5.
- 60 Belelli D, Harrison NL, Maguire J, Macdonald RL, Walker MC, Cope DW. Extrasynaptic GABA_A receptors: form, pharmacology, and function. *J Neurosci* 2009; 29: 12757–63.
- 61 Vinkers CH, Mirza NR, Olivier B, Kahn RS. The inhibitory GABA system as a therapeutic target for cognitive symptoms in schizophrenia: investigational agents in the pipeline. *Expert Opin Investig Drugs* 2010; 19: 1217–33.
- 62 Brickley SG, Mody I. Extrasynaptic GABA_A receptors: their function in the CNS and implications for disease. *Neuron* 2012; 73: 23–34.
- 63 Pouille F, Scanziani M. Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* 2001; 293: 1159–63.
- 64 Cobb SR, Buhl EH, Halasy K, Paulsen O, Somogyi P. Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* 1995; 378: 75–8.
- 65 Mody I, Pearce RA. Diversity of inhibitory neurotransmission through GABA(A) receptors. *Trends Neurosci* 2004; 27: 569–75.
- 66 Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci* 2005; 6: 215–29.
- 67 Volz A, Khorsand V, Gillies D, Leucht S. Benzodiazepines for schizophrenia. *Cochrane Database Syst Rev* 2007: CD006391.
- 68 Kehne JH, Maynard GD. GABA and schizophrenia. In: *Targets and emerging therapies for schizophrenia*. Albert JS, Wood MW, editors. USA: John Wiley & Sons, Inc; 2012.
- 69 Makaron L, Moran CA, Namjoshi O, Rallapalli S, Cook JM, Rowlett JK. Cognition-impairing effects of benzodiazepine-type drugs: role of GABA_A receptor subtypes in an executive function task in rhesus monkeys. *Pharmacol Biochem Behav* 2013; 104: 62–8.
- 70 Mohler H. Molecular regulation of cognitive functions and developmental plasticity: impact of GABA_A receptors. *J Neurochem* 2007; 102: 1–12.
- 71 Yee BK, Keist R, von Boehmer L, Studer R, Benke D, Hagenbuch N, *et al*. A schizophrenia-related sensorimotor deficit links alpha 3-containing GABA_A receptors to a dopamine hyperfunction. *Proc Natl Acad Sci U S A* 2005; 102: 17154–9.
- 72 Hauser J, Rudolph U, Keist R, Möhler H, Feldon J, Yee BK. Hippocampal alpha5 subunit-containing GABA_A receptors modulate the expression of prepulse inhibition. *Mol Psychiatry* 2005; 10: 201–7.
- 73 Gerdjikov TV, Rudolph U, Keist R, Mohler H, Feldon J, Yee BK. Hippocampal alpha 5 subunit-containing GABA A receptors are involved in the development of the latent inhibition effect. *Neurobiol Learn Mem* 2008; 89: 87–94.
- 74 Nutt D. GABA_A receptors: subtypes, regional distribution, and function. *J Clin Sleep Med* 2006; 2: S7–11.
- 75 Rudolph U, Mohler H. Analysis of GABA_A receptor function and dissection of the pharmacology of benzodiazepines and general anesthetics through mouse genetics. *Annu Rev Pharmacol Toxicol* 2004; 44: 475–98.
- 76 Bailey DJ, Tetzlaff JE, Cook JM, He X, Helmstetter FJ. Effects of hippocampal injections of a novel ligand selective for the alpha 5 beta 2 gamma 2 subunits of the GABA/benzodiazepine receptor on Pavlovian conditioning. *Neurobiol Learn Mem* 2002; 78: 1–10.
- 77 Glykys J, Mann EO, Mody I. Which GABA(A) receptor subunits are necessary for tonic inhibition in the hippocampus? *J Neurosci* 2008; 28: 1421–6.
- 78 Towers SK, Gloveli T, Traub RD, Driver JE, Engel D, Fradley R, *et al*. Alpha 5 subunit-containing GABA_A receptors affect the dynamic range of mouse hippocampal kainate-induced gamma frequency oscillations *in vitro*. *J Physiol* 2004; 559: 721–8.
- 79 Atack JR, Bayley PJ, Seabrook GR, Wafford KA, McKernan RM, Dawson GR. L-655,708 enhances cognition in rats but is not proconvulsant at a dose selective for alpha5-containing GABA_A receptors. *Neuropharmacology* 2006; 51: 1023–9.
- 80 Chambers MS, Atack JR, Broughton HB, Collinson N, Cook S, Dawson GR, *et al*. Identification of a novel, selective GABA(A) alpha5 receptor inverse agonist which enhances cognition. *J Med Chem* 2003; 46: 2227–40.
- 81 Dorow R, Horowski R, Paschelke G, Amin M. Severe anxiety induced by FG 7142, a beta-carboline ligand for benzodiazepine receptors. *Lancet* 1983; 2: 98–9.
- 82 Petersen EN. DMCM: a potent convulsive benzodiazepine receptor ligand. *Eur J Pharmacol* 1983; 94: 117–24.
- 83 Savic MM, Obradovic DI, Ugresic ND, Cook JM, Sarma PV, Bokonjic DR. Bidirectional effects of benzodiazepine binding site ligands on active avoidance acquisition and retention: differential antagonism by flumazenil and beta-Cct. *Psychopharmacology (Berl)* 2005; 180: 455–65.
- 84 Ballard TM, Knoflach F, Prinssen E, Borroni E, Vivian JA, Basile J, *et al*. RO4938581, a novel cognitive enhancer acting at GABA_A alpha5 subunit-containing receptors. *Psychopharmacology (Berl)* 2009; 202: 207–23.
- 85 Knust H, Achermann G, Ballard T, Buettelmann B, Gasser R, Fischer H, *et al*. The discovery and unique pharmacological profile of RO4938581 and RO4882224 as potent and selective GABA_A alpha5 inverse agonists for the treatment of cognitive dysfunction. *Bioorg Med Chem Lett* 2009; 19: 5940–4.
- 86 Redrobe JP, Elster L, Frederiksen K, Bundgaard C, de Jong IE, Smith GP, *et al*. Negative modulation of GABA_A alpha5 receptors by RO4938581 attenuates discrete sub-chronic and early postnatal phencyclidine (PCP)-induced cognitive deficits in rats. *Psychopharmacology (Berl)* 2012; 221: 451–68.
- 87 Green MF, Nuechterlein KH, Gold JM, Barch DM, Cohen J, Essock S, *et al*. Approaching a consensus cognitive battery for clinical trials in schizophrenia: the NIMH-MATRICES conference to select cognitive domains and test criteria. *Biol Psychiatry* 2004; 56: 301–7.
- 88 Barch DM, Braver TS, Carter CS, Poldrack RA, Robbins TW. CNTRICS final task selection: executive control. *Schizophr Bull* 2009; 35: 115–35.
- 89 Nuechterlein KH, Green MF, Kern RS, Baade LE, Barch DM, Cohen JD, *et al*. The MATRICES Consensus Cognitive Battery, part 1: test selection, reliability, and validity. *Am J Psychiatry* 2008; 165: 203–13.
- 90 Young JW, Powell SB, Risbrough V, Marston HM, Geyer MA. Using the MATRICES to guide development of a preclinical cognitive test battery for research in schizophrenia. *Pharmacol Ther* 2009; 122: 150–202.
- 91 Benejam B. Dementia symptoms in Down Syndrome. *Int Med Rev Down Syndrome* 2009; 13: 18–21.
- 92 Atack JR. GABA_A receptor subtype-selective modulators. II. alpha5-selective inverse agonists for cognition enhancement. *Curr Top Med Chem* 2011; 11: 1203–14.

- 93 Savic MM, Clayton T, Furtmuller R, Gavrilovic I, Samardzic J, Savic S, et al. PWZ-029, a compound with moderate inverse agonist functional selectivity at GABA(A) receptors containing alpha5 subunits, improves passive, but not active, avoidance learning in rats. *Brain Res* 2008; 1208: 150–9.
- 94 Harris D, Clayton T, Cook J, Sahbaie P, Halliwell RF, Furtmuller R, et al. Selective influence on contextual memory: physiochemical properties associated with selectivity of benzodiazepine ligands at GABA_A receptors containing the alpha5 subunit. *J Med Chem* 2008; 51: 3788–803.
- 95 Milic M, Timic T, Joksimovic S, Biawat P, Rallapalli S, Divljakovic J, et al. PWZ-029, an inverse agonist selective for alpha(5) GABA_A receptors, improves object recognition, but not water-maze memory in normal and scopolamine-treated rats. *Behav Brain Res* 2013; 241: 206–13.
- 96 Timic Stamenic T, Joksimovic S, Biawat P, Stankovic T, Markovic B, Cook JM, et al. Negative modulation of alpha(5) GABA_A receptors in rats may partially prevent memory impairment induced by MK-801, but not amphetamine- or MK-801-elicited hyperlocomotion. *J Psychopharmacol* 2015; 29: 1013–24.
- 97 Wong AH, Josselyn SA. Caution when diagnosing your mouse With schizophrenia: the use and misuse of model animals for understanding psychiatric disorders. *Biol Psychiatry* 2016; 79: 32–8.
- 98 Atack JR. Preclinical and clinical pharmacology of the GABA_A receptor alpha5 subtype-selective inverse agonist alpha5IA. *Pharmacol Ther* 2010; 125: 11–26.
- 99 Collinson N, Atack JR, Laughton P, Dawson GR, Stephens DN. An inverse agonist selective for alpha5 subunit-containing GABA_A receptors improves encoding and recall but not consolidation in the Morris water maze. *Psychopharmacology (Berl)* 2006; 188: 619–28.
- 100 Nutt DJ, Besson M, Wilson SJ, Dawson GR, Lingford-Hughes AR. Blockade of alcohol's amnesic activity in humans by an alpha5 subtype benzodiazepine receptor inverse agonist. *Neuropharmacology* 2007; 53: 810–20.
- 101 Atack JR, Maubach KA, Wafford KA, O'Connor D, Rodrigues AD, Evans DC, et al. *In vitro* and *in vivo* properties of 3-tert-butyl-7-(5-methylisoxazol-3-yl)-2-(1-methyl-1H-1,2,4-triazol-5-ylmethoxy)-pyrazolo[1,5-d][1,2,4]triazine (MRK-016), a GABA_A receptor alpha5 subtype-selective inverse agonist. *J Pharmacol Exp Ther* 2009; 331: 470–84.
- 102 Beinat C, Banister SD, Herrera M, Law V, Kassiou M. The therapeutic potential of alpha7 nicotinic acetylcholine receptor (alpha7 nAChR) agonists for the treatment of the cognitive deficits associated with schizophrenia. *CNS Drugs* 2015; 29: 529–42.
- 103 Stefansson H, Rujescu D, Cichon S, Pietilainen OP, Ingason A, Steinberg S, et al. Large recurrent microdeletions associated with schizophrenia. *Nature* 2008; 455: 232–6.
- 104 Leonard S, Gault J, Hopkins J, Logel J, Vianzon R, Short M, et al. Association of promoter variants in the alpha7 nicotinic acetylcholine receptor subunit gene with an inhibitory deficit found in schizophrenia. *Arch Gen Psychiatry* 2002; 59: 1085–96.
- 105 Yadon CA, Bugg JM, Kiskey MA, Davalos DB. P50 sensory gating is related to performance on select tasks of cognitive inhibition. *Cogn Affect Behav Neurosci* 2009; 9: 448–58.
- 106 Smith RC, Warner-Cohen J, Matute M, Butler E, Kelly E, Vaidyanathaswamy S, et al. Effects of nicotine nasal spray on cognitive function in schizophrenia. *Neuropsychopharmacology* 2006; 31: 637–43.
- 107 Brunzell DH, McIntosh JM. Alpha7 nicotinic acetylcholine receptors modulate motivation to self-administer nicotine: implications for smoking and schizophrenia. *Neuropsychopharmacology* 2012; 37: 1134–43.
- 108 Levin ED, McClernon FJ, Rezvani AH. Nicotinic effects on cognitive function: behavioral characterization, pharmacological specification, and anatomic localization. *Psychopharmacology (Berl)* 2006; 184: 523–39.
- 109 Wanaverbecq N, Semyanov A, Pavlov I, Walker MC, Kullmann DM. Cholinergic axons modulate GABAergic signaling among hippocampal interneurons via postsynaptic alpha 7 nicotinic receptors. *J Neurosci* 2007; 27: 5683–93.
- 110 McKay BE, Placzek AN, Dani JA. Regulation of synaptic transmission and plasticity by neuronal nicotinic acetylcholine receptors. *Biochem Pharmacol* 2007; 74: 1120–33.
- 111 Johnstone TB, Gu Z, Yoshimura RF, Villegier AS, Hogenkamp DJ, Whittemore ER, et al. Allosteric modulation of related ligand-gated ion channels synergistically induces long-term potentiation in the hippocampus and enhances cognition. *J Pharmacol Exp Ther* 2011; 336: 908–15.
- 112 Bari A, Dalley JW, Robbins TW. The application of the 5-choice serial reaction time task for the assessment of visual attentional processes and impulse control in rats. *Nat Protoc* 2008; 3: 759–67.
- 113 Buccafusco JJ. Multifunctional receptor-directed drugs for disorders of the central nervous system. *Neurotherapeutics* 2009; 6: 4–13.
- 114 Lodge DJ, Grace AA. Hippocampal dysregulation of dopamine system function and the pathophysiology of schizophrenia. *Trends Pharmacol Sci* 2011; 32: 507–13.
- 115 Gill KM, Grace AA. The role of alpha5 GABA_A receptor agonists in the treatment of cognitive deficits in schizophrenia. *Curr Pharm Des* 2014; 20: 5069–76.
- 116 Wilson IA, Ikonen S, Gallagher M, Eichenbaum H, Tanila H. Age-associated alterations of hippocampal place cells are subregion specific. *J Neurosci* 2005; 25: 6877–86.
- 117 Gill KM, Lodge DJ, Cook JM, Aras S, Grace AA. A novel alpha5GABA(A) R-positive allosteric modulator reverses hyperactivation of the dopamine system in the MAM model of schizophrenia. *Neuropsychopharmacology* 2011; 36: 1903–11.
- 118 Soto PL, Ator NA, Rallapalli SK, Biawat P, Clayton T, Cook JM, et al. Allosteric modulation of GABA_A receptor subtypes: effects on visual recognition and visuospatial working memory in rhesus monkeys [corrected]. *Neuropsychopharmacology* 2013; 38: 2315–25.
- 119 Richetto J, Labouesse MA, Poe MM, Cook JM, Grace AA, Riva MA, et al. Behavioral effects of the benzodiazepine-positive allosteric modulator SH-053-2'F-S-CH(3) in an immune-mediated neurodevelopmental disruption model. *Int J Neuropsychopharmacol* 2015; 18: pii: pyu055.
- 120 Engin E, Liu J, Rudolph U. alpha2-containing GABA(A) receptors: a target for the development of novel treatment strategies for CNS disorders. *Pharmacol Ther* 2012; 136: 142–52.
- 121 Hines RM, Hines DJ, Houston CM, Mukherjee J, Haydon PG, Tretter V, et al. Disrupting the clustering of GABA_A receptor alpha2 subunits in the frontal cortex leads to reduced gamma-power and cognitive deficits. *Proc Natl Acad Sci U S A* 2013; 110: 16628–33.
- 122 Lewis DA. The chandelier neuron in schizophrenia. *Dev Neurobiol* 2011; 71: 118–27.
- 123 Bloomfield C, French SJ, Jones DN, Reavill C, Southam E, Cilia J, et al. Chandelier cartridges in the prefrontal cortex are reduced in isolation reared rats. *Synapse* 2008; 62: 628–31.
- 124 Volk DW, Pierri JN, Fritschy JM, Auh S, Sampson AR, Lewis DA. Reciprocal alterations in pre- and postsynaptic inhibitory markers at chandelier cell inputs to pyramidal neurons in schizophrenia. *Cereb Cortex* 2002; 12: 1063–70.

- 125 Lewis DA, Volk DW, Hashimoto T. Selective alterations in prefrontal cortical GABA neurotransmission in schizophrenia: a novel target for the treatment of working memory dysfunction. *Psychopharmacology (Berl)* 2004; 174: 143–50.
- 126 Lewis DA, Hashimoto T, Volk DW. Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 2005; 6: 312–24.
- 127 Lewis DA, Cho RY, Carter CS, Eklund K, Forster S, Kelly MA, *et al*. Subunit-selective modulation of GABA type A receptor neurotransmission and cognition in schizophrenia. *Am J Psychiatry* 2008; 165: 1585–93.
- 128 Buchanan RW, Keefe RS, Lieberman JA, Barch DM, Csernansky JG, Goff DC, *et al*. A randomized clinical trial of MK-0777 for the treatment of cognitive impairments in people with schizophrenia. *Biol Psychiatry* 2011; 69: 442–9.
- 129 Whittington MA, Cunningham MO, LeBeau FE, Racca C, Traub RD. Multiple origins of the cortical gamma rhythm. *Dev Neurobiol* 2011; 71: 92–106.
- 130 Lewis DA, Fish KN, Arion D, Gonzalez-Burgos G. Perisomatic inhibition and cortical circuit dysfunction in schizophrenia. *Curr Opin Neurobiol* 2011; 21: 866–72.
- 131 Klausberger T, Somogyi P. Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* 2008; 321: 53–7.
- 132 Gonzalez-Burgos G. GABA transporter GAT1: a crucial determinant of GABA_B receptor activation in cortical circuits? *Adv Pharmacol* 2010; 58: 175–204.
- 133 Wang Y, Neubauer FB, Luscher HR, Thurley K. GABA_B receptor-dependent modulation of network activity in the rat prefrontal cortex *in vitro*. *Eur J Neurosci* 2010; 31: 1582–94.
- 134 Gaiarsa JL, Porcher C. Emerging neurotrophic role of GABA_B receptors in neuronal circuit development. *Front Cell Neurosci* 2013; 7: 206.
- 135 Lei S, McBain CJ. GABA_B receptor modulation of excitatory and inhibitory synaptic transmission onto rat CA3 hippocampal interneurons. *J Physiol* 2003; 546: 439–53.
- 136 Ghose S, Winter MK, McCarron KE, Tamminga CA, Enna SJ. The GABA_B receptor as a target for antidepressant drug action. *Br J Pharmacol* 2011; 162: 1–17.
- 137 Kantrowitz J, Citrome L, Javitt D. GABA(B) receptors, schizophrenia and sleep dysfunction: a review of the relationship and its potential clinical and therapeutic implications. *CNS Drugs* 2009; 23: 681–91.
- 138 Fatemi SH, Folsom TD, Thuras PD. GABA_A and GABA_B receptor dysregulation in superior frontal cortex of subjects with schizophrenia and bipolar disorder. *Synapse* 2017; 71: doi: 10.1002/syn.21973.
- 139 Fatemi SH, Folsom TD, Thuras PD. Deficits in GABA(B) receptor system in schizophrenia and mood disorders: a postmortem study. *Schizophr Res* 2011; 128: 37–43.
- 140 Mizukami K, Sasaki M, Ishikawa M, Iwakiri M, Hidaka S, Shiraishi H, *et al*. Immunohistochemical localization of gamma-aminobutyric acid(B) receptor in the hippocampus of subjects with schizophrenia. *Neurosci Lett* 2000; 283: 101–4.
- 141 Mizukami K, Ishikawa M, Hidaka S, Iwakiri M, Sasaki M, Iritani S. Immunohistochemical localization of GABA_B receptor in the entorhinal cortex and inferior temporal cortex of schizophrenic brain. *Prog Neuropsychopharmacol Biol Psychiatry* 2002; 26: 393–6.
- 142 Schultz H, Sommer T, Peters J. The role of the human entorhinal cortex in a representational account of memory. *Front Hum Neurosci* 2015; 9: 628.
- 143 Deng PY, Xiao Z, Yang C, Rojanathammanee L, Grisanti L, Watt J, *et al*. GABA_B receptor activation inhibits neuronal excitability and spatial learning in the entorhinal cortex by activating TREK-2 K⁺ channels. *Neuron* 2009; 63: 230–43.
- 144 Heaney CF, Kinney JW. Role of GABA_B receptors in learning and memory and neurological disorders. *Neurosci Biobehav Rev* 2016; 63: 1–28.
- 145 Iqbal F, Gillani QA, GABA_B receptor antagonists as cognition enhancers, in GABA_B receptor. Colombo G, Editor. USA: Humana Press; 2016.
- 146 Selten MM, Meyer F, Ba W, Valles A, Maas DA, Negwer M, *et al*. Increased GABA_B receptor signaling in a rat model for schizophrenia. *Sci Rep* 2016; 6: 34240.
- 147 Kleschevnikov AM, Belichenko PV, Faizi M, Jacobs LF, Htun K, Shamloo M, *et al*. Deficits in cognition and synaptic plasticity in a mouse model of Down syndrome ameliorated by GABA_B receptor antagonists. *J Neurosci* 2012; 32: 9217–27.
- 148 Liu QY, Wang CY, Cai ZL, Xu ST, Liu WX, Xiao P, *et al*. Effects of intrahippocampal GABA_B receptor antagonist treatment on the behavioral long-term potentiation and Y-maze learning performance. *Neurobiol Learn Mem* 2014; 114: 26–31.
- 149 Johnson NW, Ozkan M, Burgess AP, Prokic EJ, Wafford KA, O'Neill MJ, *et al*. Phase-amplitude coupled persistent theta and gamma oscillations in rat primary motor cortex *in vitro*. *Neuropharmacology* 2017; 119: 141–56.
- 150 Marrosu F, Santoni F, Fa M, Puligheddu M, Barberini L, Genugu F, *et al*. Beta and gamma range EEG power-spectrum correlation with spiking discharges in DBA/2J mice absence model: role of GABA receptors. *Epilepsia* 2006; 47: 489–94.
- 151 Froestl W, Gallagher M, Jenkins H, Madrid A, Melcher T, Teichman S, *et al*. SGS742: the first GABA_B receptor antagonist in clinical trials. *Biochem Pharmacol* 2004; 68: 1479–87.
- 152 Bianchi M, Panerai AE. Reversal of scopolamine-induced amnesia by the GABA_B receptor antagonist CGP 35348 in the mouse. *Brain Res Cogn Brain Res* 1993; 1: 135–6.
- 153 Farr SA, Flood JF, Morley JE. The effect of cholinergic, GABAergic, serotonergic, and glutamatergic receptor modulation on posttrial memory processing in the hippocampus. *Neurobiol Learn Mem* 2000; 73: 150–67.
- 154 Pitsikas N, Rigamonti AE, Cella SG, Muller EE. The GABA_B receptor and recognition memory: possible modulation of its behavioral effects by the nitroergic system. *Neuroscience* 2003; 118: 1121–7.
- 155 Kumar K, Kaur H, Deshmukh R. Neuroprotective role of GABA_B receptor modulation against streptozotocin-induced behavioral and biochemical abnormalities in rats. *Neuroscience* 2017; 357: 67–74.
- 156 Gillani Q, Ali M, Iqbal F. CGP 35348, GABA B receptor antagonist, has a potential to improve neuromuscular coordination and spatial learning in albino mouse following neonatal brain damage. *Biomed Res Int* 2014; 2014: 295215.
- 157 Gillani Q, Iqbal S, Arfa F, Khakwani S, Akbar A, Ullah A, *et al*. Effect of GABA_B receptor antagonist (CGP35348) on learning and memory in albino mice. *ScientificWorldJournal* 2014; 2014: 983651.
- 158 Wu N, Wang F, Jin Z, Zhang Z, Wang LK, Zhang C, *et al*. Effects of GABA_B receptors in the insula on recognition memory observed with intelligence. *Behav Brain Funct* 2017; 13: 7.
- 159 Mondadori C, Jaekel J, Preiswerk G. CGP 36742: the first orally active GABA_B blocker improves the cognitive performance of mice, rats, and rhesus monkeys. *Behav Neural Biol* 1993; 60: 62–8.
- 160 Mondadori C, Mobius HJ, Borkowski J. The GABA_B receptor antagonist CGP 36,742 and the nootropic oxiracetam facilitate the formation of long-term memory. *Behav Brain Res* 1996; 77: 223–5.
- 161 Mondadori C, Moebius HJ, Zingg M. CGP 36,742, an orally active GABA_B receptor antagonist, facilitates memory in a social recognition test in rats. *Behav Brain Res* 1996; 77: 227–9.

- 162 Carletti R LV, Bowery NG. The GABA_B antagonist CGP36742 enhances spatial learning performance and antagonises baclofen induced amnesia in mice. *Br J Pharmacol* 1993; 109: 74.
- 163 Nakagawa Y, Takashima T. The GABA_B receptor antagonist CGP36742 attenuates the baclofen- and scopolamine-induced deficit in Morris water maze task in rats. *Brain Res* 1997; 766: 101–6.
- 164 Getova D, Bowery NG, Spassov V. Effects of GABA_B receptor antagonists on learning and memory retention in a rat model of absence epilepsy. *Eur J Pharmacol* 1997; 320: 9–13.
- 165 Helm KA, Haberman RP, Dean SL, Hoyt EC, Melcher T, Lund PK, *et al*. GABA_B receptor antagonist SGS742 improves spatial memory and reduces protein binding to the cAMP response element (CRE) in the hippocampus. *Neuropharmacology* 2005; 48: 956–64.
- 166 Sabbagh MN. Drug development for Alzheimer's disease: where are we now and where are we headed? *Am J Geriatr Pharmacother* 2009; 7: 167–85.
- 167 Getova D, Bowery NG. The modulatory effects of high affinity GABA(B) receptor antagonists in an active avoidance learning paradigm in rats. *Psychopharmacology (Berl)* 1998; 137: 369–73.
- 168 Getova DP, Bowery NG. Effects of high-affinity GABA_B receptor antagonists on active and passive avoidance responding in rodents with gamma-hydroxybutyrolactone-induced absence syndrome. *Psychopharmacology (Berl)* 2001; 157: 89–95.
- 169 Davies CH, Pozza MF, Collingridge GL. CGP 55845A: a potent antagonist of GABA_B receptors in the CA1 region of rat hippocampus. *Neuropharmacology* 1993; 32: 1071–3.
- 170 Lasarge CL, Banuelos C, Mayse JD, Bizon JL. Blockade of GABA_B receptors completely reverses age-related learning impairment. *Neuroscience* 2009; 164: 941–7.
- 171 Gillani QA, Akbar A, Ali M, Iqbal F. Gender-specific effects of CGP 55845, GABA_B receptor antagonist, on neuromuscular coordination, learning and memory formation in albino mouse following neonatal hypoxia-ischemia insult. *Neurol Sci* 2015; 36: 961–9.
- 172 Hirst WD, Babbs AJ, Green A, Minton JA, Shaw TE, Wise A, *et al*. Pharmacological characterisation of a cell line expressing GABA B1b and GABA B2 receptor subunits. *Biochem Pharmacol* 2003; 65: 1103–13.
- 173 Rotaru DC, Lewis DA, Gonzalez-Burgos G. The role of glutamatergic inputs onto parvalbumin-positive interneurons: relevance for schizophrenia. *Rev Neurosci* 2012; 23: 97–109.
- 174 Bowery NG. Baclofen: therapeutic use and potential of the prototypic GABA_B receptor agonist, in GABA_B receptor. Colombo G, Editor. Switzerland: Humana Press; 2016. p337–56.
- 175 Arai S, Takuma K, Mizoguchi H, Ibi D, Nagai T, Kamei H, *et al*. GABA_B receptor agonist baclofen improves methamphetamine-induced cognitive deficit in mice. *Eur J Pharmacol* 2009; 602: 101–4.
- 176 Luo P, Chen C, Lu Y, Fu T, Lu Q, Xu X, *et al*. Baclofen ameliorates spatial working memory impairments induced by chronic cerebral hypoperfusion via up-regulation of HCN2 expression in the PFC in rats. *Behav Brain Res* 2016; 308: 6–13.
- 177 Cryan JF, Kelly PH, Chaperon F, Gentsch C, Mombereau C, Lingenhoehl K, *et al*. Behavioral characterization of the novel GABA_B receptor-positive modulator GS39783 (*N,N'*-dicyclopentyl-2-methylsulfanyl-5-nitro-pyrimidine-4,6-diamine): anxiolytic-like activity without side effects associated with baclofen or benzodiazepines. *J Pharmacol Exp Ther* 2004; 310: 952–63.
- 178 Mugnaini C, Corelli F. The allosteric modulation of the GABA_B receptor: a medicinal chemistry perspective. In: GABA_B receptor. Colombo G, Editor. Switzerland: Humana Press; 2016. p33–52.
- 179 Costa E, Dong E, Grayson DR, Guidotti A, Ruzicka W, Veldic M. Reviewing the role of DNA (cytosine-5) methyltransferase overexpression in the cortical GABAergic dysfunction associated with psychosis vulnerability. *Epigenetics* 2007; 2: 29–36.
- 180 Tsankova N, Renthal W, Kumar A, Nestler EJ. Epigenetic regulation in psychiatric disorders. *Nat Rev Neurosci* 2007; 8: 355–67.
- 181 Akbarian S. The molecular pathology of schizophrenia—focus on histone and DNA modifications. *Brain Res Bull* 2010; 83: 103–7.
- 182 Szyf M. Prospects for the development of epigenetic drugs for CNS conditions. *Nat Rev Drug Discov* 2015; 14: 461–74.
- 183 Huang HS, Matevosian A, Whittle C, Kim SY, Schumacher A, Baker SP, *et al*. Prefrontal dysfunction in schizophrenia involves mixed-lineage leukemia 1-regulated histone methylation at GABAergic gene promoters. *J Neurosci* 2007; 27: 11254–62.
- 184 Bahari-Javan S, Varbanov H, Halder R, Benito E, Kaurani L, Burkhardt S, *et al*. HDAC1 links early life stress to schizophrenia-like phenotypes. *Proc Natl Acad Sci U S A* 2017; 114: E4686–E4694.
- 185 Graff J, Tsai LH. The potential of HDAC inhibitors as cognitive enhancers. *Annu Rev Pharmacol Toxicol* 2013; 53: 311–30.
- 186 Gottlicher M, Minucci S, Zhu P, Kramer OH, Schimpf A, Giavara S, *et al*. Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells. *EMBO J* 2001; 20: 6969–78.
- 187 Su P, Jiang AL, Liu F. Clozapine modulates the glycogen synthase kinase-3 signaling partly via GABA_B receptors. *Ann Biol Sci* 2017; 5: 14–20.
- 188 Hasan A, Mitchell A, Schneider A, Halene T, Akbarian S. Epigenetic dysregulation in schizophrenia: molecular and clinical aspects of histone deacetylase inhibitors. *Eur Arch Psychiatry Clin Neurosci* 2013; 263: 273–84.
- 189 Kilgore M, Miller CA, Fass DM, Hennig KM, Haggarty SJ, Sweatt JD, *et al*. Inhibitors of class 1 histone deacetylases reverse contextual memory deficits in a mouse model of Alzheimer's disease. *Neuropsychopharmacology* 2010; 35: 870–80.
- 190 Takuma K, Hara Y, Kataoka S, Kawanai T, Maeda Y, Watanabe R, *et al*. Chronic treatment with valproic acid or sodium butyrate attenuates novel object recognition deficits and hippocampal dendritic spine loss in a mouse model of autism. *Pharmacol Biochem Behav* 2014; 126: 43–9.
- 191 Hirsch E, Schmitz B, Carreno M. Epilepsy, antiepileptic drugs (AEDs) and cognition. *Acta Neurol Scand Suppl* 2003; 180: 23–32.
- 192 Montgomery RL, Hsieh J, Barbosa AC, Richardson JA, Olson EN. Histone deacetylases 1 and 2 control the progression of neural precursors to neurons during brain development. *Proc Natl Acad Sci U S A* 2009; 106: 7876–81.
- 193 Peixoto L, Abel T. The role of histone acetylation in memory formation and cognitive impairments. *Neuropsychopharmacology* 2013; 38: 62–76.
- 194 Bieszczad KM, Bechay K, Rusche JR, Jacques V, Kudugunti S, Miao W, *et al*. Histone deacetylase inhibition via RGFP966 releases the Brakes on sensory cortical plasticity and the specificity of memory formation. *J Neurosci* 2015; 35: 13124–32.
- 195 Fischer A, Sananbenesi F, Mungenast A, Tsai LH. Targeting the correct HDAC(s) to treat cognitive disorders. *Trends Pharmacol Sci* 2010; 31: 605–17.
- 196 Rumbaugh G, Sullivan SE, Ozkan ED, Rojas CS, Hubbs CR, Aceti M, *et al*. Pharmacological selectivity within class I histone deacetylases predicts effects on synaptic function and memory rescue. *Neuropsychopharmacology* 2015; 40: 2307–16.
- 197 Wagner FF, Zhang YL, Fass DM, Joseph N, Gale JP, Weiwer M, *et al*. kinetically selective inhibitors of histone deacetylase 2 (HDAC2) as cognition enhancers. *Chem Sci* 2015; 6: 804–815.
- 198 Bahari-Javan S, Sananbenesi F, Fischer A. Histone-acetylation: a link between Alzheimer's disease and post-traumatic stress disorder?

- Front Neurosci 2014; 8: 160.
- 199 Kim D, Frank CL, Dobbin MM, Tsunemoto RK, Tu W, Peng PL, et al. Deregulation of HDAC1 by p25/Cdk5 in neurotoxicity. *Neuron* 2008; 60: 803–17.
- 200 Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, et al. HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature* 2009; 459: 55–60.
- 201 Whittle N, Singewald N. HDAC inhibitors as cognitive enhancers in fear, anxiety and trauma therapy: where do we stand? *Biochem Soc Trans* 2014; 42: 569–81.
- 202 Bredy TW, Barad M. The histone deacetylase inhibitor valproic acid enhances acquisition, extinction, and reconsolidation of conditioned fear. *Learn Mem* 2008; 15: 39–45.
- 203 Camp MC, Macpherson KP, Lederle L, Graybeal C, Gaburro S, DeBrouse LM, et al. Genetic strain differences in learned fear inhibition associated with variation in neuroendocrine, autonomic, and amygdala dendritic phenotypes. *Neuropsychopharmacology* 2012; 37: 1534–47.
- 204 Whittle N, Schmuckermair C, Gunduz Cinar O, Hauschild M, Ferraguti F, Holmes A, et al. Deep brain stimulation, histone deacetylase inhibitors and glutamatergic drugs rescue resistance to fear extinction in a genetic mouse model. *Neuropharmacology* 2013; 64: 414–23.
- 205 Wei J, Xiong Z, Lee JB, Cheng J, Duffney LJ, Matas E, et al. Histone modification of Nedd4 ubiquitin ligase controls the loss of AMPA receptors and cognitive impairment induced by repeated stress. *J Neurosci* 2016; 36: 2119–30.
- 206 Fass DM, Reis SA, Ghosh B, Hennig KM, Joseph NF, Zhao WN, et al. Crebinostat: a novel cognitive enhancer that inhibits histone deacetylase activity and modulates chromatin-mediated neuroplasticity. *Neuropharmacology* 2013; 64: 81–96.
- 207 Zhang L, Liu C, Wu J, Tao JJ, Sui XL, Yao ZG, et al. Tubastatin A/ACY-1215 improves cognition in Alzheimer's disease transgenic mice. *J Alzheimers Dis* 2014; 41: 1193–205.
- 208 Graff J, Joseph NF, Horn ME, Samiei A, Meng J, Seo J, et al. Epigenetic priming of memory updating during reconsolidation to attenuate remote fear memories. *Cell* 2014; 156: 261–76.
- 209 Hanson JE, La H, Plise E, Chen YH, Ding X, Hanania T, et al. SAHA enhances synaptic function and plasticity *in vitro* but has limited brain availability *in vivo* and does not impact cognition. *PLoS One* 2013; 8: e69964.
- 210 Alarcon JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER, et al. Chromatin acetylation, memory, and LTP are impaired in CBP^{-/-} mice: a model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron* 2004; 42: 947–59.
- 211 Vecsey CG, Hawk JD, Lattal KM, Stein JM, Fabian SA, Attner MA, et al. Histone deacetylase inhibitors enhance memory and synaptic plasticity via CREB:CBP-dependent transcriptional activation. *J Neurosci* 2007; 27: 6128–40.
- 212 Sharma S, Taliyan R, Ramagiri S. Histone deacetylase inhibitor, trichostatin A, improves learning and memory in high-fat diet-induced cognitive deficits in mice. *J Mol Neurosci* 2015; 56: 1–11.
- 213 Sung YM, Lee T, Yoon H, DiBattista AM, Song JM, Sohn Y, et al. Mercaptoacetamide-based class II HDAC inhibitor lowers Abeta levels and improves learning and memory in a mouse model of Alzheimer's disease. *Exp Neurol* 2013; 239: 192–201.
- 214 Govindarajan N, Agis-Balboa RC, Walter J, Sananbenesi F, Fischer A. Sodium butyrate improves memory function in an Alzheimer's disease mouse model when administered at an advanced stage of disease progression. *J Alzheimers Dis* 2011; 26: 187–97.
- 215 Reolon GK, Maurmann N, Werenicz A, Garcia VA, Schroder N, Wood MA, et al. Posttraining systemic administration of the histone deacetylase inhibitor sodium butyrate ameliorates aging-related memory decline in rats. *Behav Brain Res* 2011; 221: 329–32.
- 216 Fischer A, Sananbenesi F, Wang X, Dobbin M, Tsai LH. Recovery of learning and memory is associated with chromatin remodelling. *Nature* 2007; 447: 178–82.
- 217 Peleg S, Sananbenesi F, Zovoilis A, Burkhardt S, Bahari-Javan S, Agis-Balboa RC, et al. Altered histone acetylation is associated with age-dependent memory impairment in mice. *Science* 2010; 328: 753–6.
- 218 Rane P, Shields J, Heffernan M, Guo Y, Akbarian S, King JA. The histone deacetylase inhibitor, sodium butyrate, alleviates cognitive deficits in pre-motor stage PD. *Neuropharmacology* 2012; 62: 2409–12.
- 219 Valvassori SS, Varela RB, Arent CO, Dal-Pont GC, Bobsin TS, Budni J, et al. Sodium butyrate functions as an antidepressant and improves cognition with enhanced neurotrophic expression in models of maternal deprivation and chronic mild stress. *Curr Neurovasc Res* 2014; 11: 359–66.
- 220 Sandner G, Host L, Angst MJ, Guiberteau T, Guignard B, Zwiller J. The HDAC inhibitor phenylbutyrate reverses effects of neonatal ventral hippocampal lesion in rats. *Front Psychiatry* 2011; 1: 153.
- 221 Ricobaraza A, Cuadrado-Tejedor M, Perez-Mediavilla A, Frechilla D, Del Rio J, Garcia-Osta A. Phenylbutyrate ameliorates cognitive deficit and reduces tau pathology in an Alzheimer's disease mouse model. *Neuropsychopharmacology* 2009; 34: 1721–32.
- 222 Wiley JC, Pettan-Brewer C, Ladiges WC. Phenylbutyric acid reduces amyloid plaques and rescues cognitive behavior in AD transgenic mice. *Aging Cell* 2011; 10: 418–28.
- 223 Gardian G, Browne SE, Choi DK, Klivenyi P, Gregorio J, Kubilus JK, et al. Neuroprotective effects of phenylbutyrate in the N171-82Q transgenic mouse model of Huntington's disease. *J Biol Chem* 2005; 280: 556–63.
- 224 Zhou W, Bercury K, Cumiskey J, Luong N, Lebin J, Freed CR. Phenylbutyrate up-regulates the DJ-1 protein and protects neurons in cell culture and in animal models of Parkinson disease. *J Biol Chem* 2011; 286: 14941–51.
- 225 Ristic AJ, Vojvodic N, Jankovic S, Sindelic A, Sokic D. The frequency of reversible parkinsonism and cognitive decline associated with valproate treatment: a study of 364 patients with different types of epilepsy. *Epilepsia* 2006; 47: 2183–5.
- 226 Nicolai J, Aldenkamp AP, Huizenga JR, Teune LK, Brouwer OF. Cognitive side effects of valproic acid-induced hyperammonemia in children with epilepsy. *J Clin Psychopharmacol* 2007; 27: 221–4.
- 227 Shah RD, Jagtap JC, Mruthyunjaya S, Shelke GV, Pujari R, Das G, et al. Sodium valproate potentiates staurosporine-induced apoptosis in neuroblastoma cells via Akt/survivin independently of HDAC inhibition. *J Cell Biochem* 2013; 114: 854–63.
- 228 Wu P, Hong S, Zhong M, Guo Y, Chen H, Jiang L. Effect of sodium valproate on cognitive function and hippocampus of rats after convulsive status epilepticus. *Med Sci Monit* 2016; 22: 5197–205.
- 229 Bowers ME, Xia B, Carreiro S, Ressler KJ. The class I HDAC inhibitor RGFP963 enhances consolidation of cued fear extinction. *Learn Mem* 2015; 22: 225–31.
- 230 Suelves N, Kirkham-McCarthy L, Lahue RS, Gines S. A selective inhibitor of histone deacetylase 3 prevents cognitive deficits and suppresses striatal CAG repeat expansions in Huntington's disease mice. *Sci Rep* 2017; 7: 6082.
- 231 McQuown SC, Barrett RM, Matheos DP, Post RJ, Rogge GA, Alenghat T, et al. HDAC3 is a critical negative regulator of long-term memory formation. *J Neurosci* 2011; 31: 764–74.
- 232 Jia H, Kast RJ, Steffan JS, Thomas EA. Selective histone deacetylase

- (HDAC) inhibition imparts beneficial effects in Huntington's disease mice: implications for the ubiquitin-proteasomal and autophagy systems. *Hum Mol Genet* 2012; 21: 5280–93.
- 233 Castellano JF, Fletcher BR, Patzke H, Long JM, Sewal A, Kim DH, *et al*. Reassessing the effects of histone deacetylase inhibitors on hippocampal memory and cognitive aging. *Hippocampus* 2014; 24: 1006–16.
- 234 Green KN, Steffan JS, Martinez-Coria H, Sun X, Schreiber SS, Thompson LM, *et al*. Nicotinamide restores cognition in Alzheimer's disease transgenic mice via a mechanism involving sirtuin inhibition and selective reduction of Thr231-phosphotau. *J Neurosci* 2008; 28: 11500–10.
- 235 Naviaux JC, Schuchbauer MA, Li K, Wang L, Risbrough VB, Powell SB, *et al*. Reversal of autism-like behaviors and metabolism in adult mice with single-dose antipurinergic therapy. *Transl Psychiatry* 2014; 4: e400.
- 236 Naviaux JC, Wang L, Li K, Bright AT, Alaynick WA, Williams KR, *et al*. Antipurinergic therapy corrects the autism-like features in the Fragile X (*Fmr1* knockout) mouse model. *Mol Autism* 2015; 6: 1.
- 237 Naviaux RK, Curtis B, Li K, Naviaux JC, Bright AT, Reiner GE, *et al*. Low-dose suramin in autism spectrum disorder: a small, phase I/II, randomized clinical trial. *Ann Clin Transl Neurol* 2017; 4: 491–505.
- 238 Cuadrado-Tejedor M, Garcia-Barroso C, Sanchez-Arias JA, Rabal O, Perez-Gonzalez M, Mederos S, *et al*. A first-in-class small-molecule that acts as a dual inhibitor of HDAC and PDE5 and that rescues hippocampal synaptic impairment in Alzheimer's disease mice. *Neuropsychopharmacology* 2017; 42: 524–39.
- 239 Ganai SA, Ramadoss M, Mahadevan V. Histone deacetylase (HDAC) inhibitors—emerging roles in neuronal memory, learning, synaptic plasticity and neural regeneration. *Curr Neuropharmacol* 2016; 14: 55–71.
- 240 Kazantsev AG, Thompson LM. Therapeutic application of histone deacetylase inhibitors for central nervous system disorders. *Nat Rev Drug Discov* 2008; 7: 854–68.
- 241 Geffen Y, Keefe R, Rabinowitz J, Anand R, Davidson M. BL-1020, a new gamma-aminobutyric acid-enhanced antipsychotic: results of 6-week, randomized, double-blind, controlled, efficacy and safety study. *J Clin Psychiatry* 2012; 73: e1168–74.
- 242 Nudelman A, Gil-Ad I, Shpaysman N, Terasenko I, Ron H, Savitsky K, *et al*. A mutual prodrug ester of GABA and perphenazine exhibits antischizophrenic efficacy with diminished extrapyramidal effects. *J Med Chem* 2008; 51: 2858–62.
- 243 Geffen Y, Nudelman A, Gil-Ad I, Rephaeli A, Huang M, Savitsky K, *et al*. BL-1020: a novel antipsychotic drug with GABAergic activity and low catalepsy, is efficacious in a rat model of schizophrenia. *Eur Neuropsychopharmacol* 2009; 19: 1–13.
- 244 Fitzgerald PB. BL-1020, an oral antipsychotic agent that reduces dopamine activity and enhances GABA_A activity, for the treatment of schizophrenia. *Curr Opin Investig Drugs* 2010; 11: 92–100.
- 245 Conti F, Melone M, Fattorini G, Bragina L, Ciappelloni S. A role for GAT-1 in presynaptic GABA homeostasis? *Front Cell Neurosci* 2011; 5: 2.
- 246 Jensen K, Chiu CS, Sokolova I, Lester HA, Mody I. GABA transporter-1 (GAT1)-deficient mice: differential tonic activation of GABA_A versus GABA_B receptors in the hippocampus. *J Neurophysiol* 2003; 90: 2690–701.
- 247 Wilson C, Terry AV Jr. Neurodevelopmental animal models of schizophrenia: role in novel drug discovery and development. *Clin Schizophr Relat Psychoses* 2010; 4: 124–37.
- 248 Rideau Batista Novais A, Crouzin N, Cavalier M, Boubal M, Guiramand J, Cohen-Solal C, *et al*. Tiagabine improves hippocampal long-term depression in rat pups subjected to prenatal inflammation. *PLoS One* 2014; 9: e106302.
- 249 Aikia M, Jutila L, Salmenpera T, Mervaala E, Kalviainen R. Comparison of the cognitive effects of tiagabine and carbamazepine as monotherapy in newly diagnosed adult patients with partial epilepsy: pooled analysis of two long-term, randomized, follow-up studies. *Epilepsia* 2006; 47: 1121–7.
- 250 Salat K, Podkowa A, Mogilski S, Zareba P, Kulig K, Salat R, *et al*. The effect of GABA transporter 1 (GAT1) inhibitor, tiagabine, on scopolamine-induced memory impairments in mice. *Pharmacol Rep* 2015; 67: 1155–62.
- 251 Lopes da Silva F. EEG and MEG: relevance to neuroscience. *Neuron* 2013; 80: 1112–28.
- 252 Nithianantharajah J, Komiyama NH, McKechnie A, Johnstone M, Blackwood DH, St Clair D, *et al*. Synaptic scaffold evolution generated components of vertebrate cognitive complexity. *Nat Neurosci* 2013; 16: 16–24.
- 253 Wang DY, Kosowan J, Samsom J, Leung L, Zhang KL, Li YX, *et al*. Inhibition of the G9a/GLP histone methyltransferase modulates anxiety-related behavior in the mouse. *Acta Pharmacol Sin* 2018; 39. In press. doi: 10.1038/aps.2017.190.