

## ORIGINAL ARTICLE

## Coinjection of IL2 DNA enhances E7-specific antitumor immunity elicited by intravaginal therapeutic HPV DNA vaccination with electroporation

Y Sun<sup>1,2</sup>, S Peng<sup>2</sup>, A Yang<sup>2</sup>, E Farmer<sup>2</sup>, T-C Wu<sup>2,3,4,5</sup> and C-F Hung<sup>2,3</sup>

The generation and use of therapeutic human papillomavirus (HPV) DNA vaccines represent an appealing treatment method against HPV-associated cervical cancer owing to their safety and durability. Previously, we created a therapeutic HPV DNA vaccine candidate by linking the HPV16-E7 DNA sequence to calreticulin (CRT/E7), which we showed could generate significant E7-specific cytotoxic T lymphocyte (CTL)-mediated antitumor immune responses against HPV16 oncogenes expressing murine tumor model TC-1. Here we assess the therapeutic efficacy of intravaginal immunization with pcDNA3-CRT/E7 followed by electroporation. In addition, we examined whether coadministration of DNA-encoding interleukin 2 (IL2) with the pcDNA3-CRT/E7 could improve the T-cell responses elicited by pcDNA3-CRT/E7. TC-1 tumor-bearing mice vaccinated intravaginally with both pcDNA3-CRT/E7 and IL2 DNA followed by electroporation induced stronger local antitumor CTL response in comparison to mice that received other treatment regimens. Additionally, we found that coadministration of IL2 DNA with pcDNA3-CRT/E7 modified the tumor microenvironment by decreasing the population of regulatory T cells and myeloid-derived suppressor cells relative to that of CTLs. Our data demonstrate the translational potential of local administration of IL2 and pcDNA3-CRT/E7 followed by electroporation in treating cervicovaginal tumors.

*Gene Therapy* (2017) 24, 408–415; doi:10.1038/gt.2017.38

## INTRODUCTION

The generation and use of therapeutic human papillomavirus (HPV) DNA vaccines represents an appealing treatment method against HPV-associated cervical cancer owing to their safety and stability.<sup>1</sup> It has been discovered that the two oncoproteins encoded within the HPV genome, the early protein 6 (E6) and early protein 7 (E7), have vital roles in the tumorigenesis and upkeep of HPV-associated cervical cancer.<sup>2,3</sup> As such, the E6 and E7 oncoproteins are believed to be optimal targets for the generation of immunotherapeutic strategies against HPV-associated cancer because they are tumor specific, constantly expressed and are foreign proteins; therefore, they do not possess any issues regarding immune tolerance. In our previous studies, we have designed and generated a potent therapeutic HPV DNA vaccine composed of a fusion construct of HPV-16 E7 protein linked to calreticulin (CRT/E7), which we have shown to enhance the cross-presentation of the E7 antigen on to the major histocompatibility complex class I molecule.<sup>4,5</sup> Furthermore, we previously demonstrated that immunization with CRT/E7 DNA resulted in a potent cytotoxic T lymphocyte (CTL) immune response, which translates into a potent antitumor effect against TC-1, a preclinical HPV E7+ tumor model.<sup>6</sup> The promising data demonstrated in these preclinical studies have resulted in the evaluation of clinical grade CRT/E7 DNA vaccine in various clinical trials.<sup>7,8</sup>


Even though therapeutic DNA vaccines are a potentially efficacious cancer therapy, their immunogenicity is often times unimpressive due to the limited expression of the DNA-encoded protein *in vivo*. Electroporation aids in DNA plasmid delivery by creating an electrical field at the vaccination location.<sup>9</sup> This electrical field destabilizes the cell plasma membrane, inducing greater permeability and enabling more efficient passage of DNA plasmids into the target cells.<sup>10</sup> Furthermore, electroporation produces adjuvant-like effects, which enhance the immune response.<sup>9</sup> Previously, we have demonstrated that DNA immunization followed by electroporation elicited stronger antigen-specific CTL responses compared with intramuscular (IM) administration alone or gene gun vaccination.<sup>11</sup> In addition, the use of therapeutic HPV DNA vaccine with electroporation was demonstrated to be safe, well tolerated and potentially effective in humans.<sup>12</sup> These studies support the use of electroporation for the enhancement of DNA vaccine immunogenicity.

In addition to the method of therapeutic vaccination, the route of administration has also been identified to affect the location of optimal cell-mediated immune responses. It was previously demonstrated that local antitumor response is more important for tumor regression, compared with a systemic response only.<sup>13</sup> Disease clearance has been correlated with localized immune responses at the lesion site, suggesting the significance of

<sup>1</sup>Department of Obstetrics and Gynecology, Shanghai First People's Hospital, Shanghai Jiao Tong University, Shanghai, China; <sup>2</sup>Department of Pathology, Johns Hopkins Medical Institutions, Baltimore, MD, USA; <sup>3</sup>Department of Oncology, Johns Hopkins Medical Institutions, Baltimore, MD, USA; <sup>4</sup>Department of Molecular Microbiology and Immunology, Johns Hopkins Medical Institutions, Baltimore, MD, USA and <sup>5</sup>Department of Obstetrics and Gynecology, Johns Hopkins Medical Institutions, Baltimore, MD, USA. Correspondence: Dr T-C Wu, Departments of Pathology, Oncology, Molecular Microbiology and Immunology, and Obstetrics and Gynecology, The Johns Hopkins University School of Medicine, CRB II Room 309, 1550 Orleans Street, Baltimore, MD 21231, USA or Dr C-F Hung, Departments of Pathology and Oncology, The Johns Hopkins University School of Medicine, CRB II Room 307, 1550 Orleans Street, Baltimore, MD 21231, USA.

E-Mail: wutc@jhmi.edu or chung2@jhmi.edu

Received 6 December 2016; revised 16 February 2017; accepted 4 May 2017; accepted article preview online 11 May 2017; advance online publication, 1 June 2017



**Figure 1.** HPV DNA vaccine combined with IL2 intravaginal injection enhances the vaginal immune response. On day 0, 20  $\mu\text{g}$  of DNA constructs (pcDNA3-CRT/E7 and/or pcDNA3-IL2) were submucosally injected into the vagina followed by electroporation. Mice were boosted with the same regimen on day 7. **(a)** Representative flow cytometric analysis for the amount of E7-specific CD8<sup>+</sup> T cells in splenocytes. **(b)** Bar graph summary of the flow cytometric analysis. **(c)** Representative flow cytometric analysis for the amount of E7-specific CD8<sup>+</sup> T cells in the cervicovaginal tracts of mice. **(d)** Bar graph summary of the flow cytometric analysis. \* $P < 0.05$ ; \*\* $P < 0.01$ ; NS = not significant.

generating and targeting HPV-specific immune responses at the location of the lesion.<sup>14</sup> Intravaginal immunization with HPV pseudoviruses was found to induce local CTL responses and antitumor effects against cervicovaginal tumors.<sup>15</sup> We previously showed that immunization with a DNA vaccine, followed by a vaccinia virus vaccine, in the cervicovaginal region was able to stimulate an enhanced local, antigen-specific CTL response in TC-1 tumor-bearing mice when compared with mice that received IM vaccinations.<sup>16</sup> Likewise, intravaginal DNA immunization followed by electroporation enhanced E7-specific CTLs both systemically and locally and prolonged the survival of tumor-bearing mice in comparison to mice that received IM immunization followed by electroporation.<sup>17</sup>

Interleukin-2 (IL2) is a cytokine with numerous functions, including: stimulating T-cell proliferation, generating CTL and natural killer cells, and mediating B-cell proliferation and immunoglobulin synthesis.<sup>18</sup> Previously, we have successfully targeted DNA-encoding IL2 to the tumor location to provoke robust therapeutic antitumor effects in tumor-bearing mice.<sup>19</sup> Furthermore, IL2 was shown to strengthen the antigen-specific T-cell responses, specifically Th1 responses, generated by an HIV targeting DNA vaccine.<sup>20,21</sup> IL2 is currently permitted to be used in

patients with malignant melanoma and metastatic renal-cell carcinoma.<sup>18</sup> Therefore, administration of DNA-encoding IL2 may further enhance the therapeutic response produced by the CRT/E7 DNA vaccine.

Here we investigated the use of DNA-encoding IL2 to enhance the CTL responses and antitumor effects elicited by pcDNA3-CRT/E7 immunization in the cervicovaginal tract via electroporation. We established that coadministration of DNA-encoding IL2 notably enhanced the HPV antigen-specific CTL responses and antitumor effects induced by intravaginal injection of pcDNA3-CRT/E7 vaccine followed by electroporation. Additionally, we showed that concomitant injection of IL2 DNA and pcDNA3-CRT/E7 with electroporation resulted in higher numbers of E7-specific, IFN $\gamma$ <sup>+</sup> CTLs in tumor-bearing mice. Moreover, coadministration of IL2 DNA with CRT/E7 DNA improved the ratio of immunocompetent cells relative to the amount of immunosuppressive cells, in terms of the ratio of CTLs to regulatory T (Treg) cells and the ratio of CTLs to myeloid-derived suppressor cells (MDSCs). Thus our study demonstrates that IL2 can enhance the E7-specific CTL responses and antitumor effects produced by pcDNA3-CRT/E7.

## RESULTS

Coadministration of DNA-encoding IL2 significantly enhances the E7 antigen-specific CD8+ T-cell immune responses in the cervicovaginal tract induced by intravaginal injection of pcDNA3-CRT/E7

To assess the effect of pcDNA3-CRT/E7 vaccination combined with IL2 plasmid, we immunized C57BL/6 mice with various combinations of pcDNA3-CRT/E7 and pcDNA3-IL2 followed by electroporation. Mice were vaccinated submucosally in the cervicovaginal tracts with pcDNA3 empty vector, pcDNA3-IL2, pcDNA3-CRT/E7 or pcDNA3-IL2 together with pcDNA3-CRT/E7, each followed by electroporation. After 7 days, the mice were treated again with the same regimen. Splenocytes and cells from the cervicovaginal tract of treated mice were isolated and the presence of antigen-specific CD8+ T cells were analyzed using flow cytometry. The mice treated with pcDNA3-CRT/E7, both with or without IL2 DNA followed by electroporation, had a notably greater amount of E7-specific CD8+ T cells in comparison to those treated with IL2 DNA only or empty vector (Figures 1a–b). Furthermore, mice that were inoculated with pcDNA3-CRT/E7 combined with IL2 DNA, then electroporated, had more E7-specific CD8+ T cells in the cervicovaginal tract than mice inoculated with pcDNA3-CRT/E7 only (Figures 1c–d).

Coadministration of DNA-encoding IL2 significantly increases the antitumor effects elicited by intravaginal injection of pcDNA3-CRT/E7

Next we studied the antitumor effects elicited by the various DNA vaccines employed in this study, followed by electroporation. C57BL/6 mice were challenged submucosally in the cervicovaginal tract with  $2 \times 10^4$  of TC-1-Luc cells and treated with different DNA plasmids 3 days after tumor challenge and then boosted 7 days later with the same regimen. Mice treated with pcDNA3-CRT/E7 only generated enhanced antitumor effects in comparison to mice treated with IL2 DNA or empty vector alone (Figures 2a–b). Additionally, mice treated with pcDNA3-CRT/E7 and IL2 DNA vaccines had significantly less luciferase activity compared with mice treated with pcDNA3-CRT/E7 only, suggesting more significant tumor control. Additionally, Figure 2c indicates that TC-1 tumor-bearing mice immunized with pcDNA3-CRT/E7 and IL2 DNA had prolonged survival in comparison to those only treated with pcDNA3-CRT/E7.

Coadministration of DNA-encoding IL2 enhances the production of tumor-infiltrating E7-specific interferon- $\gamma$  (IFN $\gamma$ )-secreting CD8+ T cells elicited by intravaginal injection of pcDNA3-CRT/E7

Next we examined the generation of E7-specific IFN $\gamma$ -secreting CD8+ T cells following intravaginal injection of the various DNA vaccines. Tumor-bearing mice were vaccinated with the different DNA constructs and electroporated and then boosted 7 days later with the identical regimen. Cells from the spleen and the cervicovaginal tract were harvested and assessed by intracellular cytokine staining and flow cytometric analysis. Mice treated with pcDNA3-CRT/E7 alone or in combination with IL2 DNA followed by electroporation produced a larger amount of systemic E7-specific IFN $\gamma$ + CD8+ T cells compared with those treated with IL2 DNA only or empty vector (Figures 3a and b). Importantly, Figures 3c and d demonstrate that immunization with CRT/E7 DNA stimulates a significantly greater quantity of local E7-specific IFN $\gamma$ + CD8+ T cells in the cervical vaginal tract than immunization with IL2 DNA or empty vector alone. Furthermore, immunization with pcDNA3-CRT/E7 combined with IL2 DNA generates the most robust local IFN $\gamma$ + CD8+ T-cell response.

Coadministration of DNA-encoding IL2 increases the CD8+ T cell-to-Treg cell ratio among splenocytes generated by intravaginal injection of pcDNA3-CRT/E7 DNA

We then sought to gain a better understanding of the mechanism by which the pcDNA3-CRT/E7 and IL2 DNA vaccine enhances immune responses by examining its effect on the ratio of CD8+ T cells to Treg cells. TC-1 tumor-bearing mice were immunized and then boosted with the various DNA constructs shown in Figure 2. One week after vaccination boost, splenocytes and tumor cells were isolated and the lymphocytes expressing CD4 and CD25 were classified as Treg cells. Among the splenocytes of tumor-bearing mice, those administered both pcDNA3-CRT/E7 and IL2 DNA plasmids had the highest ratio of CD8+ T cells to Treg cells (Figure 4a). Furthermore, among cells located inside cervicovaginal tumor, mice administered pcDNA3-CRT/E7 had a greater ratio of CD8+ T cells to Treg cells in comparison to those treated with IL2 DNA or empty vector alone (Figure 4b). Of note, mice administered both pcDNA3-CRT/E7 and IL2 DNA plasmids had the greatest CD8+ T cell-to-Treg cell ratio. In summary, these results suggest that coadministration of IL2 DNA with pcDNA3-CRT/E7 DNA improves the ratio of immunocompetent cells relative to the number of immunosuppressive cells.


Coadministration of DNA-encoding IL2 increases the ratio of CD8+ T cells to MDSCs among cervicovaginal tumor cells generated by intravaginal injection of pcDNA3-CRT/E7 DNA

We then examined the effect of pcDNA3-CRT/E7 and IL2 DNA vaccination on the ratio of CD8+ T cells to MDSCs. Tumor-bearing mice received the same treatment regimen described above. One week after vaccination boost, splenocytes and tumor cells were isolated from the mice and cells expressing CD11b and Gr-1 were classified as MDSCs. Between the groups of treated mice, we did not observe a significant difference in the ratio of CD8+ T cells to MDSCs in the splenocytes of tumor-bearing mice (Figure 5a). However, we observed a greater ratio of CD8+ T cells to MDSCs in cervicovaginal tumor of mice treated with pcDNA3-CRT/E7 compared with those treated with IL2 DNA or empty vector alone (Figure 5b). Furthermore, of cells located in cervicovaginal tumor, mice immunized with pcDNA3-CRT/E7 and IL2 DNA plasmids had a greater ratio of CD8+ T cells to MDSCs compared with mice treated with pcDNA3-CRT/E7 only. These results provide further support that coadministration of IL2 DNA with pcDNA3-CRT/E7 followed by electroporation generates the highest amount of immunocompetent cells (CD8+ T cells) relative to the number of immunosuppressive cells in (MDSCs) the tumor microenvironment.

## DISCUSSION

We showed that coadministration of DNA-encoding IL2 enhanced the therapeutic efficacy of pcDNA3-CRT/E7 administered intravaginally followed by electroporation. Specifically, tumor-bearing mice immunized with IL2 DNA and pcDNA3-CRT/E7 produced more potent E7-specific CTL responses and antitumor effects in comparison to those treated with pcDNA3-CRT/E7 only. Additionally, IL2 DNA enhanced the amount of IFN $\gamma$ -secreting E7-specific CD8+ T cells in the cervicovaginal tumors of mice. Finally, we demonstrated that DNA-encoding IL2 coadministered with pcDNA3-CRT/E7 improved the proportion of CD8+ T cells to immunosuppressive cells, including Treg cells and MDSCs in the tumor microenvironment. In summary, these results demonstrate that IL2 DNA can greatly enhance the therapeutic effects elicited by local immunization of the pcDNA3-CRT/E7 followed by electroporation.

Electroporation in the vaginal cavity of mice has previously been used to enhance DNA transfection through the vaginal wall.<sup>10,17</sup> However, to date, there are no clinical trials investigating




**Figure 2.** HPV DNA vaccine combined with IL2 followed by electroporation generates antitumor effect and prolonged survival of tumor-bearing mice. Five-to-8-week-old female C57BL/6 mice were injected with  $2 \times 10^4$  of TC-1-Luc cells submucosally. The mice were then vaccinated with 20  $\mu$ g different DNA constructs via intravaginal injection followed by electroporation on day 3. One week later, the mice were boosted with the same regimen. The luciferase activity was measured using a Xenogen IVIS 100 imager (Alameda, CA, USA) on days 0 and 17. Mice survival were recorded. **(a)** Representative bioluminescence image of the TC-1 tumors in the cervical vaginal tract of mice vaccinated with the various regimens. **(b)** Bar graph summary of the luciferase activity of the tumors. **(c)** Survival of the orthotopic TC-1 tumor-bearing mice. \* $P < 0.05$ ; \*\* $P < 0.01$ .

the therapeutic effects of local DNA vaccination followed by intravaginal electroporation. Several preclinical animal studies have documented tissue damage after electroporation; however, this damage has been minimal and animals recovered quickly. In one study, the degree of damage was shown to be dependent upon factors, such as the electric field intensity, and length of pulse. Thus, by using a lower field intensity, tissue damage was minimized.<sup>22,23</sup> An additional study reported a correlation between increased muscle damage and increased pulse duration; however, after 2 weeks, muscle tissue appeared grossly normal, showing that damaged tissue had regenerated.<sup>24</sup>


Before the current treatment strategy can become applicable in the clinic, a suitable electroporation device for intravaginal use that is tolerable for patients will need to be created. Current electroporation devices utilized in the clinic to enhance the immunogenicity of DNA vaccines are applied conjointly with IM vaccination. A recent Phase I clinical trial tested IM administration of pNGVL4a-CRT/E7(detox) DNA vaccine followed by electroporation using the Ichor Medical Systems TriGrid Delivery System in patients with head and neck cancer (NCT01493154). In addition, the TriGrid Delivery System was combined with GX-188E, another therapeutic HPV DNA vaccine, in patients with cervical intraepithelial neoplasia 3 lesions (NCT02411019). Furthermore, a Phase II trial was conducted in patients with cervical intraepithelial neoplasia 2/3 lesions to evaluate the efficacy of IM VGX-3100 DNA vaccination followed by electroporation using CELLECTRA Constant Current Device. Participating patients showed histopathological regression and clearance after vaccination of VGX-3100 followed by electroporation.<sup>25</sup>

Although our study provides considerable incentive for the development of an electroporation device that can be safely administered to the cervicovaginal region in humans, we understand the potential concerns that may be raised regarding the clinical translation of intravaginal electroporation, including invasiveness and pain level.<sup>26</sup> In a previous study, participating patients reported pain at the vaccine injection site after electroporation. However, this pain was believed to be caused by the high voltage (>200 V) of electroporation that was administered at the injection site after vaccination.<sup>26</sup> In this study, we deliver electroporation at the injection site at a much lower voltage (72 V), which we believe would cause patients less pain at injection site during clinical trials. Furthermore, local anesthetic can be employed in clinical studies to minimize any discomfort produced by vaccination and electroporation.<sup>23</sup> Electroporation has great potential to improve the efficacy of many therapeutic vaccines; however, we believe that electroporation technology must be improved before intravaginal electroporation can be administered in clinical trial. If improved, this may be an extremely effective vaccine delivery route that can expand upon the current electroporation technologies available.


One alternative method used to improve the efficacy of therapeutic HPV DNA vaccines that may also address the concerns of using electroporation includes administering the vaccine proximal to tumor-draining lymph nodes (TDLNs). It is believed that TDLNs have a unique role in tumor immunology and malignant disease progression on several levels. It has been reported that the local microenvironment in TDLNs has a major role in local immune response to tumors.<sup>27</sup> One study investigated the effects of adjuvant therapy on the immunological crosstalk



**Figure 3.** IL2 combined with HPV DNA vaccine followed by electroporation generates significantly more tumor-infiltrating IFN- $\gamma$ -secreting CD8<sup>+</sup> T cells. Five-to-eight-week-old female C57BL/6 mice were injected with  $2 \times 10^4$  of TC-1-Luc cells submucosally. The mice were then immunized with the DNA constructs via intravaginal injection followed by electroporation on day 3. One week later, the mice were boosted with the same regimen. Seven days after the last vaccination, splenocytes and cervicovaginal cells were harvested and stimulated with HPV16 E7aa49–57 peptide in the presence of GolgiPlug, stained for surface CD8 and intracellular IFN- $\gamma$  and analyzed by flow cytometric analysis. (a) Representative flow cytometric image for the amount of IFN- $\gamma$ +CD8<sup>+</sup> T cells among splenocytes. (b) Bar graph summary of flow cytometric analysis. (c) Representative flow cytometric image for the amount of IFN- $\gamma$ +CD8<sup>+</sup> T cells among cervicovaginal cells. (d) Summary of flow cytometric analysis. \* $P < 0.05$ ; \*\* $P < 0.01$ ; NS = not significant.



**Figure 4.** Administration of HPV DNA vaccine combined with IL2 increases the ratio of CD8<sup>+</sup> T cells to Treg cells in splenocytes and cervicovaginal tumor cells. The TC-1 tumor-bearing mice were treated as described in Figure 2. Briefly, 7 days after the last vaccination, splenocytes and tumor cells were isolated from the mice. To detect Treg cells, the lymphocytes were stained with anti-mouse CD4 and CD25 antibodies. (a) Bar graph summary of the ratio of CD8<sup>+</sup> T cells/Treg cells among splenocytes. (b) Bar graph summary of the ratio of CD8<sup>+</sup> T cells/Treg cells among tumor-infiltrating lymphocytes. \* $P < 0.05$ ; \*\* $P < 0.01$ .



**Figure 5.** Administration of HPV DNA vaccine combined with IL2 increases the ratio of CD8<sup>+</sup> T cells/CD11b<sup>+</sup>Gr-1<sup>+</sup> cells in splenocytes and cervicovaginal tumor cells. The TC-1 tumor-bearing mice were treated as described in Figure 2. Seven days after the last vaccination, splenocytes and tumor cells were isolated from the mice. To detect MDSCs, the cells were stained with anti-mouse Gr-1, CD11b and CD8 antibodies. **(a)** Bar graph summary of the ratio of CD8<sup>+</sup> T cells/CD11b<sup>+</sup>Gr-1<sup>+</sup> cells. **(b)** Bar graph summary of the ratio of CD8<sup>+</sup> T cells/CD11b<sup>+</sup>Gr-1<sup>+</sup> cells. \**P* < 0.05; \*\**P* < 0.01.

between TDLN and tumor.<sup>28</sup> TDLN-targeted immunotherapy was shown to redirect tumor-associated immune suppression. Proximal vaccination to TDLNs has the potential to enhance local immune response and serve as an effective alternative to intravaginal electroporation.<sup>28</sup> As vaccination proximal to TDLN has not been widely explored, future studies are warranted to determine its therapeutic efficacy against cervicovaginal tumors.

For the clinical translation of DNA-encoding IL2, it is necessary to define the optimal dose and regimen to be used in human clinical trials. Although recombinant IL2 cytokine is currently approved by the Food and Drug Administration for the treatment of metastatic renal-cell carcinoma and malignant melanoma,<sup>18</sup> additional clinical trials are ongoing to improve IL2 treatments, limit their toxicities and examine whether IL2 may be effective in treating other cancers. High-dose IL2 treatments have been associated with high toxicities, including malaise, nausea and flu-like symptoms;<sup>29,30</sup> however, these toxicities were generally reversible after discontinuation of IL2 treatment. Alternatively, in several clinical trials low-dose IL2 was well tolerated by both patients and healthy volunteers, resulting in mild-to-moderate adverse effects.<sup>30</sup>

In order to reduce the toxicity and associated adverse effects of IL2 therapy, a previous study combined IL2 with a peptide vaccine, gp100, to treat patients with metastatic melanoma and found that gp100 improved response rates and progression-free survival.<sup>31</sup> In addition, two previous studies used recombinant adeno-associated viral vector-mediated gene transfer to both encode and release IL2.<sup>29,32</sup> One study found that a wide variety of tissues could be transduced with IL2 with minimal toxicity,<sup>32</sup> while the other study found that long-term low-dose IL2 release did not induce toxicity.<sup>29</sup> IL2 has also been examined in clinical trials in combination with monoclonal antibody for treatment of metastatic kidney cancer,<sup>33,34</sup> natural killer cells and chemotherapy to treat melanoma and kidney cancer<sup>35</sup> and in an aerosolized form to treat pulmonary metastases (NCT01590069). Although IL2 has been shown to be beneficial in the treatment of some cancers, further testing is warranted to broaden its application and improve its tolerability. Coadministration of DNA-encoding IL2 has not yet been clinically tested.

The successful use of IL2 to improve the efficacy of therapeutic HPV DNA vaccines further supports the use of additional cytokines in a similar treatment regimen. Previously, we have also tested the coadministration of DNA vaccine with cytokine granulocyte macrophage colony-stimulating factor and showed that granulocyte macrophage colony-stimulating factor was able to augment antigen-specific antitumor responses elicited by DNA vaccination.<sup>36</sup> It may be worthwhile to screen other cytokines to identify those that can elicit optimal therapeutic effects. Additional candidate molecules include those that can stimulate immune responses in the tumor microenvironment, including IL12 and IL15, which have been shown to be more effective at inhibiting tumor growth than IL2.<sup>18</sup> It will be desirable to test whether the DNA encoding any of these cytokines can be used to further enhance the therapeutic HPV DNA vaccine.

Here we demonstrated that local vaccination with IL2 and pcDNA3-CRT/E7 followed by electroporation elicited potent HPV antigen-specific immune responses, both locally and systemically. These data suggest that the cervicovaginal tract may be an ideal location for vaccine administration to generate a more potent local CTL response. Other investigators have demonstrated that intravaginal immunization with HPV pseudoviruses produced a potent HPV-specific intracervical epithelial CD103<sup>+</sup> CD8<sup>+</sup> T-cell response by stimulating and retaining primed HPV-specific CD8<sup>+</sup> T cells in the cervicovaginal tract.<sup>15</sup> Moreover, it has been demonstrated that local vaccination in the mucosa generated more potent CD8<sup>+</sup> T-cell responses against mucosal TC-1 tumors in mice.<sup>13</sup> The therapeutic effects of the coadministration of our DNA vaccine-encoding IL2 were demonstrated through immune response of CTLs and a strong antitumor response against TC-1 tumors. In this system, these data suggest that CD4<sup>+</sup>CD25<sup>+</sup> Tregs may not have been as crucial to the immunotherapeutic effect of this vaccine as CD8<sup>+</sup> T cells. However, it is believed that the generation and homing of effector memory T cells to the cervicovaginal tract epithelium may be crucial for the initiation of a potent HPV-specific immune response against cervical tumors.<sup>14</sup> Future studies are necessary in order to determine the mechanism of this process.

In summary, the current study presents a treatment regimen for HPV-associated lesions consisting of pcDNA3-CRT/E7 and IL2 DNA

administered intravaginally followed by electroporation. We showed that this treatment regimen could produce an effective therapeutic antitumor immune response and render the tumor microenvironment more immunocompetent. Owing to the importance of local T-cell responses in clearing HPV-associated tumors, the current described approach represents a potential strategy for the control of HPV-associated lesions. Future studies are warranted to develop a clinically applicable electroporation device that is safe and tolerable for vaginal use.

## MATERIALS AND METHODS

### Mice

Five-to-8-week-old female C57BL/6 mice were acquired from the National Cancer Institute and housed in the Johns Hopkins Oncology Center Animal Facility (Baltimore, MD, USA) under specific pathogen-free conditions. Only female mice were chosen for this study because cervical cancer only affects females. Mice were weighted before randomization for studies to further ensure the health of the animals and to control this variable. All animal procedures were performed in accordance with animal protocols approved by the Johns Hopkins Institutional Animal Care and Use Committee and in compliance with recommendations for the proper use and care of laboratory animals. The sample size for each experiment was five mice per group ( $n=5$ ). Based on previous experiments, this sample size was large enough to see a statistical significance between treatment groups. All mouse experiments were replicated in the laboratory two times.

### Cells

The generation of luciferase-expressing TC-1 cells (TC-1-Luc) has been described previously.<sup>37</sup> The cells were cultured in RPMI 1640 supplemented with 50 units  $\text{ml}^{-1}$  of penicillin/streptomycin, 1 mM of sodium pyruvate, 10% of fetal bovine serum, 2 mM of L-glutamine and 2 mM of non-essential amino acids, at 37 °C with 5%  $\text{CO}_2$ . The TC-1 cell line has been tested and authenticated using the method described previously.<sup>38</sup> Additionally, the TC-1 cell line is tested monthly for mycoplasma contamination.

### Peptides, antibodies and HPV16 E7 tetramer

Fluorescein isothiocyanate (FITC)- and phycoerythrin (PE)-conjugated anti-mouse CD8a (clone 53.6.7, catalog nos. 553031 and 553032, respectively) and FITC-conjugated anti-mouse IFN- $\gamma$  (clone XMG1.2, catalog no. 554411) antibodies were acquired from BD Pharmingen (San Diego, CA, USA). Purified anti-mouse CD16/32 (Fc Block, catalog no. 553141) was purchased from BD Pharmingen. FITC-conjugated anti-mouse CD4 (catalog no. 11-0042-82), allophycocyanin (APC)-conjugated CD25 (catalog no. 17-0251-82), FITC-conjugated Gr-1 (catalog no. 11-5931-82) and PE-conjugated CD11b (catalog no. 12-0112-82) antibodies were acquired from eBioscience (San Diego, CA, USA). The H-2D<sup>b</sup>-restricted HPV16 peptide, E7aa49–57 (RAHYNIVTF), was synthesized by Macromolecular Resources (Denver, CO, USA) with  $\geq 80\%$  purity. PE-conjugated HPV16 E7aa49–57 peptide-loaded H-2D<sup>b</sup> tetramer was provided by NIH Tetramer Core Facility (Atlanta, GA, USA).

### DNA vaccines

The generation of the pcDNA3-CRT/E7 DNA vaccine has been previously described.<sup>6</sup> The construction of the pcDNA3-IL2 plasmid was cloned by our laboratory. The pcDNA3-luciferase DNA plasmid was kindly provided by Dr Hyam I. Levitsky (Johns Hopkins Medical Institutions, Baltimore, MD, USA).

### DNA vaccination/electroporation

Mice were anesthetized prior to immunization. In all, 20  $\mu\text{g}$  of DNA constructs were administered into the cervicovaginal tracts by submucosal injection and followed by electroporation with an ECM830 Square Wave Electroporation System (BTX, Harvard Apparatus Inc., Holliston, MA, USA). Electroporation was administered at the site of injection using a tweezer-trode with one plate inside the vagina and the other on the ipsilateral vulva. Eight unipolar, electrical pulses of 72 V were delivered for 20 ms (pulse duration) at 200 ms intervals. The mice received the same treatment regimen 7 days later.

### Splenocyte and cervicovaginal cell preparation

Vaginal tissues were surgically removed via sterile technique, placed in RPMI-1640 medium supplemented with 100 U  $\text{ml}^{-1}$  of penicillin and 100  $\mu\text{g}$   $\text{ml}^{-1}$  of streptomycin and then washed with phosphate-buffered saline (PBS). Next the tissues were minced into 1–2-mm pieces and placed in serum-free RPMI-1640 medium supplemented with 0.05 mg  $\text{ml}^{-1}$  of collagenase I, 0.05 mg  $\text{ml}^{-1}$  of collagenase IV, 0.25 mg  $\text{ml}^{-1}$  of DNase I, 0.025 mg  $\text{ml}^{-1}$  of hyaluronidase IV, 100  $\mu\text{g}$   $\text{ml}^{-1}$  of streptomycin and 100 U  $\text{ml}^{-1}$  of penicillin. The minced tissue were incubated at 37 °C with periodic agitation for 1 h. The undigested tissue fragments in the solution were then filtered through a 70- $\mu\text{m}$  nylon filter mesh. The individual cervicovaginal cells were then washed twice using PBS, and viable cells were determined by trypan blue dye exclusion. To prepare the splenocytes, the spleen was minced using a 70- $\mu\text{m}$  nylon filter mesh. The red blood cells were lysed using ACK lysis buffer, and the cells were washed. Viable cells were determined by trypan blue dye exclusion.

### Flow cytometric analysis

All *in vitro* experiments were replicated three times in the laboratory. To analyze the HPV16 E7-specific CTL responses using tetramer, splenocytes and cervicovaginal cells were washed once with fluorescence-activated cell sorter (FACS) wash buffer (PBS+0.5 bovine serum albumin). All cells were preincubated with BD Fc Block to avoid nonspecific antibody binding through the surface Fc receptor. The cells were then stained with FITC-conjugated anti-mouse CD8a (catalog no. 553031) from BD Pharmingen and PE-conjugated HPV16 E7aa49–57 peptide loaded H-2D<sup>b</sup> E-7 tetramer. Following the wash, cells were acquired with a FACSCalibur flow cytometer (BD Biosciences, Mountain View, CA, USA) and analyzed using the FlowJo software (TreeStar Inc., Ashland, OR, USA).

To analyze the HPV16 E7-specific CTL responses using intracellular IFN- $\gamma$  staining, cervicovaginal cells and splenocytes were stimulated overnight with 1  $\mu\text{g}$   $\text{ml}^{-1}$  HPV16 E7aa49–57 peptide in the presence of 1  $\mu\text{l}$   $\text{ml}^{-1}$  of GolgiPlug (BD Pharmingen). After stimulation, the cells were washed one time with FACS wash buffer and then stained with PE-conjugated anti-mouse CD8a antibody (catalog no. 553032) acquired from BD Pharmingen. After staining, the cells were permeabilized and then fixed using the Cytofix/Cytoperm Kit following the instructions provided by the manufacturer (BD Pharmingen). Intracellular IFN- $\gamma$  staining was performed using FITC-conjugated anti-mouse IFN- $\gamma$  antibody (catalog no. 554411) purchased from BD Pharmingen. After the wash, cells were acquired with a FACSCalibur flow cytometer.

For the detection of Tregs, single-cell preparations from the mouse spleen and tumor were resuspended in FACS staining buffer and incubated with anti-CD4-FITC (catalog no. 11-0042-82, eBioscience) and CD25-APC (catalog no. 17-0251-82, eBioscience) antibodies for 30 min at 4 °C. To detect CD11b<sup>+</sup>Gr-1<sup>+</sup> MDSCs from splenocytes and single tumor cell preparations obtained from mice, cells were stained with FITC-conjugated anti-mouse Gr-1 (catalog no. 11-5931-82, eBioscience) and PE-conjugated anti-mouse CD11b (catalog no. 12-0112-82, eBioscience) antibodies. Analyses were carried out on a Becton-Dickinson FACScan with CELLQuest software (Franklin Lakes, NJ, USA).

### In vivo tumor treatment experiments

Mice were challenged with  $2 \times 10^4$  of TC-1-Luc cells injected submucosally into the wall of the vagina. Three days following the tumor challenge, mice were immunized with 20  $\mu\text{g}$  of DNA constructs as indicated. The mice received the same regimen again 7 days later. The growth of cervicovaginal tumor was monitored using bioluminescence via Xenogen IVIS 100 imager at the indicated times. The experiments were performed in a non-blind manner.

### Statistical analysis

All data are expressed as means  $\pm$  s.d., including error bars based on s.d. to account for any statistical uncertainty. Two-tailed Student's *t*-test was used to compare the differences between individual data points. Non-parametric Mann-Whitney test was used to compare two different groups. The tumor treatment experiment results were evaluated using Kaplan–Meier analysis. *P*-values < 0.05 were considered significant.

### CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGEMENTS

This work was funded by the United States National Institutes of Health (NIH) Cervical Cancer Specialized Program of Research Excellence (SPORE) (P50 CA098252), 5R21 (AI109259), 1R21 grant (CA194896) and R01 grant (CA114425).

## REFERENCES

- Yang A, Jeang J, Cheng K, Cheng T, Yang B, Wu TC *et al*. Current state in the development of candidate therapeutic HPV vaccines. *Expert Rev Vaccines* 2016; **15**: 989–1007.
- zur Hausen H. Papillomaviruses and cancer: from basic studies to clinical application. *Nat Rev Cancer* 2002; **2**: 342–350.
- Doorbar J. Model systems of human papillomavirus-associated disease. *J Pathol* 2016; **238**: 166–179.
- Spee P, Neefjes J. TAP-translocated peptides specifically bind proteins in the endoplasmic reticulum, including gp96, protein disulfide isomerase and calreticulin. *Eur J Immunol* 1997; **27**: 2441–2449.
- Sadasivan B, Lehner PJ, Ortmann B, Spies T, Cresswell P. Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* 1996; **5**: 103–114.
- Cheng WF, Hung CF, Chai CY, Hsu KF, He L, Ling M *et al*. Tumor-specific immunity and antiangiogenesis generated by a DNA vaccine encoding calreticulin linked to a tumor antigen. *J Clin Invest* 2001; **108**: 669–678.
- Center SKCCA pilot study of p nGVL4a-CRT/E7 (Detox) for the treatment of patients with HPV16+ cervical intraepithelial neoplasia 2/3 (CIN2/3) [Internet]. *ClinicalTrials.gov* [Internet]. National Library of Medicine (US): Bethesda (MD), USA, 2009.
- Center SKCC. Safety study of HPV DNA vaccine to treat head and neck cancer patients [Internet]. *ClinicalTrials.gov* [Internet]. National Library of Medicine (US): Bethesda, MD, USA, 2011.
- Kichaev G, Mendoza JM, Amante D, Smith TR, McCoy JR, Sardesai NY *et al*. Electroporation mediated DNA vaccination directly to a mucosal surface results in improved immune responses. *Hum Vaccin Immunother* 2013; **9**: 2041–2048.
- Kanazawa T, Takashima Y, Hirayama S, Okada H. Effects of menstrual cycle on gene transfection through mouse vagina for DNA vaccine. *Int J Pharm* 2008; **360**: 164–170.
- Best SR, Peng S, Juang CM, Hung CF, Hannaman D, Saunders JR *et al*. Administration of HPV DNA vaccine via electroporation elicits the strongest CD8+ T cell immune responses compared to intramuscular injection and intradermal gene gun delivery. *Vaccine* 2009; **27**: 5450–5459.
- Bagarazzi ML, Yan J, Morrow MP, Shen X, Parker RL, Lee JC *et al*. Immunotherapy against HPV16/18 generates potent TH1 and cytotoxic cellular immune responses. *Sci Transl Med* 2012; **4**: 155ra138.
- Sandoval F, Terme M, Nizard M, Badoual C, Bureau MF, Freyburger L *et al*. Mucosal imprinting of vaccine-induced CD8(+) T cells is crucial to inhibit the growth of mucosal tumors. *Sci Transl Med* 2013; **5**: 172ra20.
- Trimble CL, Clark RA, Thoburn C, Hanson NC, Tassello J, Frosina D *et al*. Human papillomavirus 16-associated cervical intraepithelial neoplasia in humans excludes CD8 T cells from dysplastic epithelium. *J Immunol* 2010; **185**: 7107–7114.
- Cuburu N, Graham BS, Buck CB, Kines RC, Pang YY, Day PM *et al*. Intravaginal immunization with HPV vectors induces tissue-resident CD8+ T cell responses. *J Clin Invest* 2012; **122**: 4606–4620.
- Sun YY, Peng S, Han L, Qiu J, Song L, Tsai Y *et al*. Local HPV recombinant vaccinia boost following priming with an HPV DNA vaccine enhances local HPV-specific CD8+ T-cell-mediated tumor control in the genital tract. *Clin Cancer Res* 2016; **22**: 657–669.
- Sun YY, Peng S, Qiu J, Miao J, Yang B, Jeang J *et al*. Intravaginal HPV DNA vaccination with electroporation induces local CD8+ T-cell immune responses and antitumor effects against cervicovaginal tumors. *Gene Therapy* 2015; **22**: 528–535.
- Waldmann TA. The biology of interleukin-2 and interleukin-15: implications for cancer therapy and vaccine design. *Nat Rev Immunol* 2006; **6**: 595–601.
- Kang TH, Mao CP, He L, Tsai YC, Liu K, La V *et al*. Tumor-targeted delivery of IL-2 by NKG2D leads to accumulation of antigen-specific CD8+ T cells in the tumor loci and enhanced anti-tumor effects. *PLoS One* 2012; **7**: e35141.
- Kim JJ, Yang JS, Manson KH, Weiner DB. Modulation of antigen-specific cellular immune responses to DNA vaccination in rhesus macaques through the use of IL-2, IFN-gamma, or IL-4 gene adjuvants. *Vaccine* 2001; **19**: 2496–2505.
- Kim JJ, Yang JS, Montaner L, Lee DJ, Chalian AA, Weiner DB. Coimmunization with IFN-gamma or IL-2, but not IL-13 or IL-4 cDNA can enhance Th1-type DNA vaccine-induced immune responses in vivo. *J Interferon Cytokine Res* 2000; **20**: 311–319.
- Khan AS, Pope MA, Draghia-Akli R. Highly efficient constant-current electroporation increases in vivo plasmid expression. *DNA Cell Biol* 2005; **24**: 810–818.
- Draghia-Akli R, Khan AS, Brown PA, Pope MA, Wu L, Hiraio L *et al*. Parameters for DNA vaccination using adaptive constant-current electroporation in mouse and pig models. *Vaccine* 2008; **26**: 5230–5237.
- Mathiesen I. Electroporation of skeletal muscle enhances gene transfer in vivo. *Gene Therapy* 1999; **6**: 508–514.
- Trimble CL, Morrow MP, Kravnyak KA, Shen X, Dallas M, Yan J *et al*. Safety, efficacy, and immunogenicity of VGX-3100, a therapeutic synthetic DNA vaccine targeting human papillomavirus 16 and 18 E6 and E7 proteins for cervical intraepithelial neoplasia 2/3: a randomised, double-blind, placebo-controlled phase 2b trial. *Lancet* 2015; **386**: 2078–2088.
- Sallberg M, Frelin L, Ahlen G, Sallberg-Chen M. Electroporation for therapeutic DNA vaccination in patients. *Med Microbiol Immunol* 2015; **204**: 131–135.
- Munn DH, Mellor AL. The tumor-draining lymph node as an immune-privileged site. *Immunol Rev* 2006; **213**: 146–158.
- Thomas SN, Vokali E, Lund AW, Hubbell JA, Swartz MA. Targeting the tumor-draining lymph node with adjuvanted nanoparticles reshapes the anti-tumor immune response. *Biomaterials* 2014; **35**: 814–824.
- Churlaud G, Jimenez V, Ruberte J, Amadouji Zin M, Fourcade G, Gottrand G *et al*. Sustained stimulation and expansion of Tregs by IL2 control autoimmunity without impairing immune responses to infection, vaccination and cancer. *Clin Immunol* 2014; **151**: 114–126.
- Klatzmann D, Abbas AK. The promise of low-dose interleukin-2 therapy for autoimmune and inflammatory diseases. *Nat Rev Immunol* 2015; **15**: 283–294.
- Schwartzentruber DJ, Lawson DH, Richards JM, Conry RM, Miller DM, Treisman J *et al*. gp100 peptide vaccine and interleukin-2 in patients with advanced melanoma. *N Engl J Med* 2011; **364**: 2119–2127.
- Goudy KS, Johnson MC, Garland A, Li C, Samulski RJ, Wang B *et al*. Inducible adeno-associated virus-mediated IL-2 gene therapy prevents autoimmune diabetes. *J Immunol* 2011; **186**: 3779–3786.
- Dandamudi UB, Ghebremichael M, Sosman JA, Clark JJ, McDermott DF, Atkins MB *et al*. A phase II study of bevacizumab and high-dose interleukin-2 in patients with metastatic renal cell carcinoma: a Cytokine Working Group (CWG) study. *J Immunother* 2013; **36**: 490–495.
- Donini M, Buti S, Lazzarelli S, Bozzetti R, Rivoltini L, Camisaschi C *et al*. Dose-finding/phase II trial: bevacizumab, immunotherapy, and chemotherapy (BIC) in metastatic renal cell cancer (mRCC). Antitumor effects and variations of circulating T regulatory cells (Treg). *Target Oncol* 2015; **10**: 277–286.
- Berend MJ, North RJ. T-cell-mediated suppression of anti-tumor immunity. An explanation for progressive growth of an immunogenic tumor. *J Exp Med* 1980; **151**: 69–80.
- Lee SJ, Song L, Yang MC, Mao CP, Yang B, Yang A *et al*. Local administration of granulocyte macrophage colony-stimulating factor induces local accumulation of dendritic cells and antigen-specific CD8+ T cells and enhances dendritic cell cross-presentation. *Vaccine* 2015; **33**: 1549–1555.
- Lin KY, Guarnieri FG, Staveley-O'Carroll KF, Levitsky HI, August JT, Pardoll DM *et al*. Treatment of established tumors with a novel vaccine that enhances major histocompatibility class II presentation of tumor antigen. *Cancer Res* 1996; **56**: 21–26.
- Kang TH, Noh KH, Kim JH, Bae HC, Lin KY, Monie A *et al*. Ectopic expression of X-linked lymphocyte-regulated protein pM1 renders tumor cells resistant to antitumor immunity. *Cancer Res* 2010; **70**: 3062–3070.