

ARTICLE



The genetic landscape of *SMARCB1* alterations in *SMARCB1*-deficient spectrum of mesenchymal neoplasms

Josephine K. Dermawan¹, Samuel Singer², William D. Tap³, Benjamin A. Nacev³, Ping Chi³, Leonard H. Wexler⁴, Michael V. Ortiz⁴, Mrinal Gounder³ and Cristina R. Antonescu¹✉

© The Author(s), under exclusive licence to United States & Canadian Academy of Pathology 2022

SMARCB1 biallelic inactivation resulting in *SMARCB1*/INI1 deficiency drives a wide range of malignancies, including many mesenchymal tumors. However, the specific types of *SMARCB1* alterations and spectrum of cooperating mutations among various types of sarcomas has not been well investigated. We profiled *SMARCB1* genetic alterations by targeted DNA sequencing and fluorescence in situ hybridization (FISH) in a large cohort of 118 soft tissue and bone tumors, including *SMARCB1*-deficient sarcomas (78, 66%): epithelioid sarcomas, epithelioid peripheral nerve sheath tumors, poorly differentiated chordomas, malignant rhabdoid tumors, and soft tissue myoepithelial tumors, as well as non-*SMARCB1*-deficient sarcomas (40, 34%) with various *SMARCB1* genetic alterations (mutations, copy number alterations). *SMARCB1* loss by immunohistochemistry was present in 94% *SMARCB1* pathogenic cases. By combined sequencing and FISH assays, 80% of *SMARCB1*-deficient tumors harbored homozygous (biallelic) *SMARCB1* loss, while 14% demonstrated heterozygous *SMARCB1* loss-of-function (LOF) alterations, and 6% showed no demonstrable *SMARCB1* alterations. FISH and sequencing were concordant in the ability to detect *SMARCB1* loss in 48% of cases. Epithelioid sarcomas most commonly (75%) harbored homozygous deletions, while a subset showed focal intragenic deletions or LOF mutations (nonsense, frameshift). In contrast, most soft tissue myoepithelial tumors (83%) harbored *SMARCB1* nonsense point mutations without copy number losses. Additionally, clinically significant, recurrent co-occurring genetic events were rare regardless of histotype. By sequencing, extended 22q copy number loss in genes flanking the *SMARCB1* locus (22q11.23) occurred in one-third of epithelioid sarcomas and the majority of poorly differentiated chordomas. Poorly differentiated chordomas and soft tissue myoepithelial tumors showed significantly worse overall and disease-free survival compared to epithelioid sarcomas. Overall, *SMARCB1* LOF alterations predominate and account for *SMARCB1* protein loss in most cases: majority being biallelic but a subset were heterozygous. In contrast, *SMARCB1* alterations of uncertain significance can be seen in diverse sarcomas types and does not indicate a *SMARCB1*-deficient entity.

Modern Pathology (2022) 35:1900–1909; <https://doi.org/10.1038/s41379-022-01148-x>

INTRODUCTION

SMARCB1 (located on chromosome 22q11.23) encodes for INI1/*SMARCB1*/BAF47/sSNF5 protein which is a core subunit of the ubiquitously expressed SWI/SNF (BAF) chromatin remodeling complex. Loss of *SMARCB1* function destabilizes the BAF complex, thereby perturbing its chromatin occupancy and opposing function of polycomb-mediated promoter repression¹. *SMARCB1* deficiency is seen in up to 20% of human malignancies^{2, 3}, including many soft tissue and bone tumors often displaying epithelioid or rhabdoid morphology^{4–7}. Loss of *SMARCB1* expression is seen in the majority of epithelioid sarcomas (distal and proximal types)^{8, 9}, 50–70% of epithelioid malignant peripheral nerve sheath tumors (eMPNST)^{9, 10}, 50% of epithelioid schwannomas (eSCHW)¹¹, 30% of soft tissue myoepithelial tumors (STME)^{9, 12}, majority of poorly differentiated chordomas (PD-CHO)^{13, 14}, and almost all malignant rhabdoid tumors (MRT)^{15, 16}. In these studies, *SMARCB1* loss was largely evidenced by loss of immunohistochemical expression of *SMARCB1*, or in a few studies, detection of homozygous or heterozygous *SMARCB1* deletions

using fluorescence in situ hybridization (FISH) or microarray-based comparative genomic hybridization (aCGH). However, the specific types and distribution of *SMARCB1* genetic alterations among the various *SMARCB1*-deficient sarcomas have not been studied in detail.

To better characterize the nature and spectrum of *SMARCB1* alterations in mesenchymal tumors, we profiled a large cohort of 118 soft tissue and bone sarcomas by targeted DNA sequencing and FISH, focusing on *SMARCB1*-deficient sarcomas, correlating *SMARCB1* copy number (homozygous/heterozygous deletion, amplification) and mutational status with histotypes, *SMARCB1* protein immunohistochemical expression, extent of chromosome 22q copy number losses, and co-occurring genomic alterations.

MATERIALS AND METHODS

Case selection and study cohort

After approval from the Institutional Review Board, cases were selected from the Pathology Department archives of Memorial Sloan Kettering

¹Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA. ²Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY, USA. ³Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA. ⁴Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA. ✉email: antonesc@mskcc.org

Received: 21 June 2022 Revised: 1 August 2022 Accepted: 4 August 2022

Published online: 10 September 2022

Cancer Center (MSKCC) from 2013 to 2021. Inclusion criteria were: 1) soft tissue and bone tumors for which SMARCB1 deficiency has been well established in the literature to be pathogenic, i.e., proximal type epithelioid sarcoma (EPS-P), distal type epithelioid sarcoma (EPS-D), eMPNST, eSCHW, PD-CHO, MRT, and STME; 2) other soft tissue and bone tumors displaying SMARCB1 genetic alterations: deletion, amplification, point mutations, etc.; 3) all cases included have already had targeted DNA sequencing performed on formalin-fixed paraffin embedded (FFPE) tissue [105 cases by clinical MSK-IMPACT (103 with matched germline control, 2 with pooled FFPE control), 13 cases by research DNA sequencing using similar platform]. A total of 118 cases were included. Note that all STME in this study were malignant tumors, and thus may also be referred to as "myoepithelial carcinomas". Immunohistochemistry (IHC) using antibody clone 25/BAF47 (BD Bioscience, catalog# 612110, 1:200 dilution) for SMARCB1 was performed in all cases where SMARCB1 deficiency is pathogenic (criteria #1) and a subset of the other cases (criteria #2) when archival tissue could be obtained. SMARCB1 IHC was available in 95 (80%) cases.

Fluorescence in situ hybridization (FISH)

SMARCB1 FISH testing was performed for 57 (48%) of cases. Details of probe hybridization conditions and signal detection were described previously¹². Custom BAC probes for SMARCB1 were designed and used to detect the presence of large homozygous or heterozygous deletions of SMARCB1 at chromosome 22q11.23. Telomeric probe against EWSR1 at chromosome 22q12 served as a reference. Normal copy number pattern was defined when two copies of the SMARCB1 gene were detected, with a 1:1 ratio to the control probe (i.e., telomeric-EWSR1 or 22q11). Heterozygous deletion was defined as only one copy of the gene of interest being present compared to the reference control probe on 22q (ratio 1:2). Homozygous deletion of SMARCB1 was interpreted when both copies of the gene were lost, compared to the control probes, either telomeric-EWSR1 or 22q11. A monosomy pattern (or large deletion) was defined if one allele copy of both the gene of interest and control were lost, with a ratio of 1:1.

Targeted DNA sequencing for mutational and copy number profiling

Detailed descriptions of MSK-IMPACT workflow and data analysis, a hybridization capture-based targeted DNA next-generation sequencing (NGS) assay for solid tumor were described previously¹⁷. All mutational and copy number calls were generated by the standard MSK-IMPACT pipeline¹⁷. Pathogenic (mostly loss-of-function) genetic alterations included copy number deletion, intragenic deletion, nonsense mutation, frameshift insertion/deletion, most splice site mutations, and missense mutations annotated as oncogenic or likely oncogenic by OncoKB¹⁸. Non-pathogenic or passenger mutations included in-frame insertion/deletion and missense mutations of unknown significance. Copy number amplification and deletion are defined as gains and losses of gene-level copy number greater than two-fold in the tumor relative to pooled FFPE normal based on NGS. Intragenic deletion denotes deletions that span one or more exons of a gene, with distinct and abrupt loss of pileup reads visualizable on integrated genome browser (IGV) but may or may not result in less than 2-fold change of gene-level copy number.

Data analysis was performed using R version 4.1.0. MSK-IMPACT data were imported using R packages "gnomeR" version 1.0.0 and "cbioportalR" version 0.0.0.9000. Mutations and gene-level copy number alterations were visualized and summarized using the R package "ComplexHeatmap" version 2.8.0¹⁹. Cohort-level copy number segmentation data were summarized and visualized using the R package "GenVisR" version 1.24.0²⁰.

Survival analysis

Survival analysis by comparison of hazard ratios using log rank *P* testing and visualization of Kaplan–Meier curves were performed using R packages "survminer" version 0.4.9 and "survival" version 3.2.13. Clinical charts were manually reviewed to document date of initial presentation, disease progression, and survival status. Median time (in years) to disease progression was defined as the time interval between initial presentation (presence of tumor seen radiographically or on physical examination) and the first instance of tumor recurrence or distant metastases after initial surgical resection and/or chemoradiation therapy with radiographically negative evidence of residual tumor.

RESULTS

Spectrum of SMARCB1 genomic alterations and correlations with various histotypes

We classified our cohort of 118 cases into: 1) SMARCB1 pathogenic (78, 66%): soft tissue and bone entities for which SMARCB1 deficiency has been well established to be pathogenic per published studies and also had DNA sequencing data available: EPS-P (30), EPS-D (14), eMPNST (11), eSCHW (7), PD-CHO (7), MRT (3), and STME (6); 2) other soft tissue and bone tumors where SMARCB1 genetic alterations were identified by targeted sequencing but likely represent non-pathogenic or passenger mutations, including SNV or intragenic deletions (27, 23%) and amplifications (13, 11%). Figure 1A summarizes the spectrum of SMARCB1 genetic alterations and corresponding histotypes.

Among the 78 SMARCB1-deficient pathogenic cases, all but four (74, 95%) showed loss of SMARCB1 nuclear expression at the protein level by immunohistochemistry. The 4 cases with retained SMARCB1 were all eSCHW, with otherwise typical histologic features. Fifty-nine (79.7%) of 74 SMARCB1-deficient pathogenic cases with SMARCB1 loss harbored homozygous (biallelic) SMARCB1 loss: mostly large homozygous deletions detected by FISH (Fig. 1B), or homozygous deletions detected by copy number profiling, and a small subset by the presence of two LOF point mutations (nonsense, frameshift, etc.) present in trans (different alleles) (2 cases) or one LOF mutation plus heterozygous deletion by FISH (2 cases).

Additionally, we found that not all SMARCB1-deficient pathogenic cases demonstrated biallelic SMARCB1 inactivation, despite loss of SMARCB1 expression at the protein level (Fig. 1A). Ten (13.5%) of 74 SMARCB1-deficient pathogenic cases with SMARCB1 protein loss (4 EPS-P, 3 STME, 1 PD-CHO, 1 MRT, 1 eSCHW) demonstrated only heterozygous loss-of-function point mutations. Five (6.8%) of 74 SMARCB1-deficient pathogenic cases with SMARCB1 protein loss (3 EPS-D, 1 EPS-P, 1 eMPNST) showed no demonstrable SMARCB1 genetic alterations in SMARCB1 or genes encoding for other SWI/SNF subunits. In total, 15 (20%) out of 74 cases with SMARCB1 loss showed no demonstrable biallelic loss based on molecular testing.

The compared sensitivity between the two assays applied (DNA sequencing vs FISH) showed that among the 53 cases of SMARCB1-deficient pathogenic cases for which FISH was performed and SMARCB1 loss was proven by IHC, FISH and sequencing were concordant in 28 (48%) cases. SMARCB1 abnormalities were not detected by FISH but present via sequencing in 14 (27%) cases: as copy number deletion in 4 cases, as intragenic deletion in 3 cases, and as loss-of-function mutations in 7 cases. On the other hand, SMARCB1 alterations was missed by sequencing but detected by FISH in 10 (19%) cases. In one case of eMPNST, neither sequencing nor FISH detected SMARCB1 alterations, despite loss of SMARCB1 by IHC. Combined, when SMARCB1 was lost by IHC, the sensitivity of sequencing and FISH in detecting SMARCB1 abnormalities was 93% (69/74).

The breakdown of SMARCB1 alterations by tumor types in the SMARCB1-deficient pathogenic category was as follows (Table 1): epithelioid sarcomas, both EPS-P and EPS-D, most frequently harbored homozygous deletions detected by FISH and/or copy number profiling (33, 75%). Within EPS-P, 7 cases (21%) harbored heterozygous intragenic deletions or loss-of-function mutations (nonsense, frameshift). For PD-CHO, 4 (57%) of 7 cases showed homozygous deletions detected by FISH and/or copy number profiling, 2 (28%) cases showed intragenic deletion, and 1 case showed only SMARCB1 frameshift deletion. For eMPNST, 7 (63%) of 11 cases showed homozygous deletions detected by FISH and/or copy number profiling, 2 (18%) cases showed biallelic LOF mutations, and 1 case showed intragenic deletion. For eSCHW, 1 (14%) case showed homozygous deletion and 3 (43%) cases showed heterozygous deletion by copy number profiling, 1 showed a monoallelic nonsense mutation. For MRT, 2 (66%) of




Fig. 1 Spectrum of SMARCB1 genomic alterations and histotypes. **A** OncoPrint showing various soft tissue and bone tumors with SMARCB1 copy number alterations detected by MSK-IMPACT or FISH and SMARCB1 point mutations detected by MSK-IMPACT, their corresponding histotypes, SMARCB1 (INI1) IHC status, with patient age and sex, categorized into: SMARCB1-deficient/pathogenic sarcomas (red), SMARCB1-altered but likely non-pathogenic sarcomas, including SMARCB1-mutated or -deleted (blue) and SMARCB1-amplified sarcomas (green). **B** FISH showing homozygous deletion of SMARCB1 (22q11, red probe) in a case of epithelioid sarcoma (left panel) and a case of epithelioid malignant peripheral nerve sheath tumor (MPNST) (right panel). EWSR1 (22q12, green probe) serves as the reference probe. Unaffected (non-neoplastic) nuclei show 2 red signals plus 2 green signals. In affected (SMARCB1-deficient) nuclei, depending on the extent of 22q deletion, there could be 0 red signals plus 2 green signals, or 0 red signals plus 1 green signal.

3 cases showed homozygous deletion by copy number profiling, 1 showed a monoallelic nonsense mutation. On the contrary, the majority of STME (5, 83%) harbored nonsense mutations in SMARCB1 without copy number losses (monoallelic alteration), while 1 case showed copy number deletion.

Additionally, we identified a cohort of 40 heterogeneous sarcoma histotypes that harbored SMARCB1 genetic alterations, including copy number changes and heterozygous non-LOF mutations. Since SMARCB1 deficiency has not been recognized as the oncogenic driver in these histotypes, this group was considered separately from the SMARCB1 pathogenic category. This was supported by the fact that among the 15 cases tested by SMARCB1 IHC, only 1 case (synovial sarcoma) showed loss of SMARCB1. As a subset of these cases showed SMARCB1 amplification, which has not yet been studied before, we further subdivided this cohort into SMARCB1-altered but nonamplified and SMARCB1-amplified cases. The first group included 27 cases (23% of total cohort) that harbored SMARCB1 copy number losses as well as SMARCB1 mutations (missense and in-frame mutation) of uncertain significance (non-LOF). These tumor types included angiosarcomas, conventional MPNST,

osteosarcomas, spindle cell rhabdomyosarcomas, myxofibrosarcomas, and sarcoma not otherwise specified. The second group included 13 cases (11% of total cohort) with SMARCB1 amplifications, and the histotypes were angiosarcoma, soft tissue leiomyosarcoma, sarcoma not otherwise specified, etc. Eight (61%) of these cases also harbored copy number amplifications at CRKL and MAPK1 (genes flanking SMARCB1 on chromosome 22q).

Co-occurring genomic alterations in SMARCB1-deficient sarcomas

Next, we investigated the types and distribution of co-occurring (secondary) genetic events in SMARCB1-deficient sarcomas (cases with SMARCB1 loss at the protein level). We focused only on recurrent events (occurring more than once) that were likely oncogenic (per OncoKb annotations)¹⁸. Overall, oncogenic, recurrent co-occurring genetic events were rare in SMARCB1-deficient sarcomas: 11 (14%) out of 74 cases of SMARCB1-deficient pathogenic cases—with mostly 1–2 alterations per case. These co-occurring alterations occurred in: TP53, RB1, CDKN2A/2B, ARID1A/B, and BRCA2. Overall, there was no obvious segregation of specific

Table 1. Incidence and types of SMARCB1 alterations across different SMARCB1-deficient histotypes in published studies compared to current study.

Tumor type	Studies	SMARCB1 homozygous deletions	SMARCB1 heterozygous deletions	SMARCB1 intragenic deletions and/or mutations	Total number of cases
Epithelioid sarcoma	Published studies	Sullivan ⁸ 8 (67%)	2 (16%)	2 (16%) – homozygous intragenic deletions	12
	Le Loarer ¹² 2014	36 (90%)			40
	Folpe ³⁸	3 (60%)	2 (40%)		5
	Jamshidi ²¹	4 (40%)	4 (40%)		10
Poorly differentiated chordoma	Current study	33 (75%)		7 (16%)	44
	Published studies	Mobley ¹³ Owosho ¹⁴ 8 (89)	3 (75%)		4
Malignant rhabdoid tumor	Current study	4 (57%)		3 (43%)	7
	Published studies	Jackson ²³ Eaton ²⁴ Lee ¹⁵ 24 (47%) 3 (9%) – germline 26 (40%)-somatic 25 (78%)	11 (22%) 9 (26%)-germline 39 (60%)-somatic	28 (55%) 25 (71%)-germline 39 (60%)-somatic	51 35-germline 65-somatic 32
	Current study	2 (67%)		1 (33%)	3
	Published studies	Schaefer ²⁵ 5 (31%)		7 (44%)	16
Epithelioid schwannoma	Current study	7 (64%)		3 (27%)	11
	Published studies	Schaefer ²⁵ 2 (40%)		3 (60%)	5
Soft tissue myoepithelial tumor	Current study	1 (14%)	3 (43%)	1 (14%)	7
	Published studies	Le Loarer ¹² Folpe ³⁸ 3 (60%) 2 (50%)	1 (25%)	1 (25%) – homozygous intragenic deletions	5 3
	Current study	1 (17%)		5 (83%)	6



Fig. 2 Co-occurring genomic alterations in SMARCB1-deficient sarcomas. A OncoPrint showing recurrent, clinically significant (by OncoKB annotation) co-occurring genetic alterations in SMARCB1-deficient sarcomas and their corresponding histotypes. TMB: tumor mutation burden. **B** OncoPrint showing acquisition of co-occurring mutations or copy number alterations in subsequent recurrence or metastases among proximal-type epithelioid sarcoma and their *SMARCB1* genetic alterations (copy number, loss-of-function mutations). This plot only included cases with *SMARCB1* deletion detected by sequencing but not those detected by FISH.

co-occurring genetic alterations by histotypes among SMARCB1-deficient sarcomas (Fig. 2A).

For five cases of EPS-P, longitudinal sequencing was available from multiple tumors from the same patient over time; we tracked whether there were additional genomic alterations acquired in subsequent recurrence or metastases. Figure 2B illustrates the acquisition of co-occurring mutations in *RB1* and *ID3* in two cases, respectively, and copy number amplifications in *AKT2*, *KIT*/*PDGFRA*, and losses in *RB1* and *PARK2* in four cases, respectively, in the subsequent recurrent/metastatic specimen that were not

identified in the primary tumor. Mutations in *ARID1B*, *TP53* and *BRCA2* were present in both the initial and subsequent samples. We did consider a potential confounder that may account for falsely negative findings: differences in % tumor purity across sequentially sequenced samples from the same patient. Fortunately, tumor purity was comparable if not higher in the primary sample than the subsequent recurrent or metastatic samples in this cohort, mostly because the primary samples were usually resection specimens while the subsequent samples were often small biopsies.




Fig. 3 Extended 22q loss in *SMARCB1*-deficient sarcomas. **A** Cohort-level copy number segmentation plot showing proportion of copy number alterations on chromosome 22q, highlighting the genes flanking *SMARCB1* (22q11.23), from centromeric to telomeric: *CRKL* (22q11.21), *MAPK1* (22q11.21), *CHEK2* (22q12.1), *NF2* (22q12.2), *EP300* (22q13.2), among *SMARCB1*-deficient sarcomas. **B** Oncoprint showing *SMARCB1*-deficient sarcomas and corresponding tumor types with or without copy number alterations at the flanking genes, *CRKL*, *MAPK1*, *CHEK2*, *NF2*, and *EP300*.

Extended 22q loss in *SMARCB1*-deficient sarcomas

Since *SMARCB1* is located on chromosome 22q, we examined the extent of 22q loss in *SMARCB1*-deficient sarcomas. Figure 3A is a copy number segmentation plot of the *SMARCB1*-deficient pathogenic cohort showing varying proportion of copy number losses at the genes flanking *SMARCB1* (22q11.23), from centromeric to telomeric side: *CRKL* (22q11.21), *MAPK1* (22q11.21), *CHEK2* (22q12.1), *NF2* (22q12.2), *EP300* (22q13.2), with *SMARCB1* showing the “deepest” (highest proportion with) copy number loss. Figure 3B illustrates the cases of *SMARCB1*-deficient sarcomas with extended 22q copy number loss beyond *SMARCB1* gene. Overall, among *SMARCB1*-deficient sarcomas where *SMARCB1* deletions were detected by sequencing and/or FISH, extended 22q copy number loss was seen in 9 of 25 (36%) EPS-P, 4 of 11 (36%) EPS-D, 5 of 6 (83%) PD-CHO, 1 of 2 (50%) MRT, 1 of 2 (50%) STME, and 1 of 7 (14%) eMPNST.

Outcome and survival

Among the various histotypes of *SMARCB1*-deficient sarcomas, PD-CHO and STME showed significantly worse overall (OS) and disease-free survival (DFS) compared to EPS-P and EPS-D: median OS for PD-CHO and STME were 3.00 and 1.28 years, respectively, compared to 7.25 and 8.10 years for EPS-P and EPS-D, respectively (log-rank $P = 0.013$). Median DFS for PD-CHO and STME were 0.73 and 1.01 years, respectively, compared to 2.78 and 5.48 years for EPS-P and EPS-D, respectively (log-rank $P = 0.023$) (Fig. 4A, B). The number of cases for eMPNST and MRT with survival data were too small for statistical comparison.

There was no significant impact on OS or DFS based on the presence of absence of co-occurring genetic alterations (possibly related to the low number of cases with co-occurring genetic hits), whether *SMARCB1* alterations were homozygous or heterozygous, or whether there was extended chromosome 22q deletions across all *SMARCB1*-deficient histotypes or within epithelioid sarcoma (Supplementary Fig. 1–3).

DISCUSSION

The mechanisms behind how *SMARCB1* biallelic inactivation within an otherwise quiet genome drives the pathogenesis of a wide spectrum of both benign and highly aggressive mesenchymal neoplasms remain unresolved. In order to gain further insight into this clinical heterogeneity, we performed genomic profiling of a large cohort of soft tissue and bone tumors, focusing on the interrelationship between the types of *SMARCB1* genetic alterations (copy number and/or mutational status), *SMARCB1* protein expression, coexisting additional genomic abnormalities and histotypes.

Among our cohort of *SMARCB1*-deficient sarcomas, in all but 5 cases (69, 93%) showing *SMARCB1* loss of expression at the protein level, *SMARCB1* copy number deletions or LOF mutations were detected by either one or both methods. However, among these 69 cases, 10 (14%) showed only monoallelic LOF mutations and 5 (7%) had no demonstrable alterations by both methods applied. The incidence and types of *SMARCB1* alterations show




Fig. 4 Survival of various histotypes of *SMARCB1*-deficient sarcomas. Kaplan Meier plots showing **A** overall survival and **B** disease-free survival of proximal-type epithelioid sarcoma ($n = 30$), distal-type epithelioid sarcoma ($n = 15$), soft tissue myoepithelial tumors ($n = 6$), and poorly differentiated chondromas ($n = 7$) [log-rank P value].

significant heterogeneity among histotypes. Table 1 summarizes this data from the published literature compared to the current study. Overall, for epithelioid sarcoma^{8, 12, 21, 22} and PD-CHO^{13, 14}, our data are largely consistent with prior findings that *SMARCB1* deletions predominate. In MRT, *SMARCB1* point mutations/intragenic deletions ranged from 55–60% in somatic cases and 71% in germline cases, whereas we observed 1 of 3 MRT with a nonsense point mutation^{15, 23, 24}. Interestingly, among epithelioid sarcomas, intragenic deletions or loss-of-function *SMARCB1* mutations (nonsense, frameshift, splice site) were only observed in EPS-P but not EPS-D. Also, four cases (29%) of EPS-D versus just one case (3%) of EPS-P showed no *SMARCB1* alterations despite *SMARCB1* protein loss. On the other hand, for eMPNST and eSCHW, we detected a higher percentage of cases with *SMARCB1* deletions compared to point mutations/intragenic deletions compared to the study by Schaefer 2019²⁵. Finally, for STME, *SMARCB1* point mutations/intragenic deletions predominate in our study (5 of 6). Although *SMARCB1* deficiency has been reported in a variant of extraskeletal myxoid chondrosarcoma (EMC)²⁶, of the 15 cases with DNA sequencing data available at our institution, we

did not identify any cases with *SMARCB1* deletions or mutations, and thus we did not include EMC in our study cohort.

We speculate that in a small subset of cases, germline LOF *SMARCB1* alterations might be missed by our MSK-IMPACT platform, which only detects somatic alterations. However, *SMARCB1* germline alterations are extremely rare outside of MRT¹². Moreover, cases with copy number neutral loss of heterozygosity on 22q with a *SMARCB1* LOF mutation may technically appear heterozygous. Further, the relatively low frequency of intragenic deletion detected in our study compared to Sullivan et al may be due to reduced sensitivity of targeted DNA sequencing compared to the gold standard method of multiplex ligation-dependent probe amplification MLPA, which was used in their study to detect intragenic deletion⁸. Additionally, *SMARCB1* gene expression could be perturbed at a post-transcriptional or epigenetic level. For example, studies have shown that *SMARCB1* gene expression in epithelioid sarcomas is regulated by specific microRNAs on both mRNA and protein levels, rather than promoter hypermethylation^{27, 28}. Conversely, in synovial sarcomas, *SMARCB1* protein is known to be reduced but not completely

lost in about 70% of cases without corresponding reduction in *SMARCB1* mRNA levels²⁹. This was postulated to be the result of a post-transcriptional mechanism where the SS18/SSX oncogenic fusion disrupts the SWI/SNF (BAF) complex by displacing wildtype SS18 from the BAF complex³⁰. This may account for the single case of synovial sarcoma in our study that showed “loss” of *SMARCB1* by IHC, which may be more accurately referred to as reduction in protein expression rather than loss.

Our study also investigated for the first time the sensitivity of FISH versus sequencing in detecting *SMARCB1* alterations in *SMARCB1*-deficient cases, showing concordant findings in 48% of cases for which FISH was available. FISH was used as a complementary method to detect large, arm-level copy number deletions of *SMARCB1*, which may not be reported in standard clinical workflow. Reasons that could account for the discrepant results between these two methods are likely related to the low resolution of FISH to detect copy number neutral cases, such as loss-of-function *SMARCB1* mutations or intragenic deletions (spanning one or more exons of a gene). By profiling the genes flanking *SMARCB1* on chromosome 22q, we also demonstrated that in about two-third of cases, *SMARCB1* was the only gene deleted on chromosome 22q. Thus, it is likely that FISH may not be able to detect some of these cases without extended 22q deletion.

Overall, in our cohort, LOF *SMARCB1* alterations predominate. Importantly, the significant number of cases with *SMARCB1* genetic alterations in the absence of *SMARCB1* loss at the protein level—mostly point mutations of uncertain significance (monoallelic missense or in-frame mutations)—suggests that the presence of a *SMARCB1* alteration detected by molecular testing does not necessarily imply a *SMARCB1*-deficient entity. Indeed, these *SMARCB1* mutations occurred in various sarcoma histotypes driven by known oncogenic drivers (such as fusions, gene amplifications), which typically are not associated with *SMARCB1* loss. The question remains if these *SMARCB1* monoallelic LOF alterations represent either passenger mutations or are functional in the context of co-occurring driver events. Among this group, a subset (11%) of cases showed *SMARCB1* amplifications. In at least 61% of these cases, copy number amplifications were seen at genes flanking *SMARCB1* on chromosome 22q, indicating that *SMARCB1* amplification could be a bystander event due to broad copy number gains on chromosome arm 22q.

Additionally, we demonstrated that clinically significant, recurrent co-occurring (secondary) genetic events were rare in *SMARCB1*-deficient sarcomas of various histotypes, regardless of outcome. This extremely low mutation rate has been well documented in malignant rhabdoid tumors¹⁵, but not in other mesenchymal neoplasms driven by the same *SMARCB1* deficiency. This is analogous to fusion-driven sarcomas, which tend to harbor relatively “quiet” genome with low tumor mutation burden³¹. These infrequent co-occurring events involved genes that are commonly altered in sarcomas: *TP53*, *RB1*, *CDKN2A/B*;³² *CDKN2A* deletions were also previously observed in epithelioid sarcomas²¹. However, in contrast to a prior study showing higher tumor mutation burden (TMB) in epithelioid sarcomas compared to MRT, we did not observe a significantly higher TMB in epithelioid sarcomas. This could be due to less precise TMB estimates for low to moderate TMB samples for targeted, panel-based NGS compared to whole exome or whole genome sequencing³³, the latter of which was used in the aforementioned study for TMB estimation²¹. In this same study, they observed *CDKN2A/B* deletion in 2 of 7 epithelioid sarcomas; we observed *CDKN2A/B* deletion in only 1 case of epithelioid sarcoma (out of 44 cases), a frequency no higher than other *SMARCB1*-deficient tumors (1 of 11 eMPNST, 1 of 7 PD-CHO). Further, an aCGH study showed *CDKN2A* deletion in a case of conventional chordoma but not in poorly differentiated chordoma (one case), which only showed 22q loss (spanning *SMARCB1* locus)³⁴. By contrast, a targeted NGS study reported 31% (4 of 16) of eMPNST and 20% (1 of 5) eSCHW

harboring *CDKN2A* deletion or nonsense mutation²⁵. On the other hand, we observed only *TP53* and *ARID1A* alterations in STME, and *RB1* alterations in epithelioid sarcoma only, in contrast to a case report of a STME showing *RB1* deletion³⁵. Overall, studies on co-occurring genetic alterations in *SMARCB1*-deficient sarcomas are largely limited to small case series and single case reports, therefore the true incidence of co-occurring genetic alterations in these tumors remains unclear. Nevertheless, in our cohort, the presence of co-occurring genetic alterations did not correlate with worse OS or DFS. Studies have demonstrated compound heterozygous mutations in more than one BAF complex subunits and intra-complex co-dependencies between different subunits of the BAF chromatin remodeling complex, e.g., *SMARCA4-ARID2*, *SMARCA4-ACTB*, etc^{2, 36}. On the MSK-IMPACT panel, the subunits of BAF complexes that were targeted are as follows: *ARID1A*, *ARID1B*, *ARID2*, *PBRM1*, *SMARCA2*, *SMARCA4*, *SMARCB1*, *SMARCD1*, *SMARCE1*. Of these, the only subunit that co-occurred with *SMARCB1* in our study was *ARID1A/B*.

Within EPS-P, longitudinal sequencing of recurrence/metastatic tumors in comparison to the initial sample showed that during tumor progression, acquisition of copy number alterations was frequent (4 of 5 cases) while mutations were less commonly acquired in subsequent recurrence/metastases (2 of 5 cases). Further, mutations in *ARID1B*, *TP53* and *BRCA2*, present in both the initial sample and subsequent recurrence/metastases, appeared to represent early genetic events, and the tumors maintained a relatively low overall tumor mutation burden even in late-stage disease. Interestingly, extended 22q copy number loss in genes flanking the *SMARCB1* locus (22q11.23) occurred in approximately one-third of epithelioid sarcomas and STME but in the majority of PD-CHO. There were no significant differences in the frequencies of coexisting mutations or extended 22q loss between EPS-P and EPS-D.

There is a paucity of large studies in the literature that specifically investigate the correlation between *SMARCB1* loss and myoepithelial tumor histology (specifically, benign vs malignant myoepithelial neoplasms). This is also compounded by the inconsistent nomenclature adopted in the literature, where “soft tissue myoepithelial tumor” or “myoepithelioma-like tumor” are used to signify malignant myoepithelial neoplasms², whereas other authors prefer the term “myoepithelial carcinoma” when referring to such neoplasms to distinguish them from benign myoepithelial tumors (i.e., myoepitheliomas)^{38, 39}. Nonetheless, the limited case series and case reports published thus far, as well as the STME cases in the current study, have demonstrated that *SMARCB1* loss are limited to malignant myoepithelial neoplasms^{9, 12, 37–39}. For example, in a study of myoepithelioma-like tumors of the vulvar region (MLTVR), *SMARCB1* IHC was lost in all 9 cases of MLTVR, but retained in all 11 cases of myoepitheliomas³⁷. To our knowledge, *SMARCB1* loss has not been reported in benign myoepithelial tumors^{4, 37}.

In summary, we demonstrated that in the majority of *SMARCB1*-deficient sarcomas, *SMARCB1* protein loss could be accounted for by the presence of biallelic *SMARCB1* deletion or LOF mutations. Further, *SMARCB1* IHC is a highly sensitive and specific surrogate for diagnostically and clinically relevant *SMARCB1* pathogenic alterations. However, a minority of cases only showed heterozygous or no detectable *SMARCB1* copy number alterations or mutations. Recurrent co-occurring genetic events are relatively rare, indicating that *SMARCB1* deficiency is the primary oncogenic driver. A subset of cases also harbored extended 22q deletion. Among *SMARCB1*-deficient sarcomas, PD-CHO and STME showed worse prognosis than epithelioid sarcomas.

DATA AVAILABILITY

The raw data generated are not publicly available due to lack of access to indefinite hosting capabilities, but are available from the corresponding author on reasonable request.

REFERENCES

- Nakayama, RT, Pulice, JL, Valencia, AM, McBride, MJ, McKenzie, ZM, Gillespie, MA, et al. SMARCB1 is required for widespread BAF complex-mediated activation of enhancers and bivalent promoters. *Nat Genet* **49**, 1613–1623 (2017)
- Kadoch, C, Hargreaves, DC, Hodges, C, Elias, L, Ho, L, Ranish, J, Crabtree, GR. Proteomic and bioinformatic analysis of mammalian SWI/SNF complexes identifies extensive roles in human malignancy. *Nat Genet* **45**, 592–601 (2013)
- McBride, MJ, & Kadoch, C. Disruption of mammalian SWI/SNF and polycomb complexes in human sarcomas: mechanisms and therapeutic opportunities. *J Pathol* **2018**;244 (5), 638–649.
- Hollmann, TJ, Hornick, JL. INI1-deficient tumors: diagnostic features and molecular genetics. *Am J Surg Pathol* **35**, 47–63 (2011)
- Nacev, BA, Jones, KB, Intlekofer, AM, Yu, JSE, Allis, CD, Tap, WD, et al. The epigenomics of sarcoma. *Nat Rev Cancer* **20**, 608–623 (2020)
- Parker, AI, Obaidi, A, Deutsch, JM. SMARCB1/INI1-deficient tumors of adulthood. *F1000Res* **9**, 662 (2020).
- Pawel, BR. SMARCB1-deficient Tumors of Childhood: A Practical Guide. *Pediatr Dev Pathol* **21**, 6–28 (2018)
- Sullivan, LM, Folpe, AL, Pawel, BR, Judkins, AR, Biegel, JA. Epithelioid sarcoma is associated with a high percentage of SMARCB1 deletions. *Mod Pathol* **26**, 385–392 (2013)
- Hornick, JL, Dal Cin, P, Fletcher, CD. Loss of INI1 expression is characteristic of both conventional and proximal-type epithelioid sarcoma. *Am J Surg Pathol* **33**, 542–550 (2009)
- Jo, VY, Fletcher, CD. Epithelioid malignant peripheral nerve sheath tumor: clinicopathologic analysis of 63 cases. *Am J Surg Pathol* **39**, 673–682 (2015)
- Jo, VY, Fletcher, CDM. SMARCB1/INI1 Loss in Epithelioid Schwannoma: A Clinicopathologic and Immunohistochemical Study of 65 Cases. *Am J Surg Pathol* **41**, 1013–1022 (2017)
- Le Loarer, F, Zhang, L, Fletcher, CD, Ribeiro, A, Singer, S, Italiano, A, et al. Consistent SMARCB1 homozygous deletions in epithelioid sarcoma and in a subset of myoepithelial carcinomas can be reliably detected by FISH in archival material. *Genes Chromosomes Cancer* **53**, 475–486 (2014)
- Mobley, BC, McKenney, JK, Bangs, CD, Callahan, K, Yeom, KW, Schneppenheim, R, Hayden, MG, Cherry, AM, Gokden, M, Edwards, MS, Fisher, PG, Vogel, H. Loss of SMARCB1/INI1 expression in poorly differentiated chordomas. *Acta Neuropathol* **120**, 745–753 (2010)
- Owosho, AA, Zhang, L, Rosenblum, MK, Antonescu, CR. High sensitivity of FISH analysis in detecting homozygous SMARCB1 deletions in poorly differentiated chordoma: a clinicopathologic and molecular study of nine cases. *Genes Chromosomes Cancer* **57**, 89–95 (2018)
- Lee, RS, Stewart, C, Carter, SL, Ambrogio, L, Cibulskis, K, Sougnez, C, et al. A remarkably simple genome underlies highly malignant pediatric rhabdoid cancers. *J Clin Invest* **122**, 2983–2988 (2012)
- Custers, L, Khabirova, E, Coorens, THH, Oliver, TRW, Calandrini, C, Young, MD, et al. Somatic mutations and single-cell transcriptomes reveal the root of malignant rhabdoid tumours. *Nat Commun* **12**, 1407 (2021).
- Cheng, D.T., Mitchell, T.N., Zehir, A., Shah, R.H., Benayed, R., Syed, A., et al. Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor Molecular Oncology. *J Mol Diagn* **17**, 251–264 (2015).
- Chakravarty, D, Gao, J, Phillips, SM, Kundra, R, Zhang, H, Wang, J, et al. OncoKB: A Precision Oncology Knowledge Base. *JCO Precis Onco* **2017**, PO.17.00011 (2017).
- Gu, Z, Eils, R, Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016)
- Skidmore, ZL, Wagner, AH, Lesurf, R, Campbell, KM, Kunisaki, J, Griffith, OL, et al. “GenVisR: Genomic Visualizations in R.” *Bioinformatics* **32**, 3012–3014 (2016)
- Jamshidi, F, Bashashati, A, Shumansky, K, Dickson, B, Gokgoz, N, Wunder, JS, et al. The genomic landscape of epithelioid sarcoma cell lines and tumours. *J Pathol* **238**, 63–73 (2016)
- Thway, K, Jones, RL, Noujaim, J, Fisher, C. Epithelioid Sarcoma: Diagnostic Features and Genetics. *Adv Anat Pathol* **23**, 41–49 (2016)
- Jackson, EM, Sievert, AJ, Gai, X, Hakonarson, H, Judkins, AR, Tooke, L, et al. Genomic analysis using high-density single nucleotide polymorphism-based oligonucleotide arrays and multiplex ligation-dependent probe amplification provides a comprehensive analysis of INI1/SMARCB1 in malignant rhabdoid tumors. *Clin Cancer Res* **15**, 1923–1930 (2009)
- Eaton, KW, Tooke, LS, Wainwright, LM, Judkins, AR, Biegel, JA. Spectrum of SMARCB1/INI1 mutations in familial and sporadic rhabdoid tumors. *Pediatr Blood Cancer* **56**, 7–15 (2011)
- Schaefer, IM, Dong, F, Garcia, EP, Fletcher, CDM, Jo, VY. Recurrent SMARCB1 Inactivation in Epithelioid Malignant Peripheral Nerve Sheath Tumors. *Am J Surg Pathol* **43**, 835–843 (2019)
- Kohashi, K, Oda, Y, Yamamoto, H, Tamiya, S, Oshiro, Y, Izumi, T, et al. SMARCB1/INI1 protein expression in round cell soft tissue sarcomas associated with chromosomal translocations involving EWS: a special reference to SMARCB1/INI1 negative variant extraskeletal myxoid chondrosarcoma. *Am J Surg Pathol* **32**, 1168–1174 (2008)
- Papp, G, Krausz, T, Stricker, TP, Szendrői, M, Sápi, Z. SMARCB1 expression in epithelioid sarcoma is regulated by miR-206, miR-381, and miR-671-5p on both mRNA and protein levels. *Genes Chromosomes Cancer* **53**, 168–176 (2014)
- Sápi, Z, Papp, G, Szendrői, M, Pápai, Z, Plótár, V, Krausz, T, et al. Epigenetic regulation of SMARCB1 By miR-206, -381 and -671-5p is evident in a variety of SMARCB1 immunonegative soft tissue sarcomas, while miR-765 appears specific for epithelioid sarcoma. A miRNA study of 223 soft tissue sarcomas. *Genes Chromosomes Cancer* **55**, 786–802 (2016)
- Kohashi, K, Oda, Y, Yamamoto, H, Tamiya, S, Matono, H, Iwamoto, Y, et al. Reduced expression of SMARCB1/INI1 protein in synovial sarcoma. *Mod Pathol* **23**, 981–990 (2010)
- Kadoch, C, Crabtree, GR. Reversible disruption of mSWI/SNF (BAF) complexes by the SS18-SSX oncogenic fusion in synovial sarcoma. *Cell* **153**, 71–85 (2013)
- Mertens, F, Antonescu, CR, Mitelman, F. Gene fusions in soft tissue tumors: Recurrent and overlapping pathogenetic themes. *Genes Chromosomes Cancer* **55**, 291–310 (2016)
- Cancer Genome Atlas Research Network. Comprehensive and Integrated Genomic Characterization of Adult Soft Tissue Sarcomas. *Cell* **171**, 950–965.e28 (2017)
- Buchhalter, I, Rempel, E, Endris, V, Allgäuer, M, Neumann, O, Volckmar, AL, et al. Size matters: Dissecting key parameters for panel-based tumor mutational burden analysis. *Int J Cancer* **144**, 848–858 (2019)
- Shih, AR, Chebib, I, Deshpande, V, Dickson, BC, lafrate, AJ, Nielsen, GP. Molecular characteristics of poorly differentiated chordoma. *Genes Chromosomes Cancer* **58**, 804–808 (2019)
- Hoggard, TM, Henderson-Jackson, E, Bui, MM, Caracciolo, J, Teer, JK, et al. Myoepithelial carcinoma with RB1 mutation: remarkable chemosensitivity to carcinoma of unknown origin therapy. *BMC Cancer* **17**, 250 (2017)
- Schick, S, Rendeiro, AF, Runggatscher, K, Ringler, A, Boidol, B, Hinkel, M, et al. Systematic characterization of BAF mutations provides insights into intracomplex synthetic lethality in human cancers. *Nat Genet* **51**, 1399–1410 (2019)
- Yoshida, A, Yoshida, H, Yoshida, M, Mori, T, Kobayashi, E, Tanzawa, Y, et al. Myoepithelioma-like Tumors of the Vulvar Region: A Distinctive Group of SMARCB1-deficient Neoplasms. *Am J Surg Pathol* **39**, 1102–1113 (2015)
- Folpe, AL, Schoolmeester, JK, McCluggage, WG, Sullivan, LM, Castagna, K, Ahrens, WA, et al. SMARCB1-deficient Vulvar Neoplasms: A Clinicopathologic, Immunohistochemical, and Molecular Genetic Study of 14 Cases. *Am J Surg Pathol* **39**, 836–849 (2015)
- Agaimy, A, Naroditsky, I, Ben-Izhak, O. Primary high-grade myoepithelial carcinoma of the lung: A study of three cases illustrating frequent SMARCB1-deficiency and review of the literature. *Ann Diagn Pathol* **53**, 151759 (2021)

ACKNOWLEDGEMENTS

We gratefully acknowledge the members of the Molecular Diagnostics Service in the Department of Pathology and would like to acknowledge the Center Core grant (P30 CA008748) and the Marie-Josee and Henry R. Kravis Center for Molecular Oncology for use of MSK-IMPACT data. We acknowledge Lei Zhang for her assistance with FISH assays. We would also like to thank Dr. Sharon Weiss for generously providing us with cases of epithelioid schwannoma and epithelioid malignant peripheral nerve sheath tumor as part of the study cohort. This work was supported by P50 CA217694 (SS, WT, CRA), P30 CA008748 (SS, WT, CRA, MVO), Kristin Ann Carr Foundation (CRA).

AUTHOR CONTRIBUTIONS

JKD performed study design, data acquisition, data analysis and interpretation, writing and revision of the paper. SS, WT, BN, PC, LHW, MVO, and MG performed data acquisition and critical review of the paper. CRA performed study design and conception, analysis and interpretation of data, writing, review and revision of paper. All authors read and approved the final manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

ETHICS APPROVAL

This study was approved by the Memorial Sloan Kettering Cancer Institute Institutional Review Board.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41379-022-01148-x>.

Correspondence and requests for materials should be addressed to Cristina R. Antonescu.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.