



## REVIEW ARTICLE

# Perinatal and childhood exposure to environmental chemicals and blood pressure in children: a review of literature 2007–2017

Alison P. Sanders<sup>1,2</sup>, Jeffrey M. Saland<sup>2</sup>, Robert O. Wright<sup>1,2</sup> and Lisa Satlin<sup>2</sup>

Exposure to environmental chemicals during periods of renal development from embryogenesis to birth and through childhood can inform critical windows of nephrotoxicity, including changes in childhood blood pressure. This review assessed recent studies that examined the relationship of air pollution, metals, and other organic pollutants with children's blood pressure outcomes. We restricted this review to peer-reviewed studies published in English between January 2007 and July 2017. We identified a total of 36 articles that estimated associations with childhood blood pressure, of which 14 studies examined the effects of air pollution, 10 examined metals, and 12 examined other organic pollutants including phthalates ( $n = 4$ ), Bisphenol A ( $n = 3$ ), polychlorinated biphenols ( $n = 2$ ), organophosphate pesticides ( $n = 2$ ), or perfluoroalkyl acids ( $n = 1$ ). Similar to the established relationship between tobacco smoke exposure and childhood blood pressure, the majority of studies that examined air pollutants, particularly exposure to PM<sub>10</sub> and PM<sub>2.5</sub>, reported associations with increased childhood blood pressure. The literature reported conflicting evidence for metals, and putative evidence of the effects of exposure to phthalates, Bisphenol A, polychlorinated biphenols, and pesticides. Overall, our review underscores the need for additional studies that assess the impact of nephrotoxicant exposure during early life, particularly the perinatal period, and blood pressure in childhood.

*Pediatric Research* (2018) 84:165–180; <https://doi.org/10.1038/s41390-018-0055-3>

## INTRODUCTION

Hypertension (HTN), an important risk factor for cardiovascular disease, affects ~30% of adults 18 years of age and older in the United States.<sup>1</sup> The prevalence of HTN in children, in contrast, is 3.5%, although it ranges from 3.8 to 24.8% in overweight or obese youth populations<sup>2</sup> (reviewed in<sup>3</sup>). Its prevalence is thus rising as child obesity rates increase.<sup>2,4</sup> Of concern, later life HTN is strongly correlated with early life blood pressure (BP) elevation and impaired renal development.<sup>5,6</sup>

The etiology of HTN is complex, and its programming is considered to reflect a complex interplay between genetic predisposition and environment, mediated in part by epigenetic factors. Although it is generally accepted that overweight/obesity,<sup>7</sup> a sedentary lifestyle,<sup>8,9</sup> and salt intake<sup>10</sup> contribute to HTN, environmental chemical exposures to heavy metals, phthalates, and arsenic are also established risk factors for development of HTN in adult populations, estimated to account for 3–19% of the population attributable risk for high BP.<sup>11</sup> Despite the predictive value of childhood BP on HTN later in life, there exist a limited number of studies that examine the effect of environmental chemical exposures on childhood BP.

The developmental origins of health and disease (DOHaD) hypothesis posits that an adverse intrauterine environment programs chronic disease including elevated BP and kidney disease later in life.<sup>12</sup> Metanephric kidney development typically begins at 5 weeks' gestation with the full complement of ~1 million nephrons per kidney largely achieved by ~35 weeks gestation.<sup>13,14</sup> Renal

blood flow and glomerular filtration, which increase significantly during fetal life, continue to mature after birth, reaching adult levels by 2 years of age.<sup>15</sup> Overall, these prenatal and postnatal periods are likely to be critical windows that are susceptible to nephrotoxic environmental chemical exposures predisposing the developing kidney for later life HTN (Fig. 1).<sup>16,17</sup> Although evidence supports this relationship for in utero tobacco exposure and child BP,<sup>18,19</sup> the effects of other highly prevalent environmental nephrotoxicants on renal development and BP are poorly understood. We previously reviewed non-BP effects of environmental chemicals on pediatric kidney function and disease.<sup>20</sup>

To summarize the state of the literature on the role of environmental chemicals as predictors of childhood BP, we conducted a review of human studies examining exposures during prenatal and postnatal susceptibility windows. We focused on exposure to air pollution, and included a wider range of publication dates and pollutants than previously reported.<sup>21,22</sup> We also included metals and organic contaminants as pervasive and/or emerging contaminants, which have not been reviewed with respect to childhood BP. We present conclusions based on this body of evidence and provide recommendations for future studies.

## METHODS


We conducted a literature search to identify human studies that examined the effects of air pollutants [particulate matter

<sup>1</sup>Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA and <sup>2</sup>Department of Pediatrics, Icahn School of Medicine at Mount Sinai, New York, NY, USA

Correspondence: Alison P. Sanders ([alison.sanders@mssm.edu](mailto:alison.sanders@mssm.edu))

Received: 16 February 2018 Revised: 26 April 2018 Accepted: 27 April 2018

Published online: 9 June 2018



**Fig. 1** The timing of exposures to environmental chemicals during periods of renal development from embryogenesis to birth and through childhood can inform critical windows of kidney susceptibility. Printed with permission of ©Mount Sinai Health System

(PM<sub>10</sub> and PM<sub>2.5</sub>), nitric oxides (NO<sub>x</sub> and NO<sub>2</sub>), ozone (O<sub>3</sub>), sulfur dioxide (SO<sub>2</sub>), carbon monoxide (CO), black carbon (BC), and polycyclic aromatic hydrocarbons (PAHs)], nephrotoxic metals [arsenic (As), cadmium (Cd), lead (Pb), or mercury (Hg)], phthalates and other organic contaminants with respect to children's BP using the following search terms (child\* OR infan\* OR school OR postnatal OR post-natal OR prenatal OR pre-natal OR fetal OR pregnan\* OR in utero) AND (blood pressure OR systolic OR diastolic OR pulse pressure) AND (metal OR "cadmium" [MeSH Terms] OR "arsenic" [MeSH Terms] OR "lead" [MeSH Terms] OR "mercury" [MeSH Terms] OR "BPA" [MeSH Terms] OR "Bisphenol A" [MeSH Terms] OR perfluoroalkyl OR pesticides OR PAH OR polycyclic aromatic hydrocarbon OR phthalate). Among additional contaminants included in the initial search terms including cannabis/marijuana, brominated flame retardants, parabens, dioxins, or furans, we identified no published studies of children's BP. We used PubMed and Web of Science search engines and restricted the search to peer-reviewed human studies published in English between January 2007 and July 2017. We summarized a total of 14, 10, and 12 studies on air pollution, metals, or phthalates and other organic contaminants, respectively. We note that this manuscript is not a systematic review, which was not possible due to the paucity of peer-reviewed studies for some exposures as well as the heterogeneity of reported exposure and outcome measures.

This review uses updated terminology for pediatric BP as defined in the 2017 report by the American Academy of Pediatrics (AAP) Subcommittee On Screening Management Of High Blood Pressure In Children and Adolescents<sup>2</sup> wherein normal BP, elevated BP (formerly called "preHTN"), Stage 1 HTN and Stage 2 HTN are defined and we refer the reader to the AAP report for details.<sup>2</sup> The term "high BP" may refer to both HTN and "elevated BP" together. To further clarify BP nomenclature, we use the term "increased" BP to describe associations with continuous BP (formerly termed "elevated" in some studies). In cases where the former terminology was used in reports herein, we have indicated the authors' intended definition of "HTN", "preHTN", and "elevated BP" and corrected linear regression interpretations to "increased BP" where appropriate.

Epidemiologic evidence of association between air pollution and childhood BP

We identified 14 studies examining the effects of air pollutants on child BP (Table 1). Exposures assessed included PM<sub>10</sub> (n = 10),

PM<sub>2.5</sub> (n = 8), NO<sub>x</sub> or NO<sub>2</sub> (n = 8), O<sub>3</sub> (n = 5), SO<sub>2</sub> (n = 3), CO (n = 2), BC (n = 2), PAHs (n = 1), distance to roadway/traffic/point source (n = 3), nano-sized ultrafine particles (n = 1); eleven studies assessed more than one pollutant. Outdoor air quality was assessed via ambient monitoring, estimates derived with geographic information system (GIS) models (distance to traffic, land-use regression), or satellite/dispersion or other national database estimates. Two studies assessed indoor air quality. Three of the study designs were prospective and 12 were cross-sectional. Among 10 studies of PM<sub>10</sub> and 8 studies of PM<sub>2.5</sub>, four and two reported significant associations with childhood BP, respectively. Five of seven studies of NO<sub>2</sub>, four of five studies of O<sub>3</sub>, two of three studies of SO<sub>2</sub>, one of two studies of CO, and one of two studies of BC reported associations with childhood BP; one study reported evidence of a relationship with nano-sized ultrafine particles (UFP) (particles with a diameter ≤100 nm). Two studies examined indoor air pollution and reported no association with children's BP.<sup>23,24</sup> A single study using school location as proxy for PAH exposure reported an association with childhood BP, but not with urine PAH biomarkers.<sup>25</sup> A study using distance to major road as a proxy for ambient PM<sub>10</sub> reported higher mean carotid arterial stiffness (derived from BP and ultrasound parameters) among children living closer to the major road.<sup>26</sup> A study of road traffic at schools near an airport found no association with child BP.<sup>27</sup>

Among the three prospective cohort studies, all identified significant associations with exposure to air pollution (PM<sub>2.5</sub> and other pollutants) and child BP. In a prospective study of 1131 infants born in the United States, higher mean PM<sub>2.5</sub> and BC in the third trimester (90 days before birth) were associated with higher SBP in newborns ~30 h after birth.<sup>28</sup> Ozone (O<sub>3</sub>) was associated with lower SBP, and no significant associations were observed for NO<sub>x</sub>, NO<sub>2</sub>, or CO averages during the third trimester. The observed significant third trimester findings with PM<sub>2.5</sub>, BC, and O<sub>3</sub> were bolstered by estimates of "moving averages" calculated from 2 to 7, 14, 30, 60, and 90 days before birth. No significant relationships were observed with first trimester pollutant averages; however, putative relationships in the second trimester were observed for NO<sub>x</sub> and CO with lower SBP. The study did not report associations with DBP. Two studies conducted in the PIAMA birth cohort in the Netherlands<sup>29,30</sup> reported associations between PM<sub>2.5</sub> and increased DBP at 12 years. In a subset of 471 children who did not move since birth, long-term NO<sub>2</sub> and PM<sub>2.5</sub> (reported as

**Table 1.** Studies assessing exposure to air pollutants and blood pressure outcomes (*n* = 14)

Authors and Year	Study design, cohort name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Confounders	Major finding(s)
Van Rossem et al. 2015 <sup>28</sup>	Prospective cohort Project Viva	US	1131	PM <sub>2.5</sub> , NO <sub>x</sub> , NO <sub>2</sub> , O <sub>3</sub> , CO, BC measured ambient monitoring station and averaged by trimester, and 2–90 days before birth. Spatiotemporally resolved estimates for PM <sub>2.5</sub> and BC.	SBP and DBP in newborns (30 ± 18 h after birth) with an automated device, measured 5 times 1-min apart, recorded measurement conditions (i.e., sleep/wake, cuff size, arm, position)	Measurement circumstance, birth weight, maternal age, race/ethnicity, SES, education level, third trimester BP time trend	An IQR increase in BC (0.33 µg/m <sup>3</sup> ) in the 3rd trimester was associated with SBP (1.0 mmHg; 95% CI: 0.2, 1.8). Similar trends observed for mean PM <sub>2.5</sub> and BC with higher SBP during the 2–90 days interval before birth. An IQR increase in O <sub>3</sub> (13.6 ppb) in the 3rd trimester was inversely associated with SBP (−2.5 mmHg; 95% CI: −4.5, −0.4). Similar trends observed for averages over 2–90 days before birth. Estimates with DBP not reported.
Bilenko et al. 2015a <sup>29</sup>	Prospective cohort PIAMA	Netherlands	1147 (subset of 471 non-movers)	Annual average levels of NO <sub>2</sub> [median: 21.8 µg/m <sup>3</sup> ; IQR: 7.6] and Cu, Fe, K, Ni, S, Si, V, and Zn components of PM <sub>2.5</sub> [median: 16.5 µg/m <sup>3</sup> ; IQR: 1.2], PM <sub>10</sub> [median: 24.7 µg/m <sup>3</sup> ; IQR 1.0] estimated by Land-Use Regression at birth and at time of BP measure.	SBP and DBP at 12 years with an automated device, sized cuff, on the non-dominant arm while seated, measured two times 5-min apart.	Sex, age, height, BMI, cuff size, weight gain 1st yr, breast feeding, maternal smoking in pregnancy, parental smoking in home, physical activity, puberty scale, maternal education, ambient temperature, and room temperature. Stratified by non-movers vs. movers	Among non-movers, an IQR increase in NO <sub>2</sub> (7.6 µg/m <sup>3</sup> ) was associated with increased DBP (0.82 mmHg, 95% CI: 0.08, 1.57). *Replicated. Among non-movers, an IQR increase in iron (119.2 ng/m <sup>3</sup> ), silicon (74.5 ng/m <sup>3</sup> ), and potassium (18.3 ng/m <sup>3</sup> ) in PM <sub>10</sub> were associated with increased DBP (0.67 mmHg, 95% CI 0.02, 1.31; 0.85 mmHg, 95% CI: 0.18, 1.52; and 0.75 mmHg, 95% CI: 0.09, 1.41). Marginal associations with iron and silicon in PM <sub>2.5</sub> were reported with DBP ( <i>p</i> < 0.1).
Bilenko et al. 2015b <sup>30</sup>	Prospective cohort PIAMA	Netherlands	1432 (471 non-movers)	NO <sub>2</sub> , PM <sub>2.5</sub> , and PM <sub>10</sub> , and PM <sub>2.5</sub> absorbance (PM <sub>2.5,abs</sub> ) assessed by ambient monitors at home and school for “short-term” (7-day average preceding BP measure) or “long-term” (annual average calculated by Land-Use Regression). Long-term: NO <sub>2</sub> [median, IQR: 9.6, 17.5–25.3 µg/m <sup>3</sup> ] PM <sub>2.5</sub> [16.5, 15.6–16.7 µg/m <sup>3</sup> ], PM <sub>10</sub> [24.5, 24.0–25 µg/m <sup>3</sup> ], and PM <sub>2.5,abs</sub> [1.2, 1.0–1.3 (10 <sup>−5</sup> m)]	SBP and DBP at 12 years with an automated device, sized cuff, on the non-dominant arm while seated, measured two times 5-min apart.	Sex, age, height, BMI, cuff size, GA at birth, birth weight, weight gain 1st year, breast feeding, maternal smoking in pregnancy, parental smoking in home, physical activity, puberty scale, maternal education, and room temperature. Stratified by non-movers vs. movers, and sex.	Among non-movers, an IQR increase in long-term NO <sub>2</sub> (7.8 µg/m <sup>3</sup> ) and PM <sub>2.5,abs</sub> (0.3 × 10 <sup>−5</sup> m) were associated with increased DBP (0.83, 95% CI: 0.06, 1.61 and 0.75, 95% CI −0.08, 1.58, respectively), but not SBP. No association with short-term exposures.
Zeng et al. 2017 <sup>31</sup>	Cross-sectional SNECCS	China	9354 (4771 elementary and 4583 middle school-aged children)	PM <sub>10</sub> [median: 108.6 µg/m <sup>3</sup> ], SO <sub>2</sub> [24.6 µg/m <sup>3</sup> ], NO <sub>2</sub> [25.5 µg/m <sup>3</sup> ], and O <sub>3</sub> [69.4 µg/m <sup>3</sup> ] 1–5 days preceding BP examine measured by ambient air monitors	SBP and DBP at age 5–17 years using a standard mercury sphygmomanometer after 5 min rest, seated, on the right arm, three times at 2-min intervals. Elevated BP defined as either SBP or DBP >95th percentile based on sex, age, and height (4th report guides) [66].	Temperature, age, sex, BMI, breast feeding, birth weight, exercise time, passive smoke exposure, parental education, family income, family history of HTN, and district.	And IQR increase in 5-day mean PM <sub>10</sub> (47.4 µg/m <sup>3</sup> ) and O <sub>3</sub> (51.4 µg/m <sup>3</sup> ) was associated with increased odds ratio of elevated BP 2.17 (95% CI: 1.61, 2.93) and 2.77 (95% CI: 1.94, 3.95), respectively and increased SBP of 2.07 mmHg (95% CI: 1.71, 2.44) and 3.29 mmHg (95% CI: 2.86, 3.72).

Table 1 continued

Authors and Year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Confounders	Major finding(s)
Dong et al. 2015 <sup>33</sup>	Cross-sectional SNECCS	China	9354 Categorized as normal weight (6311), overweight (1479) and obese (1564)	4-year average PM <sub>10</sub> , SO <sub>2</sub> , NO <sub>2</sub> , and O <sub>3</sub> measured at ambient monitoring stations [see below].	SBP and DBP at age 5–17 years using a standard mercury sphygmomanometer after 5 min rest, seated, on the right arm, three times at 2-min intervals. HTN defined as SBP and DBP ≥95th percentile for gender, age, and height (4th report guides) <sup>66</sup> .	Age, sex, parental education, LBW, PTB, breast feeding, income, passive smoke exposure, home coal use, exercise time, home size, family HTN history, distance to school, distance between air monitor and school, temperature, and district.	respectively. Significant associations also observed for DBP, but not reported. Stratified by sex, an IQR increase in NO <sub>2</sub> (18.5 µg/m <sup>3</sup> ) was associated with elevated BP in males (OR: 1.38; 95% CI: 1.00, 1.91), but not females. No associations with SO <sub>2</sub> . Among normal weight children, an IQR increase in PM <sub>10</sub> (30.6 µg/m <sup>3</sup> ), NO <sub>2</sub> (13.0 µg/m <sup>3</sup> ), and O <sub>3</sub> (46.3 µg/m <sup>3</sup> ) were associated with increased odds of HTN [OR: 1.21 95% CI: 1.07, 1.38; OR: 1.19, 95% CI: 1.07, 1.38; OR: 1.08 95% CI: 1.06, 1.10, respectively. Among normal weight children, a IQR increase of SO <sub>2</sub> (23.4 µg/m <sup>3</sup> ) was associated with decreased odds of HTN (0.82 95% CI: 0.72, 0.93). No associations were observed among normal weight children and SBP or DBP. There was evidence of an interaction effect with overweight and obese groups for PM <sub>10</sub> , SO <sub>2</sub> , and O <sub>3</sub> and both SBP and DBP.
Dong et al. 2014 <sup>32</sup>	Cross-sectional SNECCS	China	9354	4-year average PM <sub>10</sub> [median: 90.4 µg/m <sup>3</sup> ], SO <sub>2</sub> [48.4 µg/m <sup>3</sup> ], NO <sub>2</sub> [35.0 µg/m <sup>3</sup> ], O <sub>3</sub> [43.8 µg/m <sup>3</sup> ], and CO [1289.5 µg/m <sup>3</sup> ] measured at ambient monitoring stations.	HTN defined as SBP and DBP ≥95 <sup>th</sup> percentile for gender, age and height (4 <sup>th</sup> report guides) <sup>66</sup> .	Age, sex, BMI, parental education, LBW, PTB, breastfeeding, income, passive smoking exposure, home coal use, physical activity, home sq ft per person, family HTN history, and district. Stratified by breast feeding status and sex.	Increases in O <sub>3</sub> , PM <sub>10</sub> , CO, SO <sub>2</sub> , and NO <sub>2</sub> were associated with increased odd ratios for HTN. An IQR increase in O <sub>3</sub> (46.3 µg/m <sup>3</sup> ) was associated with an OR for HTN of 1.12 (95% CI: 1.10–1.13). An IQR increase in PM <sub>10</sub> (30.6 µg/m <sup>3</sup> ) was associated with an OR for HTN of 1.68 (95% CI: 1.53–1.86). An IQR increase in O <sub>3</sub> (46.3 µg/m <sup>3</sup> ) was associated with a 0.50 mmHg increase in SBP (95% CI: 0.43, 0.57) and a 0.58 mmHg increase in DBP (95% CI: 0.52, 0.63). An IQR increase in PM <sub>10</sub> (30.6 µg/m <sup>3</sup> ) was associated with a 2.10 mmHg increase in SBP (95% CI: 1.73, 2.47) and a 1.93 increase in DBP (95% CI 1.62, 2.25). An IQR increase in CO (563.4 µg/m <sup>3</sup> ) was associated with a 2.89 mmHg increase in DBP (95% CI: 2.53, 3.24). Effect sizes remained significant and of greater magnitude when restricted to non-breastfed group.

**Table 1** continued

Authors and Year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Confounders	Major finding(s)
Liu et al. 2013 <sup>34</sup>	Cross-sectional GINIplus & LISApilus	Germany	2368	"Long-term" (1-year) averages from ambient monitors of traffic-related NO <sub>2</sub> [median: 21.5 µg/m <sup>3</sup> ], PM <sub>2.5</sub> [median: 13.97 µg/m <sup>3</sup> ], PM <sub>10</sub> [median: 21.77 µg/m <sup>3</sup> ], and PM <sub>2.5</sub> abs. [median: 1.52 × 10 <sup>-5</sup> m] and for 'short-term' (7-day) for NO <sub>2</sub> [mean: 27.7 µg/m <sup>3</sup> ] and PM <sub>10</sub> [mean: 24.6 µg/m <sup>3</sup> ]	SBP and DBP at 10 years, measured using an automated device after 5 min rest, using a sized cuff, twice on the right arm 2-min apart while seated.	Study site, gender, age, BMI, physical activity, maternal smoking in pregnancy, parental education, parental HTN history, 7-day air pollutant, 7-day temperature.	No associations were observed for NO <sub>2</sub> , PM <sub>2.5</sub> or PM <sub>10</sub> . Noise was significantly associated with increased DBP.
Clark et al. 2012 <sup>27</sup>	Cross-sectional RANCH	UK	719 (subset of 276 with NO <sub>2</sub> )	Traffic-related noise and NO <sub>2</sub> at school locations near London Heathrow airport	SBP and DBP at 9–10 years using an automated device with sized cuffs, seated after 5 min of rest; three measures on the right arm with 1–2 min intervals (reported in <sup>36</sup> ).	Age, gender, employment status, crowding, home ownership, maternal education, chronic illness, language spoken, parental support, classroom window glaze, as well as preterm birth (<36 weeks), parental high BP, low birth weight (<2500 g), cuff size, temperature during testing, and BMI	Among children with air pollution data, no associations were observed for NO <sub>2</sub> with SBP or DBP. No associations observed for road traffic noise or aircraft noise with SBP or DBP.
Trasande et al. 2015 <sup>25</sup>	Cross-sectional	Saudi Arabia	184 (males only)	Ambient levels of TSP [mean 444.09 µg/m <sup>3</sup> ] and 14 PAHs [summed mean 36.75 ng/m <sup>3</sup> ] were monitored at three school locations at varying proximity to oil refinery. [concentration shown is for closest school] PAH metabolites were measured in urine (1-hydroxypyrene [median: 261.4 ng/g creatinine] and 1-hydroxyphenanthrene [549.6 ng/g creatinine])	SBP and DBP at 10–14 years using a standard cuff sphygmomanometer and DynaPulse instrument. BP Z-scores were calculated according to age, height, and sex (4th report) <sup>36</sup> . PreHTN defined as z-score SBP or DBP >90th percentile.	BMI z-score, SES, smokers in home (yes/no), use of incense, and age.	Proximity to oil refinery was associated with a 0.55 mmHg (95% CI: 0.26, 0.84) increase in z-score DBP and a 0.56 mmHg (95% CI: 0.16, 0.96) increase in SBP z-score. Significant associations were also observed with continuous SBP and DBP. A 4.35-fold odds of preHTN (95% CI: 1.42, 13.3) was associated with attending school closest to the refinery compared to the background school. Urine biomarkers were not significantly associated with BP outcomes. No significant associations observed for indoor PM <sub>2.5</sub> or BC with SBP or DBP.
Baumgartner et al. 2012 <sup>24</sup>	Cross-sectional	China	180	Indoor 24-h PM <sub>2.5</sub> [range: 14–393 µg/m <sup>3</sup> ] and BC [range: 2.0–9.6 µg/m <sup>3</sup> ] measured with personal monitors in rural households (reported in ref. <sup>35</sup> )	SBP and DBP at 5–14 years using an automated device.	Sex, age, height, BMI, passive smoking, SES, salt intake, MSG use, physical activity, day of week, time and day of BP measure	Higher mean SBP ( <i>p</i> < 0.0001) and DBP ( <i>p</i> < 0.0038) observed in areas with high traffic pollution vs. low. Increased, and marginally significant odds ratio of elevated BP (2.56; 95% CI: 0.96, 6.78).
Sughis et al. 2012 <sup>23</sup>	Cross-sectional	Pakistan	166 Categorized as low- ( <i>n</i> = 73) or high- ( <i>n</i> = 93) traffic areas	Indoor and outdoor PM <sub>10</sub> [mean "low" vs. "high" 223.0 vs. 728.6 µg/m <sup>3</sup> ] and PM <sub>2.5</sub> [28.5 vs. 183 µg/m <sup>3</sup> ] measured with ambient monitors at school sites. Exposure categorized by site.	SBP and DBP at 8–12 years using an automated device after 5 min rest, single-sized cuff, seated, measured 5 times. PreHTN defined as 120–139 mmHg SBP and 80–89 mmHg DBP.	Age, gender, height, weight, SES, passive smoke exposure, urinary Na, and creatinine.	An IQR increase in nano-size UFP of 20–30, 30–50, and 70–100 nm was associated with 6.35 mmHg (95% CI: 1.56, 11.47), 1.18 mmHg (95% CI: 0.05, 2.31), and 0.86 mmHg (95% CI: 0.05, 1.68) increase in SBP, respectively. An IQR increase in total UFP (1666/cm <sup>3</sup> ) associated with a 0.79 mmHg (95% CI: 0.07,
Pieters et al. 2015 <sup>37</sup>	Longitudinal HEAPS	Belgium	130	Ambient monitors of nano-sized ultrafine particles (UFP) [range: 2020–17,701 particles/cm <sup>3</sup> ], PM <sub>2.5</sub> [2–53 µg/m <sup>3</sup> ], PM <sub>10</sub> [5–64 µg/m <sup>3</sup> ], and PM <sub>10-coarse</sub> [1–34 µg/m <sup>3</sup> ] in school playgrounds. PM <sub>10-coarse</sub> defined as PM <sub>10</sub> –PM <sub>2.5</sub> fraction.	SBP and DBP measured twice, ~6 months apart, at 6–12 years of age using an automated device following 5 min of rest, while seated, with a pediatric cuff, 5–7 measures.	Sex, age, height, weight, parental education, SES, fish consumption, heart rate, school, day of week, season, wind speed, humidity, and ambient temperature day of exam.	An IQR increase in nano-size UFP of 20–30, 30–50, and 70–100 nm was associated with 6.35 mmHg (95% CI: 1.56, 11.47), 1.18 mmHg (95% CI: 0.05, 2.31), and 0.86 mmHg (95% CI: 0.05, 1.68) increase in SBP, respectively. An IQR increase in total UFP (1666/cm <sup>3</sup> ) associated with a 0.79 mmHg (95% CI: 0.07,

**Table 1** continued

Authors and Year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Confounders	Major finding(s)
Calderon et al. 2007 <sup>38</sup>	Cross-sectional	Mexico	81 Categorized by residence in NE Mexico City (n = 19), SW Mexico City (n = 40), Polotitlán (n = 22, ref.)	Ambient PM <sub>10</sub> , PM <sub>2.5</sub> , and O <sub>3</sub> measured over 1-, 2-, and 7-day periods preceding blood draw exam (not BP exam).	Systolic pulmonary arterial pressure (PAP) at 6–13 years using Doppler echocardiography in supine left position, with caffeine avoidance for 24-hrs before exam. Mean pulmonary arterial pressure (MPAP) calculated as 0.61*SPAP + 2 mmHg. <sup>65</sup>	None.	1.51) increase in SBP. No associations with DBP. No association with PM measures. Average MPAP was higher in both NE (p < 0.01) and SW (p < 0.05) Mexico City compared to Polotitlán. Among all children, increased plasma endothelin-1 levels were associated with higher MPAP (p < 0.0001). An increase in 7-day mean PM <sub>2.5</sub> was positively correlated with endothelin-1 levels (p = 0.03). No association observed with MAP. Mean carotid arterial stiffness was higher in both groups nearer to road vs. farthest (p < 0.008).
Ianuzzi et al. 2010 <sup>66</sup>	Cross-sectional	Italy	52 Categorized by distance tertiles as farthest (n = 17), middle (n = 17), or shortest (n = 18) distance from major road.	Distance to major road as a proxy from PM <sub>10</sub> , ambient PM <sub>10</sub> monitoring farthest (>1 km) from road [median: 10 µg/m <sup>3</sup> ], middle distance (300 m–1 km) [22 µg/m <sup>3</sup> ], or closest (<300 m) to the trafficked roads [40 µg/m <sup>3</sup> ]	SBP and DBP at 6–14 years of age was measured using a standard sphygmomanometer following 10 min rest, seated, with a sized cuff, three times with 1-min intervals, to the nearest 2 mmHg. Mean arterial pressure (MAP) calculated as DBP + 1/3 (SBP-DBP). Carotid arterial parameters measured using ultrasound. Stiffness was calculated as [(ln (SBP) - ln (DBP))/(Systolic diam. - Diastolic diam.)/Diastolic Diam.	Age, gender, and BMI.	

PIAMA Prevention and Incidence of Asthma and Mite Allergy, *SEMCSS* Seven Northeastern Cities Chinese Children's Study, *GINplus* German Infant Nutritional Intervention plus environmental and genetic influences on allergy development study, *LISAplus* Lifestyle-Related factors on the Immune System and the Development of Allergies in Childhood plus the influence of traffic emissions and genetics, *RANCH* Road Traffic and Aircraft Noise Exposure and Children's Cognition and Health, *MSG* monosodium glutamate, *HEAPS* Health Effects of Air Pollution in Antwerp Schools

<sup>a</sup>Represents the number of subjects included in analyses of exposure and BP if this was different from the overall study number and reported in the original article

annual averages at participants' home and school addresses estimated by land use regression) were associated with increased DBP.<sup>30</sup> No associations were observed with short-term (7-day average preceding BP measure) exposure. In a concurrent study, the composition of PM<sub>2.5</sub> and PM<sub>10</sub> (estimated as annual averages at participants' home addresses), reported that increased DBP was associated with levels of iron, silicon, and potassium in PM<sub>10</sub>, and showed marginal associations ( $p < 0.1$ ) with iron and silicon in PM<sub>2.5</sub>.<sup>29</sup> The authors attributed part of the observed effect to NO<sub>2</sub> (a marker of traffic exhaust emissions), and noted that the possible association with iron suggested non-exhaust emissions were present.

Among the 10 cross-sectional studies, 8 reported significant relationships between exposure to air pollution and children's BP. Three of these studies were based in a large cross-sectional study of 9354 children ages 5–17 years old in China, the Seven Northeastern Cities Chinese Children's Study (SNECCS) cohort.<sup>31–33</sup> The studies assessed the effects of long-term<sup>32,33</sup> or short-term exposure.<sup>31</sup> Four-year average concentrations of PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, O<sub>3</sub>, and CO were calculated from monitoring stations. One analysis reported the increased odds of childhood HTN with O<sub>3</sub> and PM<sub>10</sub>. Increased DBP was associated with O<sub>3</sub>, PM<sub>10</sub>, and CO; increased SBP was associated with O<sub>3</sub> and PM<sub>10</sub>.<sup>32</sup> These relationships were attenuated when stratified by children who were breastfed versus those who were not. Another analysis conducted in the SNECCS cohort reported the interaction effects of obesity and air pollutant exposure on BP outcomes, and found significant relationships between NO<sub>2</sub>, O<sub>3</sub>, and PM<sub>10</sub> with the odds of HTN that were consistently larger effect sizes among overweight/obese children compared to normal weight children.<sup>33</sup> The findings reported evidence of an interactive effect with child weight and exposure to PM<sub>10</sub>, SO<sub>2</sub>, and O<sub>3</sub> for both SBP and DBP. A third study assessed short-term effects using 5-day mean exposures to PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub>, and reported increased odds of elevated BP (defined as either SBP or DBP >95<sup>th</sup> percentile) for both O<sub>3</sub> and PM<sub>10</sub>.<sup>31</sup> SBP and DBP were positively associated with all four pollutants at various lag times.<sup>31</sup> When stratified by sex the relationship between elevated BP and NO<sub>2</sub> was observed only in males. Another large cross-sectional study of 2368 German children at 10 years of age exposed to air and noise pollution due to traffic<sup>34,35</sup> found no association between NO<sub>2</sub>, PM<sub>2.5</sub>, or PM<sub>10</sub> and BP overall, however, when restricted to the 605 participants with noise information a significant relationship with NO<sub>2</sub> and increased DBP was observed.<sup>34</sup>

In a secondary analysis of a subsample of 276 children age 9 and 10 years in the UK attending school near a major airport, no association was observed between NO<sub>2</sub> levels and child BP with or without adjustment for noise,<sup>27</sup> and the study replicated previous findings that aircraft and road traffic noise were not associated with child BP.<sup>36</sup> In a cross-sectional study of 184 adolescent males ages 10–14 years in Saudi Arabia, increased SBP and DBP and odds of "preHTN" were associated with attendance at schools located closest to an oil refinery, compared to a school location further from the refinery.<sup>25</sup> Concentrations of PM<sub>10</sub> and PAH metabolites were measured including total hydroxyphenanthrenes and 1-hydroxypyrene, however no relationships were identified. In a longitudinal study of 130 children in Belgium ages 6–12 years, increased total ultrafine particle fraction as well as fraction of nano-sized (20–30 nm) ultrafine particles were associated with increased SBP. No effects were identified for nano-sized fractions with diameter >100 nm (PM<sub>0.1–2.5</sub>), nor PM<sub>2.5</sub>, PM<sub>course</sub> or PM<sub>10</sub>.<sup>37</sup> In a cross-sectional study of 81 Mexican children 6–13 years of age, 7-day average PM<sub>2.5</sub> and O<sub>3</sub> were measured/estimated based on residence in Polotitlan ( $n = 22$ , control), Northeast ( $n = 19$ ) or Southwest Mexico City ( $n = 40$ ). Doppler echocardiography was used to measure systolic pulmonary arterial pressure (PAP) and diastolic PAP, and calculate mean pulmonary arterial pressure (MPAP).<sup>38</sup> MPAP was increased in NE

and SW Mexico City locations compared to Polotitlan, and the 7-day mean PM<sub>2.5</sub> was associated with MPAP. The assessment of MPAP, rather than systemic BP, provides insight into potential pulmonary effects of PM<sub>2.5</sub>, but may be less informative for developmental renal effects.

In addition to two studies discussed above,<sup>25,27</sup> three of the cross-sectional studies of indoor air quality or pollution assessed by proxy measures (e.g., distance, location) reported no association between exposure to air pollution and children's BP. A cross-sectional study of 240 Chinese study children 10 years of age, identified no association between 24-h indoor PM<sub>2.5</sub> or BC exposure,<sup>24</sup> using previously reported assessment techniques.<sup>39</sup> A cross-sectional study of 179 Pakistani children 8–12 years of age measured indoor and outdoor PM<sub>10</sub> and PM<sub>2.5</sub> at schools located in geographic regions with low ( $n = 79$ ) and high ( $n = 93$ ) air pollution.<sup>23</sup> Although direct associations were not reported for indoor or outdoor PM measures, the authors observed higher SBP and DBP in areas with high pollution compared to controls (both indoor and outdoor levels were increased in "high" pollution area). Finally, a cross-sectional analysis of 52 Italian children, which assessed distance to major road as a proxy for PM<sub>2.5</sub> exposure, reported no significant association with MAP, or BP, however the authors could not adjust for covariates.<sup>26</sup>

#### Epidemiologic evidence of association between metals and childhood BP

We identified 10 studies that examined the effects of selected nephrotoxic metals comprised of As ( $n = 2$ ), Pb ( $n = 2$ ), Cd ( $n = 4$ ), and Hg ( $n = 3$ ) on child BP (Table 2). One study examined both Cd and As and one study examined Cd and Pb.

Of the two studies examining early life As exposure, one identified a significant relationship with childhood BP, for both prenatal and postnatal As exposure.<sup>40</sup> In a prospective cohort study of 1887 mother–baby pairs in Bangladesh, an increase in pregnancy urine-As (U-As) (average measure at weeks 8 and 30 of gestation) was associated with increased SBP and DBP at 4.5 years of age. Additionally, children's U-As at 18 months was associated with an increase in SBP at 4.5 years. In a subsequent cross-sectional study of 1356 children aged 4.5 years from the same population, no association was observed with U-As levels (reported as the sum of inorganic, methylarsonic acid, and dimethylarsonic metabolites) measured approximately 8 months following BP assessment.<sup>41</sup> Although U-As typically reflects ongoing exposure, the study aimed to capture spot urine measures as a proxy for consistent As exposure through drinking water, as well as chronic Cd exposure (discussed below).

Of the two studies examining early life Pb exposure, both were conducted in prospective birth cohorts; however, only one identified a significant relationship with childhood BP.<sup>42</sup> In a prospective birth cohort of 457 mother–child pairs in Mexico City, an increase in maternal tibia Pb was associated with increased SBP and DBP in girls, but not boys. No associations were observed with patella Pb or cord blood Pb levels. In a prospective cohort of 1511 dyads in Bangladesh, no association was observed with SBP or DBP at 4.5 years with maternal blood Pb measured at 14 and 30 weeks gestation.<sup>43</sup>

Among the four studies examining early life Cd exposure, none reported a significant relationship with childhood BP.<sup>41,44–46</sup> Among these studies, two measured urine Cd (U-Cd), in either maternal<sup>41</sup> or children's samples,<sup>46</sup> and two measured blood Cd in children's samples.<sup>44,45</sup> Two studies were prospective (including a secondary analysis of a randomized control trial), and two were cross-sectional (including a case–control design measured at two time points). The prospective study of 1356 Bangladeshi children reported no association with maternal U-Cd levels in pregnancy.<sup>41</sup> In a prospective cohort (secondary analysis of randomized control trial data) of 223 control and 218 US children who received succimer treatment, blood Cd measured at 2 years of age was not

**Table 2.** Studies assessing exposure to nephrotoxic metals and children's blood pressure outcomes ( $n = 10$ )

Authors and Year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Hawkesworth et al. 2013 <sup>40</sup>	Prospective cohort MINIMat	Bangladesh	1887	Maternal urinary As (8th [median: 83 µg/L] and 30th week [83 µg/L]) and urinary Cd (week 8 [0.63 µg/L] Child U-As at 18 mo [34 µg/L]).	SBP and DBP at 4.5 years using an automated device, after 5 min rest, while seated, in triplicate with 1-min intervals.	Child's sex, age, height, SES, season of birth, maternal early pregnancy BP.	Each 1 mg/L increase in pregnancy U-As was associated with 3.69 mmHg (95% CI: 0.74, 6.63) increase in SBP and 2.91 (95% CI: 0.41, 5.42) increase in DBP. Child U-As at 18 mo was associated with 8.25 mmHg (95% CI: 1.37, 15.1) increase SBP. No associations observed with U-Cd.
Skroder et al. 2015 <sup>41</sup>	Cross-sectional MINIMat	Bangladesh	1356	Children's urine iAs [Mean: 54 µg/L, range: 16–343 µg/L] and urine Cd [Mean: 0.22 µg/L, range: 0.0078–0.63 µg/L] at ~5 years of age, collected ~8 mo after BP evaluation	SBP and DBP at 4.4–5.4 years using an automated device, after 5 min rest, while seated, in triplicate with 1-min intervals. HTN classified according to height and age 1996 report. <sup>67</sup>	Sex, birth weight, season of birth, age, weight for age z-score, maternal early pregnancy BMI, parity and SES; additionally adjusted for arsenic, cadmium and selenium. Stratified by sex. ( $p = 0.12$ ).	No observed significant associations. A moderate association was observed for each 0.5 µg/L increase in U-Cd and a 0.60 (95% CI: -0.16, 1.4) increase DBP ( $p = 0.12$ ).
Skroder et al. 2016 <sup>43</sup>	Prospective cohort MINIMat	Bangladesh	1511 or 713 (with Pb data at 14 or 30 weeks)	Maternal blood erythrocyte Pb at gestational weeks 14 [median: 73 µg/kg, range: 1.6–368] and 30 [85 µg/kg; range: 19–370]	SBP and DBP at 4.5 years using an automated device, after 5 min rest, while seated, in triplicate with 2–3 min intervals. HTN classified according to height and age 1996 report <sup>67</sup> .	Gender, birth weight, season of birth, age, height for age z-score, maternal early pregnancy BMI, parity, SES, and supplementation group. Stratified by sex.	No observations.
Zhang et al. 2012 <sup>42</sup>	Prospective cohort ELEMENT	Mexico	457	Maternal tibia Pb [median: 9.3 µg/g] and patella Pb [1.6 µg/g] measured 1-mo postpartum, and cord blood Pb [mean: 5.51 µg/dL]	SBP and DBP at 10 years using a standard sphygmomanometer after 5 min rest, on the left arm, with a sized cuff.	Maternal education, smoke during pregnancy, caloric intake, calcium and iron intake, infant birth order, gestational age, weight at birth, and child age, height, and BMI. Stratified by sex.	And IQR increase in maternal tibia Pb (13 µg/g) was associated with 2.11 mmHg (95% CI: 0.69, 3.52) increased SBP and 1.60 mmHg (95% CI: 0.28, 2.91) DBP in girls, but not boys. No associations observed with patella or cord blood Pb.
Swaddiwudhipong et al. 2015 <sup>46</sup>	Cross-sectional	Thailand	594 (301 vs. 293 Cd-exposed vs. comparison)	Urinary Cd [geometric mean: 0.57 µg/L vs. 0.39 µg/L comparison]	SBP and DBP measured twice on the right arm while seated.	Age, sex, blood Pb	No association between continuous or tertiled Cd with BP.
Cao et al. 2009 <sup>44</sup>	Prospective cohort (secondary analysis of RCT) TLC	US	441 (combined 223 placebo and 218 succimer groups)	Blood Cd at 2 years [mean: 0.21 µg/L]	SBP and DBP measured at 2, 5, and 7 years using an automated device, while seated, measured up to three times.	None reported for the relationship with BP.	No association with SBP or DBP at 2, 5, or 7 years.
Kelishadi et al. 2013 <sup>45</sup>	Case-control CASPIAN-III	Iran	320 (160 with metabolic	SBP and DBP at 10–18 years measured using a protocol	None	None	



**Table 2** continued

Authors and Year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Kalish et al. 2014 <sup>48</sup>	Prospective Cohort Project Viva	US	1103	Blood Cd [mean MetS: 10.09 µg/L; mean Control: 9.97 µg/L]	reported in ref <sup>46</sup> (National High Blood Pressure Education Program Working Group on High Blood Pressure in Children Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Pediatrics 114, 555–76 (2004). SBP and DBP at early (median 3.2 years) and mid-childhood (median 7.7 years) using an automated device with a sized cuff, arm and position noted, up to five times in 1-min intervals	Measurement conditions (position, sleep/wake state, arm), child age, sex, BMI z-score, fetal growth z-score, maternal age, race/ethnicity, education, marital status, pre-pregnancy BMI, smoking status, and second trimester BP second trimester fish consumption	No significant association with DBP or SBP.  No association with SBP observed at age 3 or 7 years, comparing Q4 to Q1.
Thurston et al. 2007 <sup>47</sup>	Prospective cohort SCDS	Seychelles	644 and 559 (age 12 and 15 years)	Total Hg assessed in maternal hair during pregnancy [mean: ~6.6 ppm for male offspring, and 7.0 for girls]	SBP and DBP at 12 and 15 years using an automated device, with a sized cuff following a few min of rest, while seated, measured in duplicate.	Maternal HTN during pregnancy, child sex, birth weight, age, BMI, height.	A 1-ppm increase in prenatal meHg was associated with a 0.17 mmHg increase in DBP at age 15 years among boys ( $p < 0.02$ ), but not girls. No associations observed at age 12 years.
Valera et al. 2012 <sup>49</sup>	Prospective cohort	Canada, Nunavik Inuit	226	Cord blood Hg [median: 81.5 nmol/L], blood at 11 years [14.5 nmol/L], and hair at 11 years [4.75 nmol/g]	SBP and DBP at 11 years using a mercury sphygmomanometer after 5 min rest, while seated, measured in triplicate (averaged last two measures).	Age, sex, birth weight, BMI, height, total cord n-3 fatty acids, PBC 153, cord blood Pb, selenium and maternal smoking in pregnancy	No association observed with SBP or DBP for either blood Hg assessment time point or hair levels at age 11 years.

*MINIMat* Maternal and Infant Nutrition Interventions, Matlab, *iAs* inorganic arsenic, *ELEMENT* Early Life Exposures in Mexico to Environmental Toxicants, *TLC* Treatment of Lead-Exposed Children, *CASPIAN-III* Childhood and adolescence surveillance and prevention of adult non-communicable disease, *SCDS* Seychelles Child Development Study

<sup>a</sup>Represents the number of subjects included in analyses of exposure and BP if this was different from the overall study number and reported in the original article

associated with BP at 2, 5, or 7 years.<sup>44</sup> A decrease in SBP was observed 1-week following succimer treatment, although the authors note the fall in BP may be attributable to reduction in Pb burden and there was no clear association with Cd and BP. In a cross-sectional study of 594 Thai school-age children randomly selected from 10 Cd-contaminated schools and 3 outside the contamination area, no significant association was found for U-Cd when analyzed as a continuous variable or categorized by tertile with SBP or DBP.<sup>46</sup> Blood Pb was measured in this study but the relationship with BP was not reported. Lastly, in a case-control study of 320 children from Iran (160 normal weight, 160 cases with metabolic syndrome), blood Cd measured at 10 and 18 years showed a positive, but not significant association with DBP. The analysis did not include adjustment for covariates, and did not report the method of BP measurement. Notably, the Cd levels were relatively high among the control group (mean was 10.09 µg/L for cases and 9.97 µg/L for controls), and the analysis included linear regression across all cases and controls and reported null associations for the relationship between blood Cd and DBP or SBP.

We identified three studies examining early life Hg exposure, and only one reported an association with childhood BP. All three studies were prospective cohorts. A prospective cohort study of 645 and 561 children in the Seychelles that assessed maternal hair Hg and followed children to 12 and 15 years, respectively, reported that prenatal maternal Hg levels were associated with increased DBP among 15-year-old boys but not girls.<sup>47</sup> No associations were observed among 12-year-old children. A large prospective cohort study of 1103 mother-child pairs in the United States, found no association with maternal blood erythrocyte Hg measured during the second trimester and SBP at 3.2 or 7.7 years.<sup>48</sup> Another prospective cohort study of 226 Nunavik Inuit children, measured cord blood Hg, as well as blood and hair Hg at 11 years of age, and reported no association with Hg exposure at any time point and BP at 11 years.<sup>49</sup> Notably, the studies above reported lower Hg levels than a 1999 study (outside the range of years reviewed herein) conducted of 917 children in the Faroe Islands.<sup>50</sup> The study identified a significant relationship between cord blood Hg and SBP at 7 years, and we note that Hg levels were 2–3× higher in cord or maternal hair levels than those reported in refs.<sup>49,47</sup> which may account for the lack of statistically significant findings in the later studies. Additionally, the selection of appropriate biomarker matrices for nephrotoxicity may have affected the results (i.e., hair vs. urine).

#### Epidemiologic evidence of association between phthalates and other organic compounds and childhood BP

We identified 12 studies that examined the effects of other potential environmental nephrotoxics on child BP, comprised of phthalates ( $n = 4$ ), Bisphenol A (PBA) ( $n = 3$ ), polychlorinated biphenols (PCBs) ( $n = 2$ ), pesticides ( $n = 4$ , two also measured PCBs) and flame retardants ( $n = 1$ ) (Table 3). The studies were either cross-sectional ( $n = 6$ ) or prospective ( $n = 6$ ) in design. Studies of phthalates, pesticides, and BPA used urinary levels of the parent compound or metabolites, whereas PCBs and flame retardants were measured in blood.

Of the four studies examining early life phthalate exposure, all reported a significant relationship with subsequent childhood BP, for both prenatal and postnatal exposure. In a prospective study of 379 mother-child pairs in Spain, maternal phthalate exposure (averaged from first and third trimester urine measures, grouped by high or low molecular weight), showed significantly lower SBP z-score among girls comparing the 2nd and 3rd tertiles to the first tertile of summed molar concentrations of high molecular weight phthalates (HMWP) and comparing only the 3rd tertile to first for summed molar concentrations of low molecular weight phthalates (LMWP).<sup>51</sup> In another prospective cohort study of 258 children in the United States, prenatal levels of mono(2-ethyl-5-oxohexyl)

phthalate (MEOHP) and mono(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP) measured in maternal urine (at 12 and 20 weeks gestation) were associated with DBP at 14 years of age, but not at 5 or 9 years of age.<sup>52</sup> This study did not directly report estimates of the association between phthalates and SBP or DBP. Associations were also noted in two large cross-sectional studies in the United States.<sup>53,54</sup> Among 2447 adolescents participating in NHANES 2003–2008, grouped urine phthalates and metabolite molar sums (HMWP, LMWP, and diethylhexyl phthalate [DEHP]) were assessed in relation to SBP z-score and “preHTN”.<sup>54</sup> An increase in DEHP metabolites was associated with a significantly higher SBP z-score. Individual metabolites including MEHP, MBP, MEHHP, and MEOHP were associated with increased SBP z-score and MEP was associated with increased odds of preHTN. No association was observed with the LMWP or HMWP groups. In a subsequent study of 1329 adolescents participating in NHANES 2009–2012, molar sums of HMWP and urinary DEHP metabolites as well as the parent compound for its replacements, di-isononyl phthalate (DINP) and di-isodecyl phthalate (DIDP), were each associated with increased SBP z-score.<sup>53</sup> No associations were observed with the LMWP group. An increase in summed DEHP metabolites was also associated with increased DBP z-score. This study further identified the significant metabolites within each group and we refer the reader to Table 3 and the original article for details.

Among three studies examining early life BPA exposure, two were conducted in prospective cohorts and one was a cross-sectional study. Two reported significant associations with childhood BP. In a prospective cohort of 486 mother-child pairs in Korea, higher prenatal BPA exposure (measured at 20 weeks gestation) was associated with a significant increase in DBP at levels above an identified threshold of 4.5 µg BPA per g creatinine.<sup>55</sup> Additionally, for prenatal BPA levels above 5.0 µg/g creatinine, a relationship with decreased pulse pressure was observed. Among girls, a U-shaped relationship with SBP was identified above and below the level of 0.7 µg BPA/g creatinine. However, in a prospective cohort study of 500 mother-child pairs in Greece, urine BPA (measured during the first trimester, as well as at 2.5 and 4 years) was not associated with child BP at 4 years.<sup>56</sup> Lastly, a limited cross-sectional study of 39 obese or overweight children ages 3–8 years in the United States, urinary BPA adjusted for creatinine was associated with elevated DBP among males, but not girls.<sup>57</sup> No associations were observed with SBP.

Among two studies examining early life PCB exposure (a persistent organic pollutant banned in the 1970s), one reported evidence of association with BP. In a prospective cohort study of 427 Greek children, the persistent organic pollutant hexachlorobenzene (HCB) was associated with higher SBP, and p,p'-DDE, a metabolite of DDT, was associated with higher DBP at 4 years of age.<sup>58</sup> No association was observed for a group of summed PCBs comprised of six congeners. In a cross-sectional study of 158 Korean children aged 7–9 years at baseline, the sum of 32 measured PCBs and 2 individual PCBs (congeners 138 and 153) were associated with significantly higher DBP at 1-year follow-up, after adjustment for sex, age, household income, and BMI.<sup>59</sup> No associations were identified with SBP or DBP and 19 organochlorines measured.

Among four studies examining early life organochlorine compound exposure, three reported significant associations with BP. The prospective cohort study by Lee et al. (described above) reported no association with organochlorine pesticides,<sup>59</sup> whereas the study by Vefaidi et al. (described above),<sup>58</sup> and two cross-sectional studies from the ESPINA cohort in Ecuador reported associations.<sup>60,61</sup> In a cross-sectional examination of 87 children, prenatal pesticide exposure assessed by parental occupation interview were marginally associated with higher SBP at 6–8 years among a subset of 69<sup>60</sup>; no relationships were observed with urine organophosphate metabolites. In a follow-up study of 271 children, duration of living with a flower worker as a

**Table 3.** Studies assessing exposure to phthalates ( $n = 4$ ), bisphenol A ( $n = 3$ ), polychlorinated biphenols ( $n = 2$ ), pesticides ( $n = 3$ ), one also measured PCBs), and flame retardants/perfluoroalkyl chemicals ( $n = 1$ ) and children's blood pressure outcomes ( $n = 12$ )

Authors and year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Valvi et al. 2015 <sup>51</sup>	Prospective cohort INMA	Spain	379	Maternal urine MBzP, MEHP, MEHHP, MEOHP, MECPP, MEP, MIBP, and MEP in the first and third trimesters of pregnancy; creatinine-adjusted and converted to molar sums for: $\Sigma$ DEHPm [median: 95.1 $\mu$ g/g creatinine], $\Sigma$ HMWp [112 $\mu$ g/g creatinine], and $\Sigma$ LMWp [472 $\mu$ g/g creatinine]	SBP and DBP at age 4 and 7 years using an automated device after 5 min rest, with a sized cuff, on the right arm, age- and height-specific z-scores, stratified by sex. High BP defined as any BP z-score $\geq$ 90th percentile.	Child sex and age, maternal country of origin, age at delivery, parity, education, social class, pre-pregnancy BMI, and smoking in pregnancy.	Across all ages, $\Sigma$ HMWp were associated with lower SBP z-scores in girls (adjusted $\beta = -0.39$ ; 95% CI: $-0.65$ , $-0.12$ for the 2nd tertile and $-0.28$ ; $-0.55$ , $-0.01$ for the 3rd tertile of exposure), but not boys. Associations with $\Sigma$ HMWp were driven by DEHP. $\Sigma$ LMWp were associated with lower SBP z-scores in girls (adjusted $\beta = -0.23$ ; 95% CI: $-0.50$ , $-0.04$ in 2nd tertile and $-0.40$ ; $-0.66$ , $-0.12$ in 3rd tertile of exposure), but not boys. Associations with $\Sigma$ LMWp were driven by MEP. No association observed with DBP z-scores.
Tran et al. 2017 <sup>52</sup>	Prospective cohort CHAMACOS	US	258	Maternal urine assessed for 11 phthalate metabolites at $\sim$ 13 and 26 weeks' gestation (LMW: MEP, MBP, MIBP; $\Sigma$ DEHP: MEHP, MEHHP, MEOHP, MECPP; HMW: MBzP, MCPP, MCOP, MCNP) and 8-isoprostane creatinine-adjusted.	SBP and DBP at age 5, 9, and 14 years using an automated device after rest.	BMI, sex, and creatinine	Prenatal MEOHP and MEHHP were associated with DBP at 14 years (betas not reported). Cross-sectionally, SBP and DBP were positively associated with 8-isoprostane at 14 years (SBP beta: 0.008 95% CI: 0.002, 0.015; DBP beta: 0.011 95% CI: 0.001, 0.02). Isoprostane at 14 years was positively associated with MEHP, MECPP, and $\Sigma$ DEHP at 12 weeks of gestation. The effect of 13-week MEOHP and MEHHP on 14-year DBP mediated by 14-year 8-isoprostane was not significant.
Trasande et al. 2013 <sup>54</sup>	Cross-sectional NHANES (2003–2008)	US	2447	Spot urine levels converted to molar sums of $\Sigma$ LMWP (comprising MEP, MBP, MIBP), $\Sigma$ HMWp (comprising MECPP, MEHHP, MEOHP, MEHP, MBzP), and $\Sigma$ DEHP (comprising MEHP, MECPP, MEHHP, MEOHP).	SBP and DBP 8–19 years using a standard sphygmomanometer, following 5 min rest, in triplicate. Age and height z-scores calculated according to 4th report [66]. PreHTN defined as $\geq$ 90th percentile age and height z-score stratified by sex.	Urine creatinine, BMI, race/ethnicity, age, caregiver education, poverty-income ratio, sex, serum cotinine, caloric intake, and TV watching	Each log-unit ( $\sim$ 3-fold) increase in $\Sigma$ DEHP was associated with a 0.041 SD unit increase in SBP z-score ( $p = 0.047$ ). $\Sigma$ DEHP relationships with SBP z-score driven by MEHP, MBP, MEHHP, and MEOHP (all $p < 0.05$ ). MEP associated with

**Table 3** continued

Authors and year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Trasande et al. 2015 <sup>53</sup>	Cross-sectional NHANES (2009-2012)	US	1329	Urine phthalates converted to molar sums: $\Sigma$ LMW, $\Sigma$ HMW, $\Sigma$ DEHP, ( $\Sigma$ DINP), and ( $\Sigma$ DIDP).	SBP and DBP at 8-19 years using a standard sphygmomanometer, following 5 min rest, in triplicate. Age- gender- and height-standardized z-scores calculated according to 4th report <sup>66</sup> . PreHTN defined as $\geq$ 90th percentile age and height z-score stratified by sex.	Urine creatinine, BMI, race/ethnicity, age, poverty-income ratio, gender, serum cotinine, caloric intake, and physical activity.	increased odds of preHTN (OR: 1.20, $p = 0.04$ ). No association observed with $\Sigma$ LMWP or $\Sigma$ HMW. A 1-log unit increase in $\Sigma$ HMW, $\Sigma$ DINP, $\Sigma$ DIDP, and $\Sigma$ DEHP associated with $\sim$ 0.10 SD increase in SBP z-score ( $p < 0.05$ ). No association observed with $\Sigma$ LMWP. Associations with SBP z-score driven by MCOP, MNP, MCNP, MEHP, MECPP ( $p < 0.05$ ). A 1-log unit increase in $\Sigma$ DEHP associated with a 0.09 SD increase in DBP z-score ( $p = 0.04$ ). Associations with DBP z-score driven by MEHP, MEOHP ( $p < 0.05$ ).
Bae et al. 2017 <sup>55</sup>	Prospective cohort EDC	Korea	486	BPA after 8 h fasting in maternal urine at 20 weeks gestation [geometric mean: 1.2 $\mu$ g/g creatinine] and child urine at age 4 years [geometric mean: 3.3 $\mu$ g/g creatinine].	DBP and SBP at 4 years using an automated device following 10 min rest, while seated, on the same arm, with a pediatric cuff, measured twice in 5 min interval. PP calculated as SBP-DBP.	Age, sex, height, weight, gestational age, birth weight, parental history of HTN, paternal education, exposure to environmental tobacco, pulmonary/enteric infection, physical activity (min/week). Stratified by sex.	A 1-log unit of prenatal BPA was associated with 7.9 mmHg increase in DBP ( $p = 0.0001$ ), at BPA levels above an identified threshold level of 4.5 $\mu$ g/g creatinine ( $n = 41$ ). A 1-log unit of prenatal BPA was associated with 8.0 mmHg decrease in PP ( $p = 0.0015$ ), at BPA levels above a threshold level of 5.0 $\mu$ g/g creatinine ( $n = 31$ ). Among boys, similar relationships were observed. Among girls, significant U-shaped relationships were observed for SBP above and below a 0.7 $\mu$ g/g creatinine threshold ( $p < 0.05$ ). A negative relationship below a 0.9 $\mu$ g/g creatinine threshold with DBP ( $p = 0.03$ ).
Vafeiadi et al. 2016 <sup>56</sup>	Prospective cohort rhea	Greece	500 (4-year cross section) 235 (prenatal and 2.5 years)	BPA in maternal urine in first of pregnancy trimester [geometric mean: 1.2 $\mu$ g/g creatinine, $n = 235$ ], and child urine at 2.5 [geometric mean: 5.1 $\mu$ g/g creatinine $n = 235$ ] and 4 years [geometric mean: 1.9 $\mu$ g/g creatinine $n = 500$ ]	SBP and DBP at 4 years using an automated device, with a sized cuff, following 5 min rest, while seated, measured five times in 1-min intervals.	Maternal education, age, pre-pregnancy BMI, employment status during pregnancy, child sex, z-score birth weight for gestational age, breastfeeding status, height z-score, time spent watching	No significant associations observed with any exposure time point.

**Table 3** continued

Authors and year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Khalil et al. 2014 <sup>57</sup>	Cross-sectional	US	39 obese/overweight children	Urine BPA [median: 1.82 µg/g creatinine]	SBP and DBP at 3–8 years using an automated device with two cuff sizes.	TV, and energy intake at 4 years. Age, ethnicity	Among males ( <i>n</i> = 17), a 1-unit increase in log BPA was associated with 4.01 mmHg increase in DBP ( <i>p</i> = 0.01). Associations with SBP not reported.
Vafeiadi et al. 2016 <sup>58</sup>	Prospective cohort Rhea	Greece	427	POPs (PCBs, DDE [median: 1981.2 pg/mL], and HCB [median: 82.5 pg/mL]) in first trimester maternal serum ΣPCB [median: 319.4 pg/mL] comprising congeners 118, 138, 153, 156, 170, and 180.	SBP and DBP at 4 years using an automated device with a sized cuff, after 5 min rest, while seated, measured 5 times in 1-min intervals.	Maternal serum triglycerides and cholesterol, age, pre-pregnancy BMI, parity, education, smoking status during pregnancy, breastfeeding duration, child sex, age, birth weight, and gestational age.	A 10-fold increase in HCB was associated with a 4.34 mmHg (95% CI: 0.63, 8.05) increase in SBP. A 10-fold increase in DDE was associated with higher DBP (1.79 mmHg 95% CI: 0.13, 3.46) ΣPCB was not associated with SBP or DBP.
Lee et al. 2016 <sup>59</sup>	Prospective Ewha Birth and Growth Cohort	Korea	158	POPs [32 PCBs and 19 OCPs] in blood 7 PCBs (52, 101, 118, 138, 153, 156, and 180) and 5 OCPs (HCB, <i>trans</i> -Nonachlor, β-HCH, <i>p,p'</i> -DDT and <i>p,p'</i> -DDE) were considered "detectable" and analyzed individually. "Marker PCBs" category comprised of 6 congeners (28,52,101,138,153,180) and ΣPCB was comprising all 32 congeners.	BP at 7–9 years and after 1-year follow-up using an automated device with a sized cuff, after rest, in a "stable" position, measured twice in 5-min intervals. MAP was calculated as [(SBP-DBP)/3] + DBP.	Sex, age, household income, and change in BMI	PCB 138 and 153 were associated with an increase in relative change in 1-year DBP (3.01: 95% CI 0.20–5.82 and 2.33: 95% CI: 0.21, 4.45). ΣPCB and "marker PCB" were associated with an increase in relative change in 1-year DBP (4.84: 95% CI: 1.11, 8.57, and 4.44 95% CI: 0.85, 8.03). No association with organochlorine pesticides associated with DBP. No associations observed with SBP.
Suarez-Lopez et al. 2013 <sup>61</sup>	Cross-sectional ESPINA	Ecuador	271 A subset of 138 reported cohabitation with a flower plantation worker.	Secondary pesticide exposure defined as living with a flower worker [mean no. of workers at home: 1.64], duration of cohabitation with a flower worker [mean: 5.3 years], and number of "bad practices" likely to result in home contamination with pesticides [mean: 2.9]. Erythrocyte AChE activity was measured by finger prick as a proxy for exposure to pesticides that inhibit cholinesterase activity [mean: 3.14 U/mL].	SBP and DBP at 4–9 years with a pediatric aneroid sphygmomanometer, following 3–5 min rest, while seated, both feet on floor, measured twice). BP z-scores for age, sex and height for age for calculated according to the 4th Report <sup>66</sup>	Age, sex, race, height-for-age z-score, exam date, heart rate, number of smokers in home, residence to nearest flower plantation, pesticide use in household, and pesticide use by neighbors.	Each year longer duration of cohabitation with a flower worker and increase in number of bad practices were associated with a 0.32 mmHg (95% CI: –0.64, –0.02) and 0.82 mmHg (95% CI: –1.45, –0.20) decrease in SBP. Living with flower worker was associated with a 1.72 mmHg decrease in SBP (95% CI: –3.53, 0.08; <i>p</i> = 0.06). A 1 U/mL decrease in AChE activity was associated with a 2.86 mmHg decrease in SBP (95% CI: –5.20, –0.53) and a 2.89 mmHg decrease in DBP

**Table 3** continued

Authors and year	Study design, name	Location	Sample size <sup>a</sup>	Exposure measure [reported level]	BP evaluation	Covariates	Major finding
Harari et al. 2010 <sup>60</sup>	Cross-sectional	Ecuador	87	Prenatal exposure assessed by interview of maternal/paternal occupational exposure. Children's spot urine assessed for 6 organophosphate metabolites, individually and summed as $\Sigma$ dimethyl, $\Sigma$ diethyl, or $\Sigma$ all molar concentrations. Erythrocyte AChE activity measured by finger prick.	SBP and DBP at 6–8 years using a standard sphygmomanometer, while seated with a child-sized cuff, measured twice.	Age, sex, race, BMI, stunting, and cohort	(−5.00, −0.78). Similar results observed with BP z-scores. Prenatal maternal occupational pesticide exposure was associated with a 3.6 mmHg (95% CI: −0.1, 7.2) increase in SBP ( $p \leq 0.05$ ) compared to those without maternal exposure. No associations observed with paternal occupational exposure or cross-sectional children's exposure.
Geiger et al. 2014 <sup>62</sup>	Cross-sectional NHANES (1999–2000, 2003–2008)	US	1655	PFOS and PFOA in serum	SBP and DBP at 12–18 years using a standard sphygmomanometer, following 5 min rest while seated, measured three times. HTN defined as BP $\geq$ 95th percentile adjusted for age, height, and sex according to 1996 report <sup>7</sup> .	Age, sex, race/ethnicity, BMI, household income, physical activity, total cholesterol, and serum cotinine.	No association was observed for SBP or DBP.

<sup>a</sup>MMA Infancia y Medio Ambiente (Environment and Childhood), CHAMACOS Center for Health Assessment of Mothers and Children of Salinas, EDC Environment and Development of Children, ESPINA Estudio de la Exposición Secundaria a Plaguicidas en Infantes, Niños y Adolescentes (Secondary Exposure to Pesticides among Infants, Children and Adolescents Study), DIMP di-isononyl phthalate, DIDP di-isodecyl phthalate, DDE dichlorodiphenyl-dichloroethylene, HCB hexachlorobenzene, NHANES National Health and Nutrition Examination Survey, PFOA perfluorooctanoic acid, PFOS perfluorooctane sulfonate

<sup>7</sup>Represents the number of subjects included in analyses of exposure and BP if this was different from the overall study number and reported in the original article

proxy for exposure as well as number of “bad practices” likely to result in home contamination were significantly associated with lower SBP.<sup>61</sup>

Finally, we identified a single study examining early life exposure to perfluoroalkyl flame retardants with childhood BP. A cross-sectional analysis of 1655 children participating in NHANES in years 1999–2000 and 2003–2008, reported no association with serum PFOA or PFOS and continuous BP measure or HTN.<sup>62</sup>

## DISCUSSION

Summary and recommendations for future directions

There is a growing body of literature supporting adverse cardiorenal effects of air pollution exposure. For metals, the literature is sparse in children, but suggests that As, Hg, and Pb may affect childhood BP. Among the persistent organic pollutants there is limited but consistent evidence that phthalates and pesticide exposure may alter childhood BP. There is mixed evidence for BPA and PCB effects, and there is no current evidence to support an association between childhood PFC exposure and BP. There remain many gaps in our knowledge, including the importance of developmental susceptibility windows (i.e., the role of exposure timing in predicting effects), appropriate study design such as prospective studies that can avoid the issue of reverse causality (in which renal damage alters biomarker levels—a potential issue in cross-sectional studies), more standardized measurement and definition of BP outcomes, need for more complex statistical modeling approaches that factor in developmental trajectories or mixed exposure, and consideration of genetic susceptibility and epigenetic mediation as a way to define susceptible subpopulations and mechanisms of action. Future studies should also consider possible effects of emerging contaminants given that over 100,000 chemicals are registered for industrial use in the United States yet only a handful have been studied with regards to renal toxicity. Additional attention is also needed to examine potential maternal BP effects from exposures in pregnancy. Incorporating these considerations into new studies will enable the research community to address existing limitations and enhance our understanding of early life environmental exposures on BP. Given the heterogeneity of reported exposures and outcomes in the peer-reviewed studies summarized herein, we also recommend that exposure-specific systematic reviews and meta-analyses are needed in the advancing field of developmental exposures and childhood BP.

## CONCLUSIONS

This review of the recent literature suggests that early life exposure to air pollutants, metals, and organic pollutants may contribute to changes in childhood BP. Limited evidence suggests that prenatal exposure may be more deleterious to BP than postnatal exposures and requires additional investigation. Because childhood BP tracks with adult BP and HTN<sup>63,64</sup>, it is imperative that hypertensive children are identified early for intervention and potential environmental, behavioral, and biological risk factors.

## ACKNOWLEDGEMENTS

This work was supported in part by funding from the Children’s Center Foundation and NIH: K99ES027508, P30ES23515, and P30DK079307 (The Pittsburgh Center for Kidney Research), U2CE5026561 R01ES013744, R01ES14930, R01ES020268, and R01ES021357. The illustrations in Fig. 1 were created by Jill Gregory, Department of Academic Medical Illustration, Icahn School of Medicine at Mount Sinai.

## ADDITIONAL INFORMATION

**Competing interests:** The authors declare no competing interests.

**Publisher’s note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## REFERENCES

1. Yoon, S. S., Carroll, M. D., Fryar, C. D. Hypertension prevalence and control among adults: United States, 2011–2014. *NCHS Data Brief* **220**, 1–8 (2015).
2. Flynn, J. T. et al. Clinical practice guideline for screening and management of high blood pressure in children and adolescents. *Pediatrics* **140**, e20171904 (2017).
3. Rao, G. Diagnosis, epidemiology, and management of hypertension in children. *Pediatrics* **138**, e20153616 (2016).
4. Rosner, B., Cook, N. R., Daniels, S. & Falkner, B. Childhood blood pressure trends and risk factors for high blood pressure: the NHANES experience 1988–2008. *Hypertension* **62**, 247–254 (2013).
5. Lee, M. H. et al. A 24-year follow-up study of blood pressure tracking from childhood to adulthood in Korea: the Kangwha Study. *Yonsei Med. J.* **55**, 360–366 (2014).
6. Dorey, E. S., Pantaleon, M., Weir, K. A. & Moritz, K. M. Adverse prenatal environment and kidney development: Implications for programming of adult disease. *Reproduction* **147**, R189–198 (2014).
7. Lauer, R. M. & Clarke, W. R. Childhood risk factors for high adult blood pressure: the Muscatine Study. *Pediatrics* **84**, 633–641 (1989).
8. Gidding, S. S. et al. Higher self-reported physical activity is associated with lower systolic blood pressure: the Dietary Intervention Study in Childhood (DISC). *Pediatrics* **118**, 2388–2393 (2006).
9. Carnethon, M. R. et al. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *JAMA* **290**, 3092–3100 (2003).
10. Yang, Q. et al. Sodium intake and blood pressure among US children and adolescents. *Pediatrics* **130**, 611–619 (2012).
11. Shiue, I. & Hristova, K. Higher urinary heavy metal, phthalate and arsenic concentrations accounted for 3–19% of the population attributable risk for high blood pressure: US NHANES, 2009–2012. *Hypertens. Res.* **37**, 1075–1081 (2014).
12. Luyckx, V. A. & Brenner, B. M. Birth weight, malnutrition and kidney-associated outcomes—a global concern. *Nat. Rev. Nephrol.* **11**, 135–149 (2015).
13. Benz, K. & Amann, K. Maternal nutrition, low nephron number and arterial hypertension in later life. *Biochim. Biophys. Acta* **1802**, 1309–1317 (2010).
14. Solhaug, M. J., Bolger, P. M. & Jose, P. A. The developing kidney and environmental toxins. *Pediatrics* **113**, 1084–1091 (2004).
15. Rubin, M. I. et al. Maturation of renal function in childhood: clearance studies. *J. Clin. Invest.* **28**, 1144–1162 (1949).
16. Baum, M. Role of the kidney in the prenatal and early postnatal programming of hypertension. *Am. J. Physiol. Ren. Physiol.* **298**, F235–247 (2010).
17. Yun, M. et al. Tobacco smoking strengthens the association of elevated blood pressure with arterial stiffness: the Bogalusa Heart Study. *J. Hypertens.* **33**, 266–274 (2015).
18. Juonala, M., Magnussen, C. G. & Raitakari, O. T. Parental smoking produces long-term damage to vascular function in their children. *Curr. Opin. Cardiol.* **28**, 569–574 (2013).
19. Banderali, G. et al. Short and long term health effects of parental tobacco smoking during pregnancy and lactation: a descriptive review. *J. Transl. Med.* **13**, 327 (2015).
20. Zheng, L. Y., Sanders, A. P., Saland, J. M., Wright, R. O. & Arora, M. Environmental exposures and pediatric kidney function and disease: a systematic review. *Environ. Res.* **158**, 625–648 (2017).
21. Liang, R. et al. Effect of exposure to PM2.5 on blood pressure: a systematic review and meta-analysis. *J. Hypertens.* **32**, 2130–2140 (2014).
22. Giorgini, P. et al. Air pollution exposure and blood pressure: an updated review of the literature. *Curr. Pharm. Des.* **22**, 28–51 (2016).
23. Sughis, M., Nawrot, T. S., Ihsan-ul-Haque, S., Amjad, A. & Nemery, B. Blood pressure and particulate air pollution in schoolchildren of Lahore, Pakistan. *BMC Public Health* **12**, 378 (2012).
24. Baumgartner, J. et al. Household air pollution and children’s blood pressure. *Epidemiology* **23**, 641–642 (2012).
25. Trasande, L. et al. Polycyclic aromatic hydrocarbons, brachial artery distensibility and blood pressure among children residing near an oil refinery. *Environ. Res.* **136**, 133–140 (2015).
26. Iannuzzi, A. et al. Air pollution and carotid arterial stiffness in children. *Cardiol. Young.* **20**, 186–190 (2010).
27. Clark, C. et al. Does traffic-related air pollution explain associations of aircraft and road traffic noise exposure on children’s health and cognition? A secondary analysis of the United Kingdom sample from the RANCH project. *Am. J. Epidemiol.* **176**, 327–337 (2012).
28. van Rossem, L. et al. Prenatal air pollution exposure and newborn blood pressure. *Environ. Health Perspect.* **123**, 353–359 (2015).

29. Bilenko, N. et al. Associations between particulate matter composition and childhood blood pressure—the PIAMA study. *Environ. Int.* **84**, 1–6 (2015).
30. Bilenko, N. et al. Traffic-related air pollution and noise and children's blood pressure: results from the PIAMA birth cohort study. *Eur. J. Prev. Cardiol.* **22**, 4–12 (2015).
31. Zeng, X. W. et al. Positive association between short-term ambient air pollution exposure and children blood pressure in China—Result from the Seven Northeast Cities (SNEC) study. *Environ. Pollut.* **224**, 698–705 (2017).
32. Dong, G. H. et al. Air pollution associated hypertension and increased blood pressure may be reduced by breastfeeding in Chinese children: the Seven Northeastern Cities Chinese Children's Study. *Int. J. Cardiol.* **176**, 956–961 (2014).
33. Dong, G. H. et al. Interactions between air pollution and obesity on blood pressure and hypertension in Chinese children. *Epidemiology* **26**, 740–747 (2015).
34. Liu, C. et al. The associations between traffic-related air pollution and noise with blood pressure in children: results from the GINIplus and LISAPlus studies. *Int. J. Hyg. Environ. Health* **217**, 499–505 (2014).
35. Markevych, I. et al. A cross-sectional analysis of the effects of residential greenness on blood pressure in 10-year old children: results from the GINIplus and LISAPlus studies. *BMC Public Health* **14**, 477 (2014).
36. van Kempen, E. et al. Noise exposure and children's blood pressure and heart rate: the RANCH project. *Occup. Environ. Med.* **63**, 632–639 (2006).
37. Pieters, N. et al. Blood pressure and same-day exposure to air pollution at school: associations with nano-sized to coarse PM in children. *Environ. Health Perspect.* **123**, 737–742 (2015).
38. Calderon-Garciduenas, L. et al. Elevated plasma endothelin-1 and pulmonary arterial pressure in children exposed to air pollution. *Environ. Health Perspect.* **115**, 1248–1253 (2007).
39. Baumgartner, J. et al. Patterns and predictors of personal exposure to indoor air pollution from biomass combustion among women and children in rural China. *Indoor Air* **21**, 479–488 (2011).
40. Hawkesworth, S. et al. Early exposure to toxic metals has a limited effect on blood pressure or kidney function in later childhood, rural Bangladesh. *Int. J. Epidemiol.* **42**, 176–185 (2013).
41. Skroder, H. et al. Kidney function and blood pressure in preschool-aged children exposed to cadmium and arsenic—potential alleviation by selenium. *Environ. Res.* **140**, 205–213 (2015).
42. Zhang, A. et al. Association between prenatal lead exposure and blood pressure in children. *Environ. Health Perspect.* **120**, 445–450 (2012).
43. Skroder, H. et al. Prenatal lead exposure and childhood blood pressure and kidney function. *Environ. Res.* **151**, 628–634 (2016).
44. Cao, Y. et al. Postnatal cadmium exposure, neurodevelopment, and blood pressure in children at 2, 5, and 7 years of age. *Environ. Health Perspect.* **117**, 1580–1586 (2009).
45. Kelishadi, R. et al. Association of blood cadmium level with cardiometabolic risk factors and liver enzymes in a nationally representative sample of adolescents: the CASPIAN-III study. *J. Environ. Public Health* **2013**, 142856 (2013).
46. Swaddiwudhipong, W. et al. Renal and blood pressure effects from environmental cadmium exposure in Thai children. *Environ. Res.* **136**, 82–87 (2015).
47. Thurston, S. W. et al. Does prenatal methylmercury exposure from fish consumption affect blood pressure in childhood? *Neurotoxicology* **28**, 924–930 (2007).
48. Kalish, B. T. et al. Associations of prenatal maternal blood mercury concentrations with early and mid-childhood blood pressure: A prospective study. *Environ. Res.* **133**, 327–333 (2014).
49. Valera, B. et al. Cardiac autonomic activity and blood pressure among Inuit children exposed to mercury. *Neurotoxicology* **33**, 1067–1074 (2012).
50. Sorensen, N., Murata, K., Budtz-Jorgensen, E., Weihe, P. & Grandjean, P. Prenatal methylmercury exposure as a cardiovascular risk factor at seven years of age. *Epidemiology* **10**, 370–375 (1999).
51. Valvi, D. et al. Prenatal phthalate exposure and childhood growth and blood pressure: evidence from the Spanish INMA-Sabadell Birth Cohort Study. *Environ. Health Perspect.* **123**, 1022–1029 (2015).
52. Tran, V. et al. Prenatal phthalate exposure and 8-isoprostane among Mexican-American children with high prevalence of obesity. *J. Dev. Orig. Health Dis.* **8**, 196–205 (2017).
53. Trasande, L. & Attina, T. M. Association of exposure to di-2-ethylhexylphthalate replacements with increased blood pressure in children and adolescents. *Hypertension* **66**, 301–308 (2015).
54. Trasande, L. et al. Urinary phthalates are associated with higher blood pressure in childhood. *J. Pediatr.* **163**, 747–753 (2013). e741.
55. Bae, S. et al. Maternal urinary bisphenol A concentration during midterm pregnancy and children's blood pressure at age 4. *Hypertension* **69**, 367–374 (2017).
56. Vafeiadi, M. et al. Association of early life exposure to bisphenol A with obesity and cardiometabolic traits in childhood. *Environ. Res.* **146**, 379–387 (2016).
57. Khalil, N. et al. Bisphenol A and cardiometabolic risk factors in obese children. *Sci. Total Environ.* **470–471**, 726–732 (2014).
58. Vafeiadi, M. et al. Association of prenatal exposure to persistent organic pollutants with obesity and cardiometabolic traits in early childhood: the Rhea Mother-Child Cohort (Crete, Greece). *Environ. Health Perspect.* **123**, 1015–1021 (2015).
59. Lee, H. A., Park, S. H., Hong, Y. S., Ha, E. H. & Park, H. The effect of exposure to persistent organic pollutants on metabolic health among KOREAN children during a 1-year follow-up. *Int. J. Environ. Res. Public Health* **13**, E270 (2016).
60. Harari, R. et al. Neurobehavioral deficits and increased blood pressure in school-age children prenatally exposed to pesticides. *Environ. Health Perspect.* **118**, 890–896 (2010).
61. Suarez-Lopez, J. R., Jacobs, D. R. Jr, Himes, J. H. & Alexander, B. H. Acetylcholinesterase activity, cohabitation with floricultural workers, and blood pressure in Ecuadorian children. *Environ. Health Perspect.* **121**, 619–624 (2013).
62. Geiger, S. D., Xiao, J. & Shankar, A. No association between perfluoroalkyl chemicals and hypertension in children. *Integr. Blood Press Control* **7**, 1–7 (2014).
63. Toschke, A. M., Kohl, L., Mansmann, U. & von Kries, R. Meta-analysis of blood pressure tracking from childhood to adulthood and implications for the design of intervention trials. *Acta Paediatr.* **99**, 24–29 (2010).
64. Chen, X. & Wang, Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation* **117**, 3171–3180 (2008).
65. Chemla, D. et al. New formula for predicting mean pulmonary artery pressure using systolic pulmonary artery pressure. *Chest* **126**, 1313–1317 (2004).
66. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The Fourth Report on the Diagnosis, Evaluation, and Treatment of High Blood Pressure in Children and Adolescents. *Pediatrics* **114** (2 Suppl 4th Report), 555–576 (2004).
67. Update on the 1987 Task Force Report on High Blood Pressure in Children and Adolescents: a working group report from the National High Blood Pressure Education Program. National High Blood Pressure Education Program Working Group on Hypertension Control in Children and Adolescents. *Pediatrics* **98** (4 Pt 1), 649–658 (1996).