



OPEN Identification of redox activators for continuous reactivation of glyoxal oxidase from *Trametes versicolor* in a two-enzyme reaction cascade

Saadet Alpdagtas^{1,2}, Nina Jankowski², Vlada B. Urlacher² & Katja Koschorreck²✉

Glyoxal oxidases, belonging to the group of copper radical oxidases (CROs), oxidize aldehydes to carboxylic acids, while reducing O_2 to H_2O_2 . Their activity on furan derivatives like 5-hydroxymethylfurfural (HMF) makes these enzymes promising biocatalysts for the environmentally friendly synthesis of the bioplastics precursor 2,5-furandicarboxylic acid (FDCA). However, glyoxal oxidases suffer from inactivation, which requires the identification of suitable redox activators for efficient substrate conversion. Furthermore, only a few glyoxal oxidases have been expressed and characterized so far. Here, we report on a new glyoxal oxidase from *Trametes versicolor* (TvGLOX) that was expressed at high levels in *Pichia pastoris* (reclassified as *Komagataella phaffii*). TvGLOX was found to catalyze the oxidation of aldehyde groups in glyoxylic acid, methyl glyoxal, HMF, 2,5-diformylfuran (DFF) and 5-formyl-2-furancarboxylic acid (FFCA), but barely accepted alcohol groups as in 5-hydroxymethyl-2-furancarboxylic acid (HMFCFA), preventing formation of FDCA from HMF. Various redox activators were tested for TvGLOX reactivation during catalyzed reactions. Among them, a combination of horseradish peroxidase and its substrate 2,2'-azino-di-(3-ethylbenzthiazoline sulfonic acid) (ABTS) most efficiently reactivated TvGLOX. Through continuous reactivation of TvGLOX in a two-enzyme system employing a recombinant *Moesziomyces antarcticus* aryl-alcohol oxidase (MaAAO) almost complete conversion of 8 mM HMF to FDCA was achieved within 24 h.

Keywords Glyoxal oxidase, Copper radical oxidase, Redox activators, FDCA (2,5-furandicarboxylic acid)

Due to climate change and fossil fuel deficiency, the demand for renewable and sustainable equivalents instead of fossil-derived commodities is increasing day by day¹. In this context, waste plant biomass with its heterogenous composition has a promising potential for utilization in industrial biotechnology². Lignocellulose accounts for the majority of this plant biomass and thus represents a readily available, sustainable, and renewable source for providing bio-based chemicals³. The environmental production of value-added commodities or building blocks from biomass is quite attractive and has gained increasing attention over the last years. For instance, the synthesis of renewable building blocks for polymeric materials like bioplastics has been intensively studied. FDCA (2,5-furandicarboxylic acid) is a promising precursor for the sustainable synthesis of bioplastics. FDCA can be derived from FFCA (5-formyl-2-furancarboxylic acid), that is produced from HMF (5-hydroxymethylfurfural) either over HMFCFA (5-hydroxymethyl-2-furancarboxylic acid) or over DFF (2,5-diformylfuran) (Fig. 1). HMF is produced during the pretreatment of lignocellulosic biomass⁴. Several studies have dealt with the development of enzymatic routes for the environmentally friendly and sustainable production of FDCA from HMF^{5–10}. Fungal enzymes involved in lignin degradation have been thoroughly investigated for this purpose^{11–13}.

Several organisms, particularly white-rot fungi, are efficient degraders of lignocellulose and the fungal secretome comprises a variety of enzymes like laccases, peroxidases, and H_2O_2 -generating auxiliary enzymes, that contribute to lignin degradation¹⁴. Glyoxal oxidase (GLOX), discovered in secretomes of several fungi, is suggested to be involved in this process by fueling peroxide-depending ligninolytic peroxidases with H_2O_2 . GLOX belongs to the family of copper radical oxidases (CROs) and catalyzes the reduction of O_2 to H_2O_2 while oxidizing

¹Department of Biology, Van Yuzuncu Yil University, Van 65080, Turkey. ²Institute of Biochemistry, Heinrich Heine University Düsseldorf, Universitätsstraße 1, 40225 Düsseldorf, Germany. ✉email: Katja.Koschorreck@hhu.de




Figure 1. Reaction routes for HMF conversion into FDCA.

several aldehydes to carboxylic acids¹⁵. While other members of CROs like galactose oxidase (GalOx) have been extensively studied and some crystal structures of these enzymes are available, much less is known about GLOX enzymes. Similar to all CROs, these enzymes have a conserved active site with a mononuclear copper ion coordinated to an axial tyrosine, two histidines, and a cross-linked cysteine-tyrosyl radical cofactor^{11,16}. Notably, GLOX as well as GalOx is inactivated during purification and requires initial activation by a strong oxidizing agent^{17,18}. Inactivation is caused by the reduction of the active free-radical Cu(II)-state to the catalytically inactive non-radical Cu(I)-state (Fig. 2), as well as by high H₂O₂ concentrations. Activation of GLOX by regeneration of the oxidized radical form could be achieved by treating GLOX with high-redox potential inorganic oxidants like Na₂IrCl₆¹⁸ or other single oxidants generated by lignin peroxidase¹⁹. However, the activated enzyme is unstable with a half-life of 4 h for the radical form, and returns to the reduced inactive form¹⁸. Up to now, only GLOX from *Phanerochaete chrysosporium*^{20,21}, from *Ustilago maydis*²², from *Myceliophthora thermophila*³, and three GLOX isoenzymes from *Pycnoporus cinnabarinus*^{1,23} were characterized. Nevertheless, the usefulness of GLOX for the production of value-added commodities from biomass has been demonstrated. For instance, GLOX was reported to oxidize glycerol and produce glyceraldehyde and glyceric acid^{23,24}. In 2019, Daou et al. investigated the ability of three GLOX isoenzymes for the bioconversion of HMF to yield FDCA, when combined with an aryl-alcohol oxidase (AAO) to prevent HMFCa accumulation¹. In this setup, 16% FDCA was formed from HMF along with 84% FFCA. However, further optimization of this promising two-enzyme approach for the complete conversion of HMF to FDCA is still required.

In the present study, we produced a new glyoxal oxidase from *Trametes versicolor* (*Tv*GLOX) in *Pichia pastoris* and investigated this enzyme for the production of FDCA from HMF in a two-enzyme reaction cascade. A suitable redox activator for continuous GLOX reactivation in the course of the reaction was identified and applied in our setup, giving almost complete conversion of 8 mM HMF to FDCA within 24 h.

Results and discussion

*Tv*GLOX production and enzyme purification

*Tv*GLOX was identified by protein BLAST search, using GLOX from *P. chrysosporium* (UniProtKB: Q01772.1) as a query. The *tv*glox encoding DNA sequence with the native signal sequence for secretion (NCBI accession number XM_008037054) was codon optimized using the online tool JCat²⁵ and cloned into the pPICZA plasmid




Figure 2. Reaction cycle of GLOX and reactivation by ABTS and horseradish peroxidase. I, oxidative half-reaction, II, reductive half-reaction.

under the control of the *AOX1* promoter. The resulting plasmid was integrated into the *P. pastoris* X-33 genome by homologous recombination. Among 48 transformants, four *P. pastoris* colonies showing the most intensive green halo formation on methylglyoxal/HRP/ABTS-containing agar plates were selected for expression in shaking flasks. The transformant with the highest volumetric activity towards methylglyoxal of 0.23 U/ml was chosen for fed-batch cultivation in a 7.5 L bioreactor. After 9 days of fermentation, a volumetric activity of 19,000 U/l and a protein concentration of 1.6 g/l were obtained. Similar volumetric activity of 22.7 U/ml²¹ and protein concentration of 1–2 g/l¹⁵ were reported for expression of *P. chrysosporium* GLOX in *P. pastoris*.

TvGLOX was concentrated from the supernatant and purified by hydrophobic interaction chromatography. The purified enzyme displayed a specific activity towards methylglyoxal of 4.0 U/mg. During purification, TvGLOX changed its color from light green, attributed to the oxidized form of the enzyme, to purple color, which was observed for GLOX from *P. cinnabarinus* as well²³.

Purified TvGLOX had a molecular mass of around 80 kDa (Fig. S1), while the calculated molecular mass of native TvGLOX is 58 kDa. Four potential *N*-glycosylation sites were found in TvGLOX (Fig. S2). *N*-deglycosylation of TvGLOX shifted the band to around 58 kDa (Fig. S1), giving about 29% *N*-glycosylation of TvGLOX which is higher than ~18% *N*-glycosylation of GLOX from *P. chrysosporium*, heterologously expressed in *P. pastoris*²⁶.

Biochemical characterization of TvGLOX

TvGLOX showed the highest activity towards methylglyoxal at pH 6.5 (Fig. S3) which is similar to other GLOX enzymes^{1,3,21,23} and retained about 100% of its initial activity after 5 h incubation at 25 °C and pH 6.5 (Fig. S4). After 3 h incubation of TvGLOX between 40 °C and 60 °C, about 90% of its initial activity remained, whereas at 70 °C, it lost its activity immediately (Fig. S4). The thermostability of other GLOX enzymes like *Pci*GLOX1 and *Pci*GLOX2 at 40 °C and 50 °C was comparable to that of TvGLOX, while at 60 °C *Pci*GLOX1 and *Pci*GLOX2 retained only 50% of their initial activity after 2 h of incubation²³.

The substrate spectrum of TvGLOX was estimated in a coupled assay, including HRP and ABTS, to measure H₂O₂ generated by TvGLOX during substrate oxidation. A total of 15 compounds including HMF, HMFCa, DFF, FFCA, furfural, glycerol, formaldehyde, glyoxal, methylglyoxal, glyoxylic acid, D-galactose, D-glucose, benzyl alcohol, veratryl alcohol, and veratral were tested. As shown in Fig. 3, TvGLOX possessed the highest activity towards glyoxylic acid followed by methylglyoxal, glyoxal, formaldehyde and DFF. Activity towards glycerol, HMF, FFCA and furfural was also detected, while no activity was found towards D-glucose, D-galactose, benzyl alcohol, veratryl alcohol, veratral and HMFCa. These results confirm that TvGLOX is a typical glyoxal oxidase, that mainly accepts aldehyde groups and hardly oxidizes alcohol groups. In comparison, *Pci*GLOX1 also demonstrated the highest activity towards glyoxylic acid, whereas other GLOX enzymes such as *Pci*GLOX3 or GLOX from *P. chrysosporium* preferred methylglyoxal over glyoxylic acid^{1,19,23}. Homology modeling confirmed that the active site and overall structure of TvGLOX is quite similar to other GLOX enzymes such as GLOX from *P. chrysosporium* (Fig. 4). Like in other CROs, the active site of TvGLOX contains an axial tyrosine (Tyr399), two histidines (His400 and His493), and a cross-linked cysteine-tyrosyl radical cofactor formed by Cys93 and Tyr158.

Oxidation of HMF and derivatives thereof by TvGLOX

Although GLOX typically oxidizes aldehydes to the corresponding acids and thus converts HMF to HMFCa, this enzyme is also capable of oxidizing alcohols to aldehydes, like glycerol to glyceraldehyde. While *Pci*GLOX1, *Pci*GLOX2, and *Pci*GLOX3 have been reported to predominantly oxidize the aldehyde group of HMF and to produce HMFCa¹, *Mt*GLOx from *Myceliophthora thermophila* M77 has been shown to oxidize the alcohol group of HMF leading to DFF, but failed to further oxidize DFF³. We investigated the activity of TvGLOX towards




Figure 3. Substrate spectrum of TvGLOX (activity towards glyoxylic acid was set to 100%).




Figure 4. Structural model of TvGLOX and GLOX from *P. chrysosporium*. Homology models of both enzymes were built using GalOx (PDB: 2EIE) as template. Catalytic site of (a) TvGLOX and (b) GLOX from *P. chrysosporium*. The crosslink between the cysteine-tyrosine cofactor is missing due to the method of homology model building. Cartoon and surface representation of (c) TvGLOX and (d) GLOX from *P. chrysosporium*. The copper ion is depicted as a blue sphere.

HMF, DFF, HMFCA, and FFCA in more detail and determined the corresponding product profiles in these reactions (Table 1). With HMF, TvGLOX was mainly active on the aldehyde group and converted HMF to 88% HMFCA and about 12% FDCA after 24 h. Conversion of HMF by *Pci*GLOX2 and *Pci*GLOX3 reached only 39% and 41%, respectively after 24 h¹. In contrast to *Mt*GLOX, which failed to oxidize DFF³, TvGLOX showed the highest activity towards DFF among the investigated furan derivatives and completely converted DFF via FFCA to FDCA within 24 h.

With HMFCA no product formation was detected after 24 h, which is consistent with the preference of TvGLOX for the aldehyde group (as in HMF) and low acceptance of alcohol groups (as in HMFCA). According to this, the aldehyde group of FFCA was oxidized very well, leading to 100% FDCA after 24 h. As HMFCA was

Detected compounds	Initial substrate							
	HMF		DFF		HMFCA		FFCA	
	2 h	24 h	2 h	24 h	2 h	24 h	2 h	24 h
	Product distribution (%)							
HMF	57.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0
DFF	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HMFCA	42.6	88.1	0.0	0.0	100.0	100.0	0.0	0.0
FFCA	0.0	0.0	20.9	0.0	0.0	0.0	14.7	0.0
FDCA	0.0	11.9	79.1	100.0	0.0	0.0	85.3	100.0

Table 1. Conversion of furan derivatives catalyzed by TvGLOX.

not accepted at all, the formation of small amounts of FDCA in the conversion of HMF with TvGLOX might be explained by very slow oxidation of HMF to DFF (not observed after 2 h of reaction), followed by a fast conversion of DFF to FFCA and further to FDCA within 24 h of reaction. In contrast, with PciGLOX1-3 only a low conversion of HMFCFA to FFCA and FDCA within 24 h has been reported¹. The absence of oxidation products in the reaction of TvGLOX with HMFCFA may also be related to stability issues caused by this furan derivative. Therefore, we investigated the influence of the furan derivatives on the stability of TvGLOX. Only a slight decrease in activity was observed when TvGLOX was incubated for 24 h with HMFCFA or FFCA, while no loss in activity was observed with HMF and DFF (Fig. S5). Other GLOX enzymes are more sensitive to furan derivatives, e.g. all three GLOX enzymes from *P. cinnabarinus* almost completely lost their activity after 24 h incubation with HMFCFA¹. From our results, it can be concluded, that the lack of product formation in the reaction of TvGLOX with HMFCFA can be attributed to the inability of this enzyme to oxidize the alcohol group of HMFCFA.

Redox activators for TvGLOX

As known from the literature, GLOX easily undergoes inactivation by reduction of the active free-radical Cu(II)-state to the catalytically inactive non-radical Cu(I)-state requiring reactivation in the course of substrate oxidation reaction^{17,19,24}. High-redox potential inorganic oxidants like potassium octacyanomolybdate ($K_3Mo(CN)_8$), sodium hexachloroiridate (Na_2IrCl_6) or Mn^{3+} can reactivate GLOX^{18,24}. In ligninolytic cultures, GLOX from *P. chrysosporium* was fully activated in the presence of lignin peroxidase combined with its substrate veratryl alcohol¹⁹. In this system, lignin peroxidase consumed H_2O_2 produced by GLOX. HRP can also be used for GLOX activation when applied together with a suitable HRP substrate such as ABTS or veratryl alcohol^{18,27}. Although the activating effect has not been completely understood, it has been suggested that a single-electron oxidant generated by HRP or other peroxidases, is used to regenerate the oxidized radical form of GLOX²⁸. Recently, this hypothesis has been supported by the finding that ABTS cation radicals, generated through the oxidation by HRP in HRP/GLOX-coupled reactions, were in fact the species responsible for GLOX activation during methylglyoxal oxidation (Fig. 2)¹⁷. In order to find the most suitable redox activator for TvGLOX, we tested different HRP substrates—ABTS, 2,6-dimethoxyphenol, catechol, methyl syringate, syringaldazine, *p*-hydroquinone, guaiacol, veratryl alcohol and Mn^{2+} , respectively—as redox activators of TvGLOX during oxidation of FFCA (Table 2). Mn^{2+} is not a common HRP substrate, but Mn^{3+} -oxalate complexes were previously shown to strongly activate GLOX when HRP was included in the reaction, due to the HRP-catalyzed oxidation of Mn^{2+} to Mn^{3+} ²⁴.

In the absence of both, HRP and a redox activator, a slight formation of FDCA by TvGLOX was detected after 24 h, which may indicate that TvGLOX was not completely inactive after purification. MtGLOx from *M. thermophila* M77 was found to be fully active even without the addition of peroxidases or oxidizing agents³. Addition of commercial HRP Type II improved FDCA formation to a certain extent (18% FDCA with HRP vs. 5% FDCA w/o HRP). Whether TvGLOX was activated by any compounds contained in the commercially available HRP preparation or activated by the enzyme itself, as it was reported for GLOX from *P. chrysosporium*, which was moderately activated when incubated with lignin peroxidase¹⁹, remains elusive. The HRP/ABTS-system gave the best results with complete FFCA conversion within 24 h, followed by the HRP/methyl syringate system (94% FDCA). When FFCA conversion was analyzed after 2 h, it became apparent that HRP/ABTS overperformed HRP/methyl syringate in activating TvGLOX (85% FDCA vs. 62% FDCA, respectively). HRP/ Mn^{2+} was also able to activate TvGLOX, but only half the FFCA conversion was achieved compared to the reaction with ABTS. Veratryl alcohol only moderately activated TvGLOX as was also reported by Kersten et al.²⁷, and resulted only in 25% conversion of FFCA. Syringaldazine did not improve FFCA conversion compared to the reaction without any redox activator, whereas 2,6-dimethoxyphenol, guaiacol, *p*-hydroquinone and catechol reduced FFCA conversion. Inhibition of GLOX by guaiacol and catechol when peroxidase was added to the reaction has been described before²⁷. Consequently, the best HRP substrate, ABTS, with the strongest activating effect on TvGLOX was used in further experiments.

Entry	HRP	Redox activator	FDCA (%)
1	–	–	5
2	+	–	18
3	+	ABTS	100
4	+	Methyl syringate	94
5	+	Mn^{2+}	52
6	+	Veratryl alcohol	25
7	+	Syringaldazine	17
8	+	2,6-Dimethoxyphenol	10
9	+	Guaiacol	6
10	+	<i>p</i> -Hydroquinone	4
11	+	Catechol	3

Table 2. Formation of FDCA (%) from 2 mM FFCA by TvGLOX in the presence of HRP and different redox activators after 24 h.

Reaction cascade for the production of FDCA from HMF

In order to achieve the complete conversion of HMF to FDCA and avoid the accumulation of the dead-end product HMFCa, a cascade involving a second enzyme, the aryl-alcohol oxidase *MaAAO* from *Moesziomyces antarcticus*, was established (Fig. S6). In our previous work, *MaAAO* was found to convert HMF to FFCA within 24 h, but was not able to convert FFCA to FDCA²⁹. We proposed that by combining the activities and substrate preferences of *MaAAO* and *TvGLOX*, complete conversion of HMF to FDCA can be achieved. Several experimental setups were tested. *TvGLOX* was added after 2 h and 24 h of reaction with *MaAAO*, respectively, and in one setup both enzymes were added from the beginning of the reaction. When *TvGLOX* and *MaAAO* were added at the same time as the substrate HMF, complete conversion of 2 mM HMF to FDCA was achieved after 24 h (Fig. 5). In a similar approach, Daou et al. combined *UmAAO* from *Ustilago maydis* and *PciGLOX3* from *P. cinnabarinus* in a cascade for HMF conversion to FDCA, but detected only 16% FDCA along with 84% FFCA¹. In that study, catalase was also added to decompose high concentrations of H₂O₂ but conversion could be only marginally improved. Comparison with our results allows us to suggest that continuous reactivation of *GLOX* has a strong effect on the cascade outcome and *GLOX* inactivation by H₂O₂ only plays a marginal role.

In the next step, we increased the HMF concentration while maintaining the enzyme load, and observed complete conversion of up to 6 mM HMF to FDCA. With 8 mM HMF 7.8 mM FDCA and 0.2 mM FFCA were measured and with 10 mM HMF 7.8 mM FDCA and 2.2 mM FFCA (Figs. 5 and S7). For comparison, in a three-enzyme system consisting of a galactose oxidase, an unspecific peroxxygenase, and an aryl-alcohol oxidase, 7.9 mM FDCA was achieved after 24 h starting from 9.7 mM HMF⁷. Our results show that *TvGLOX* is a promising biocatalyst for the enzymatic synthesis of bioplastics precursors, and the continuous reactivation of this Cu-radical oxidase represents a critical factor during process optimization.

Conclusion

In this study, a new glyoxal oxidase from *T. versicolor*—*TvGLOX*—was heterologously expressed at high levels in *P. pastoris* and characterized regarding its substrate spectrum, stability and reactivation by different redox activators. *TvGLOX* was most efficiently reactivated by ABTS in the presence of horseradish peroxidase (HRP) and preferred aldehydes over alcohols, leading to HMFCa formation from HMF. HMFCa was not accepted by this enzyme preventing further oxidation to yield FDCA. To achieve complete conversion to FDCA, a biocatalytic system consisting of *MaAAO* and *TvGLOX*, the latter one continuously reactivated through HRP and ABTS, was developed. With this system, up to 8 mM HMF was almost completely converted to FDCA within 24 h, demonstrating that *TvGLOX* is a promising biocatalyst for biotechnological applications.

Methods and material

Construction of recombinant *Pichia pastoris* strain

TvGLOX from *Trametes versicolor* FP-101664 SS1 (NCBI accession number XM_008037054) was identified by protein BLAST search using *GLOX* from *T. cinnabarina* (GenBank: ANJ20632.1) as query (88.91% identity, E-value 0.0, query coverage 100%). The *tvglx* encoding sequence with its native signal sequence for secretion was synthesized by BioCat GmbH (Heidelberg, Germany) after codon optimization for expression in *Saccharomyces cerevisiae* using the online tool JCat²⁵. The gene was cloned into the *Pichia pastoris* expression vector pPICZA between *EcoRI* and *NotI* restriction sites (pPICZA_ *tvglx*). Chemically competent *E. coli* DH5a cells were used for plasmid propagation. Transformants were selected on low salt lysogeny broth agar plates including 25 µg/ml Zeocin. Plasmid isolation was performed by using the ZR Plasmid Miniprep Kit (Zymo Research, USA). Electrocompetent *P. pastoris* X-33 cells were transformed by electroporation with 10 µg pPICZA_ *tvglx* plasmid after linearization with *MssI*. Positive transformants were selected on yeast extract peptone dextrose sorbitol (YPDS) agar plates supplemented with 100 µg/ml Zeocin after incubation at 30 °C for 3 days.




Figure 5. Dependence of FDCA production in the *TvGLOX/MaAAO* reaction system on increasing HMF concentrations.

Heterologous expression of TvGLOX in shaking flask and bioreactor

P. pastoris transformants with TvGLOX activity were selected on buffered minimal methanol (BMM) screening agar plates (1.34% yeast nitrogen base, 100 mM potassium phosphate buffer pH 6.0, $4 \cdot 10^{-5}$ biotin, 0.5% methanol, 2% agar, 0.5 mM 2,2'-azino-di-(3-ethylbenzthiazoline sulfonic acid (ABTS)), 0.006 mg/ml HRP, 2 mM methylglyoxal). Colonies with the strongest green halo formation after 72 h incubation at 30 °C were selected for cultivation in shaking flasks. For this purpose, a colony from the agar plate was grown in 10 ml BMGY (buffered complex glycerol) medium overnight at 30 °C and 200 rpm. The overnight-culture was used for the inoculation of 200 ml BMMY (buffered complex methanol) medium to an OD₆₀₀ of 1 and incubated for 3–4 days at 25 °C and 200 rpm. Every 24 h, the cultures were fed with 0.5% (v/v) methanol. The OD₆₀₀ value and volumetric activity of the supernatant were measured daily using methylglyoxal as substrate.

The most active *P. pastoris* transformant was used for fed-batch fermentation. Fed-batch fermentation was performed in a 7.5 L bioreactor (Infors, Switzerland) as previously described³⁰. The temperature was set to 25 °C after induction of TvGLOX expression and the fermentation proceeded for 9 days with daily sampling to follow OD₆₀₀ and volumetric activity towards methylglyoxal.

Enzyme purification

P. pastoris cells were separated from fermentation broth by centrifugation at 4 °C and $11,325 \times g$ for 15 min, and the collected supernatant containing secreted TvGLOX was concentrated and rebuffed with 25 mM sodium phosphate pH 6.0 by tangential flow filtration (TFE, 10 kDa molecular cut-off; Pall, Port Washington, USA). TvGLOX was purified via hydrophobic interaction chromatography (HIC) on an ÄKTA purifier FPLC system (GE Healthcare, USA). At first, 10 ml concentrated supernatant was incubated in 25 mM sodium phosphate buffer pH 6.0 containing 1.5 M ammonium sulfate at 10 °C overnight. Then, the sample was centrifuged at 4 °C and $18,000 \times g$ for 30 min, filtered and applied on the column with butyl sepharose HP medium. Elution was performed by decreasing ammonium sulfate concentration by adding 25 mM sodium phosphate buffer pH 6.0. Fractions with activity towards methylglyoxal were concentrated and desalted via an ultrafiltration membrane filter (Vivaspin Turbo 15, 10 kDa molecular cut-off). The purified TvGLOX enzyme was kept at 4 °C until use.

Biochemical characterization

The activity of TvGLOX was routinely measured in 96 well microtiter plates in a coupled assay using horseradish peroxidase (HRP, Type II; purchased from Sigma-Aldrich (Schnellendorf, Germany)) and ABTS for initial activation of GLOX and monitoring of substrate oxidation through H₂O₂ formation. Reaction mixtures contained 20 µl TvGLOX solution, 20 µl 100 mM methylglyoxal, 20 µl 0.06 mg/ml HRP, 20 µl 5 mM ABTS, and 120 µl 50 mM sodium phosphate buffer pH 6.5, including 0.5 µM H₂O₂. ABTS oxidation was followed at 420 nm using an Infinite M200 Pro plate reader (Tecan, Switzerland).

Protein concentration was measured using the Bradford method³¹. Deglycosylation of purified TvGLOX (20 µg) was carried out under native and denaturing conditions (for up to 24 h) by using Peptide N-Glycosidase F (PNGase F, New England Biolabs, Germany) according to the manufacturer's instructions. Purified and deglycosylated protein was visualized by SDS-PAGE³². Potential N-glycosylation sites were predicted by using the NetNGlyc 1.0 server prediction tool³³.

pH optimum of purified TvGLOX was measured in 40 mM Britton-Robinson buffer at several pH values between pH 2.0–11.5 by using methylglyoxal as substrate as described above. The stability of TvGLOX at pH 6.5 was investigated by incubating the enzyme in 50 mM sodium phosphate buffer pH 6.5 at 25 °C for 5 h. Thermal stability was determined by incubating TvGLOX in 50 mM sodium phosphate buffer pH 6.5 at different temperatures (40 °C, 50 °C, 60 °C, and 70 °C) for 3 h. After incubation, samples were taken, incubated on ice and the remaining activity was measured at room temperature using methylglyoxal as substrate.

The enzyme activity towards 15 compounds (HMF, HMFCFA, DFF, FFCA, furfural, glycerol, formaldehyde, glyoxal, methylglyoxal, glyoxylic acid, D-galactose, D-glucose, benzyl alcohol, veratryl alcohol, and veratral) was estimated at a substrate concentration of 10 mM in a coupled assay as described above.

The enzyme stability towards furan derivatives was assessed by incubating TvGLOX in 50 mM sodium phosphate buffer pH 6.5 with 2 mM substrate (HMF, DFF, HMFCFA, and FFCA, respectively). Samples were taken after 4 h, 24 h, and 48 h incubation, and the remaining activity towards methylglyoxal was measured as described above.

Oxidation of furan derivatives

Reactions were performed in a total volume of 200 µl in 50 mM sodium phosphate buffer pH 6.5 with 2 mM substrate (HMF, DFF, HMFCFA and FFCA, respectively), 2 µM TvGLOX, 0.5 µM H₂O₂, 0.1 mg/ml HRP and 0.5 mM ABTS. Reactions were incubated at 25 °C and 500 rpm for up to 24 h. Fifty µl samples were taken after 15 min, 2 h and 24 h, acidified with 10 µl 6 M HCl and 2-furoic acid (20 mM) was added as an internal standard. Methyl *tert*-butyl ether (MTBE) was used for sample extraction, each sample was dried over MgSO₄, evaporated to dryness and resuspended in 50 µl of *N,O*-bis (trimethylsilyl) trifluoroacetamide (BSTFA), and incubated at 60 °C for 30 min before GC–MS analysis.

Investigation of redox activators

Reactions were performed in a total volume of 200 µl in 50 mM sodium phosphate buffer pH 6.5 with 2 mM FFCA, 2 µM TvGLOX, 0.5 µM H₂O₂, 0.1 mg/ml HRP and 2 mM redox activator (2,6-dimethoxyphenol, catechol, methyl syringate, syringaldazine, *p*-hydroquinone, guaiacol, veratryl alcohol and MnSO₄, respectively). Reactions with HRP and Mn²⁺ were conducted in 50 mM sodium malonate buffer pH 6.5. Reactions were incubated

at 25 °C and 500 rpm for up to 24 h. Fifty µl samples were taken in the course of the reaction and prepared for analysis via GC–MS measurements as described above.

Optimization of HMF conversion

For optimization of the reaction cascade *Moesziomyces antarcticus* aryl alcohol oxidase (MaAAO) was used²⁹. Reactions were performed in a total volume of 200 µl in 50 mM sodium phosphate buffer pH 6.5 containing 2–10 mM HMF, 0.5 µM H₂O₂, 2 µM MaAAO, 0.1 mg/ml HRP, and 0.5 mM ABTS. All reactions were incubated at 25 °C, 500 rpm for up to 24 h. Samples were taken in the course of the reaction and prepared for analysis via GC–MS measurements as described above.

GC–MS analysis

Oxidation of furan derivatives was analyzed using a GC–MS–QP-2010 (Shimadzu, Japan) on an FS-Supreme-5 ms column (CS Chromatographie Service GmbH, Germany). The injection, interface, and ion source temperatures were set to 250 °C, 285 °C, and 200 °C, respectively. The temperature for the column was adjusted to 110 °C, maintained for 2 min, then increased to 300 °C gradually with 20 °C/min. Substrates and products were identified by comparing the measured mass spectra with authentic standards. Substrate conversion and product formation were calculated from substrate depletion (control set to 100%), using 2-furoic acid as the internal standard and from external calibration curves of furan derivatives (0–2 mM) and 2-furoic acid as internal standard.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Received: 20 December 2023; Accepted: 6 March 2024

Published online: 11 March 2024

References

- Daou, M. *et al.* *Pycnoporus cinnabarinus* glyoxal oxidases display differential catalytic efficiencies on 5-hydroxymethylfurfural and its oxidized derivatives. *Fungal. Biol. Biotechnol.* **6**, 4 (2019).
- Daou, M. & Faulds, C. B. Glyoxal oxidases: Their nature and properties. *World J. Microbiol. Biotechnol.* **33**, 87 (2017).
- Kadowaki, M. A. S. *et al.* Characterization of a new glyoxal oxidase from the thermophilic fungus *Myceliophthora thermophila* M77: Hydrogen peroxide production retained in 5-hydroxymethylfurfural oxidation. *Catalysts* **8**, 476 (2018).
- Agblevor, F. A. & Jahromi, H. Aqueous-phase synthesis of hydrocarbons from furfural reactions with low-molecular-weight biomass oxygenates. *Energy Fuel.* **32**, 8552–8562 (2018).
- Sayed, M. *et al.* Oxidation of 5-hydroxymethylfurfural with a novel aryl alcohol oxidase from *Mycobacterium* sp. MS1601. *Microb. Biotechnol.* **15**, 2176–2190 (2022).
- Vinambres, M., Espada, M., Martinez, A. T. & Serrano, A. Screening and evaluation of new hydroxymethylfurfural oxidases for furandicarboxylic acid production. *Appl. Environ. Microbiol.* **86**, e00842–00820 (2020).
- Karich, A., Kleeborg, S. B., Ullrich, R. & Hofrichter, M. Enzymatic preparation of 2,5-furandicarboxylic acid (FDCA)-a substitute of terephthalic acid-by the joined action of three fungal enzymes. *Microorganisms* **6**, 12 (2018).
- Dijkman, W. P., Groothuis, D. E. & Fraaije, M. W. Enzyme-catalyzed oxidation of 5-hydroxymethylfurfural to furan-2,5-dicarboxylic acid. *Angew. Chem. Int. Ed.* **53**, 6515–6518 (2014).
- Milic, M., Byström, E., de María, P. D. & Kara, S. Enzymatic cascade for the synthesis of 2,5-furandicarboxylic acid in biphasic and microaqueous conditions: “Media-agnostic” biocatalysts for biorefineries. *ChemSuschem.* <https://doi.org/10.1002/cssc.202102704> (2022).
- Serrano, A. *et al.* Complete oxidation of hydroxymethylfurfural to furandicarboxylic acid by aryl-alcohol oxidase. *Biotechnol. Biofuels* **12**, 217 (2019).
- Mathieu, Y. *et al.* Discovery of a fungal copper radical oxidase with high catalytic efficiency toward 5-hydroxymethylfurfural and benzyl alcohols for bioprocessing. *ACS Catal.* **10**, 3042–3058 (2020).
- Dijkman, W. P. & Fraaije, M. W. Discovery and characterization of a 5-hydroxymethylfurfural oxidase from *Methylovorus* sp. strain MP688. *Appl. Environ. Microbiol.* **80**, 1082–1090 (2014).
- Carro, J. *et al.* 5-hydroxymethylfurfural conversion by fungal aryl-alcohol oxidase and unspecific peroxygenase. *FEBS J.* **282**, 3218–3229 (2015).
- Dashtban, M., Schraft, H., Syed, T. A. & Qin, W. Fungal biodegradation and enzymatic modification of lignin. *Int. J. Biochem. Mol. Biol.* **1**, 36–50 (2010).
- Whittaker, M. M., Kersten, P. J., Cullen, D. & Whittaker, J. W. Identification of catalytic residues in glyoxal oxidase by targeted mutagenesis. *J. Biol. Chem.* **274**, 36226–36232 (1999).
- Whittaker, J. W. Galactose oxidase. *Adv. Protein Chem.* **60**, 1–49 (2002).
- Wohlschlager, L., Kracher, D., Scheiblbrandner, S., Csarman, F. & Ludwig, R. Spectroelectrochemical investigation of the glyoxal oxidase activation mechanism. *Bioelectrochemistry* **141**, 107845 (2021).
- Whittaker, M. M. *et al.* Glyoxal oxidase from *Phanerochaete chrysosporium* is a new radical-copper oxidase. *J. Biol. Chem.* **271**, 681–687 (1996).
- Kersten, P. J. Glyoxal oxidase of *Phanerochaete chrysosporium*: Its characterization and activation by lignin peroxidase. *Proc. Natl. Acad. Sci. USA.* **87**, 2936–2940 (1990).
- Kersten, P. J. & Kirk, T. K. Involvement of a new enzyme, glyoxal oxidase, in extracellular H₂O₂ production by *Phanerochaete chrysosporium*. *J. Bacteriol.* **169**, 2195–2201 (1987).
- Wohlschlager, L., Csarman, F., Zrilic, M., Seiboth, B. & Ludwig, R. Comparative characterization of glyoxal oxidase from *Phanerochaete chrysosporium* expressed at high levels in *Pichia pastoris* and *Trichoderma reesei*. *Enzyme Microb. Technol.* **145**, 109748 (2021).
- Leuthner, B. *et al.* A H₂O₂-producing glyoxal oxidase is required for filamentous growth and pathogenicity in *Ustilago maydis*. *Mol. Genet. Genomics* **272**, 639–650 (2005).
- Daou, M., Piumi, F., Cullen, D., Record, E. & Faulds, C. B. Heterologous production and characterization of two glyoxal oxidases from *Pycnoporus cinnabarinus*. *Appl. Environ. Microbiol.* **82**, 4867–4875 (2016).
- Roncal, T., Munoz, C., Lorenzo, L., Maestro, B. & de Guerenú, M. D. D. Two-step oxidation of glycerol to glyceric acid catalyzed by the *Phanerochaete chrysosporium* glyoxal oxidase. *Enzyme Microb. Technol.* **50**, 143–150 (2012).

25. Grote, A. *et al.* JCat: A novel tool to adapt codon usage of a target gene to its potential expression host. *Nucleic Acids Res.* **33**, 526–531 (2005).
26. Son, Y. L., Kim, H. Y., Thiyagarajan, S., Xu, J. J. & Park, S. M. Heterologous expression of *Phanerochaete chrysosporium* glyoxal oxidase and its application for the coupled reaction with manganese peroxidase to decolorize malachite green. *Mycobiology* **40**, 258–262 (2012).
27. Kurek, B. & Kersten, P. J. Physiological regulation of glyoxal oxidase from *Phanerochaete chrysosporium* by peroxidase systems. *Enzyme Microb. Technol.* **17**, 751–756 (1995).
28. Pedersen, A. T. *et al.* Process requirements of galactose oxidase catalyzed oxidation of alcohols. *Org. Process Res. Dev.* **19**, 1580–1589 (2015).
29. Lappe, A., Jankowski, N., Albrecht, A. & Koschorreck, K. Characterization of a thermotolerant aryl-alcohol oxidase from *Moesziomyces antarcticus* oxidizing 5-hydroxymethyl-2-furancarboxylic acid. *Appl. Microbiol. Biotechnol.* **105**, 8313–8327 (2021).
30. Jankowski, N., Koschorreck, K. & Urlacher, V. B. High-level expression of aryl-alcohol oxidase 2 from *Pleurotus eryngii* in *Pichia pastoris* for production of fragrances and bioactive precursors. *Appl. Microbiol. Biotechnol.* **104**, 9205–9218 (2020).
31. Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254 (1976).
32. Laemmli, U. K. Cleavage of structural proteins during assembly of head of bacteriophage-T4. *Nature* **227**, 680–685 (1970).
33. Kukuruzinska, M. A., Bergh, M. L. E. & Jackson, B. J. Protein glycosylation in yeast. *Annu. Rev. Biochem.* **56**, 915–944 (1987).

Acknowledgements

The authors wish to thank the Bioeconomy Science Center (BioSC, Germany) through the Ministry of Innovation, Science and Research of the German State of North Rhine-Westphalia within the framework of the NRW-Strategieprojekt BioSC (No. 313/323-400-00213) for financial support.

Author contributions

S.A.: investigation, formal analysis, data collection, conceptualization, validation, visualization, writing—original draft; writing—review and editing. N.J.: conceptualization, investigation. V.B.U.: project administration, supervision, conceptualization, writing—original draft; writing—review and editing. K.K.: conceptualization, supervision, investigation, data collection, validation, writing—original draft; writing—review and editing.

Funding

Open Access funding enabled and organized by Projekt DEAL. Funding of this project was provided by the Scientific and Technological Research Council of Turkey (TUBITAK) 2219-International Research Fellowship Program for Saadet Alpdağtaş.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-024-56429-z>.

Correspondence and requests for materials should be addressed to K.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024